

## **EFFECT OF STOCKING DENSITY AND FEEDING RATE ON PRODUCTION OF EARTHEN PONDS CULTURED WITH NILE TILAPIA (*Oreochromis niloticus*).**

**Ramdan, M.EL-SH.<sup>1</sup>; N.F. Abdel-Hakim<sup>2</sup>; M.M.S. Lachine<sup>3</sup> ; D.E.EL-Azab<sup>1</sup> and A.M. Zaghoul<sup>1</sup>**



<sup>1</sup> Anim. Prod. Dept., Fish production, Faculty of Agriculture, Al-Azhar University, Cairo.

<sup>2</sup> Animal Production Department, Animal Nutrition, Faculty of Agriculture, Al-Azhar University, Cairo.

<sup>3</sup> Animal Production Department, Animal Care, Faculty of Agriculture, Al-Azhar University, Cairo.

### **ABSTRACT**

The study was conducted over a 98 days period in order to study effects of stocking density and feeding rate on production of earthen ponds cultured with Nile tilapia (*Oreochromis niloticus*). Each hapa measuring 2\*4\*1m was suspended in an earthen pond (4000m<sup>2</sup>). There were 6 treatments, each consisting of three replicates, stocked with fish of mean individual initial bodyweight 19g. A total number of 32, 64 and 128 Nile tilapia were randomly distributed into 6 treatment groups, representing 3 stocking rates, and were fed daily at rates (zero, 3, and 4%) of fish live bodyweight. Fish were fed a balanced diet of 28.5% crude protein along the period of the experiment. The treatments were stocking densities (4, 8, and 16 fish/m<sup>3</sup>) and feeding rates (zero, 3, and 4%). The results indicate that the best final bodyweight, feed utilization, physiological status, and chemical composition was obtained by group T7 (stocking density 4fish/m<sup>3</sup> and feeding rate 4%).

**Keywords:** Nile tilapia, stocking density, feeding rate, earthen pond, growth performance, chemical composition.

### **INTRODUCTION**

Tilapias have a high reproduction capacity, fast growth, extensive feeding, and high resistance to disease; therefore, they have been recommended by the FAO to be a good culture species. There is a great need to increase fish production via aquaculture in Egypt to fill the gap in animal protein in the country. In 2006, fish production from aquaculture activities contributed by 61.28% of total fish production which amounted 970.913 metric tons. GAFRD (2006) stated that the full utilization of space for maximum fish production through intensive culture can improve the profitability of the fish farm. Fish intensification by increasing stocking density is also found suitable to overcome the problem of land shortage (Chakraborty and Banerjee, 2010). However, Chang (1988) reported that fish stocking density and feeding rates are important factors used in aquaculture as it can affect natural food availability, the efficient utilization of food resource and total fish yield in ponds. The objective of the present study was to determine the effect of different stocking densities and feeding rates on growth

performance, feed utilization, carcass composition and blood analysis of Nile tilapia reared in floating hapas in an earthen pond.

## MATERIALS AND METHODS

The present work was conducted in a fish farm in Kafr El-Sheikh governorate during season 2013 in order to evaluate daily feeding rates (0, 3, and 4%) and stocking densities (4, 8, and 16 fish /m<sup>3</sup>) on growth performance and chemical composition under Egyptian conditions.

### **Fish culture system:**

A total number of 1344 Nile tilapia fish (*Oreochromis niloticus*) with an average bodyweight of 19 g were obtained from (a private fish farm at Metobese area, belonging to Kafr El-Sheikh governorate, Egypt. Each size of fishes was randomly distributed among 18 hapa. Each hapa measuring 2\*4\*1m as width, length, and depth, respectively. Hapas were suspended in an earthen pond of 4200 m<sup>2</sup>. The stocking densities were 4, 8 and 16 fish/m<sup>3</sup>). Thirteen fish were kept frozen at -20 C<sup>0</sup> for chemical analysis at the beginning of the experiment. The pond was supplied with fresh water from Metobese area. The water exchange rate was 15% of total pond's water volume/day. The experimental period was 98 days (from 1/6 to 30/8/ 2014). Water pH was measured using pH meter (model 68 Engineered system and Designs). Water dissolved oxygen and temperature were measured p.m. by oxygen meter (WPA 20 scientific instrument).

### **Experimental diets:**

Fingerlings were fed a commercial diet (Go-Traid, Industrial Area, 6<sup>th</sup> of october) containing 28.5% crude protein and 4000 Kcal/Kg at feeding rates of 3%, and 4% of fresh biomass in each hapa (six days per week). Fish were fed two times daily at 8 am and 2 pm with feed amounts adjusted at approximately 14 day-intervals in response to the actual weight. Ingredients and composition of the experimental diet are presented in Table 1.

**Table 1: Ingredients and chemical analysis (%) of the commercial diet.**

Ingredients	Commercial diet
Fish meal	14.0
Soybean meal	40.0
Yellow corn	20.0
Wheat bran	11.0
Vegetable oil	4.0
Vitamins & Minerals premix	0.6
Rice bran	10.0
Limestone	0.4
Total	100
Chemical analysis	
Dry matter (DM)	90
Crude protein (CP)	28.5
Ether extract (EE)	6.8
Crude fiber (CF)	6.2
Ash	8.9
Nitrogen-free extract (NFE)*	49.6
GE (kcal/kg)**	3755
DG (kcal/kg)	2816
ME (kcal/kg)	3100
P/E (mg CP/Kcal GE)	99.7

\* Nitrogen. Free extract  $NFE=100(CP\%+EE\%+CF\%+ash\%)$

\*\*GE= Gross energy was calculated by multiply the Factors 4.1, 6.5 and 9.44 Kilo GE/g DM for carbohydrate, protein and Fat, respectively(Jobling,1983).

**Growth parameters:**

Average total gain (ATG), average daily gain (ADG), specific growth rate (SGR), feed conversion ratio (FCR), protein efficiency ratio (PER), protein productive value (PPV) and survival rate (SR) were calculated according to the following equations:

$$ATG (g/fish) = [Average\ final\ weigh (g) - Average\ initial\ weigh (g)].$$

$$ADG (g/fish/ day) = ATG / experimental\ period (d).$$

$$SGR (\%/day) = [Ln\ final\ body\ weight - Ln\ initial\ body\ weight * 100 / experimental\ period (d)].$$

FCR= feed intake (g)/live weight gain.

PER = Live weight gain (g)/protein intake (g).

PPV (%) =  $100 [final\ fish\ body\ protein (g) - initial\ fish\ body\ protein (g) / crude\ protein\ intake (g)]$ .

SR (%) =  $100 [total\ No\ of\ fish\ at\ the\ end\ of\ the\ experimental\ period / total\ No\ of\ fish\ at\ the\ start\ of\ the\ experiment]$ .

**Proximate analysis:**

Dry matter, crude protein, ether extract, (crude fiber), and ash contents of the commercial diet and whole body of fish at the beginning and at the end of the experimental were preformed according to A.O.A.C (1990).

**Blood parameters determination:**

At the end of the experimental period, blood samples from the different groups were taken from the caudal vein for analysis. Adequate amounts of whole blood in small plastic vials containing heparin were used. Blood plasma (after the centrifugation of the whole blood samples) was sent immediately to a private human clinical laboratory for determination of the biochemical parameters (using commercial kits obtained from Diamond Diagnostic Company, Egypt) including total proteins (TP) after the method of Henry (1964) and albumin according to the method of Kohin (1958). Whereas globulin was calculated by mathematical subtraction of albumin value from total proteins (Kohin 1958). Alanine aminotransferase (ALT) and aspartate aminotransferase (AST) activities were assayed according to the method of Reitman and Frankel (1957) using commercial kits purchased from Randox Company. Glucose was determined calorimetrically according to Trinder (1969). Cortisol levels were determined by an electrochemiluminometric assay using the Elecsys and Cobas e 411 Immunoassay Analyzer (Roche Diagnostics, Indianapolis, IN, USA). Other biochemical parameters were determined using commercial kits too.

**Statistical analysis:**

Analysis of variance (ANOVA) was used for analyzing collected data with the aid of statistical analysis system (SAS, 2006), Duncan's (1955) test was used to assess the significance between different stocking densities and feeding regimes.

## RESULTS AND DISCUSSION

**Chemical composition of the experiment of diets:**

Chemical composition and calculated energy of different diets are presented in Table 1. The CP content was between 28 to 28.5% on DM basis. Such level was within the range suggested by NRC (1993). The calculated gross energy (GE) was 375.5 Kcal/100g.

**Quality Parameters of rearing water:**

All tested water quality criteria (Table 2) were suitable for rearing Nile tilapia fingerlings as cited by Abdel-Hakim *et al.* (2002). Since water temperature ranged between 24.9 and 26.5°C, pH values 6.8 and 6.9, ammonia ranged between 0.09 and 0.11 mg/l, nitrate ranged between 0.23 and 0.26 mg/l, and nitrite ranged between 0.15 and 0.26. Also, Abdelhamid *et al.* (2002) suggested that these values are suitable for rearing Nile tilapia.

**Table 2: Effect of stoking density and feeding rate on water quality of Nile tilapia (*Oreochromis niloticus*).**

Stocking density	Feeding rate	Temperature (C°)	Dissolved Oxygen (mg l <sup>-1</sup> )	PH	Ammonia (mg l <sup>-1</sup> )	Nitrate (mg l <sup>-1</sup> )	Nitrite (mg l <sup>-1</sup> )
4/m <sup>3</sup>	3%	24.9± 0.1	3.8± 0.22	6.9± 0.103	0.11± 0.017	0.242± 0.82	0.20± 0.038
	4%	25.5± 0.0	4.1± 0.22	6.8± 0.103	0.09± 0.017	0.234± 0.82	0.26± 0.038
		25.1± 0.0	4.0 ± 0.15	6.9± 0.07	0.10± 0.012	0.240± 0.58	0.23± 0.027
8/m <sup>3</sup>	3%	26.1± 0.1	3.8± 0.22	6.8± 0.103	0.09± 0.017	0.250± 0.82	0.25± 0.038
	4%	25.5± 0.02	4.0± 0.22	6.9± 0.103	0.09± 0.017	0.250± 0.82	0.26± 0.038
		26.2± 0.1	3.9± 0.15	6.80± 0.07	0.09± 0.012	0.250± 0.58	0.20± 0.027
16/m <sup>3</sup>	3%	25.5 ± 0.6	3.8± 0.22	6.9± 0.103	0.10± 0.017	0.260± 0.82	0.26± 0.038
	4%	26.5± 0.1	4.1± 0.22	6.8± 0.103	0.09± 0.017	0.234± 0.82	0.26± 0.038
		26.4± 0.8	3.9± 0.15	6.9± 0.07	0.1± 00.012	0.250± 0.58	0.26± 0.027

**Growth performance and survival rate:**

At the end of the experimental period, both groups T6 and T1 (4 fish/m<sup>3</sup>, and feeding rate 4% and 4 fish/ m<sup>3</sup>, and feeding rate zero). followed by T4 (4 fish/m<sup>3</sup> and feeding rate 3%) gave significant increases in total weight gain (TWG), average daily gain (ADG), specific growth rate (SGR), and survival rate (SR). There were significant (P≤ 0.05) differences among various groups of fish concerning final body weight, daily gain, specific growth rate, and survival rate of the experimented fish, being the best values in favor of T6 and T1, which seem even more better than the T2, T3, T5, T6, T8, and T9. These results are demonstrated in Table 3. These results are supported by those of Amar (2009) and Awad 92015). Abdel-Hakim *et al.* (1995) who reported that growth performance of Nile tilapia cultured in earthen ponds decreased as the stoking density increased from 3000 to 4500 or 6000 fish/feddan. In this connection, Pagan (1970), Suwanasart (1971), and Cohche (1976) showed that growth rate of fish in general decreased with increasing stocking rate. Furthermore, Moav *et al.* (1977) and Barlarin (1979) showed that body weight and growth rate of fish stocked at higher rates showed depression, may be attributed to crowding, social interaction and aggression. Also, results in Table 3 are in agreement with the finding of Abdel-Hakim *et al.* (2008) who reported that lower stocking densities (16000 fish/feddan) resulted in significantly higher final weights and lengths of fish compared with the 19000 and 22400 fish/feddan. Furthermore, Zannatul *et al.* (2014) indicated in general that the highest initial weights influences the bodyweight during the growth period, it gives the highest final weight and highest SGR. Results of Abdel- Hakim *et al.* (2001) showed that increasing

tilapia stocking density from 50 to 100 or 150 fish /m<sup>3</sup> of tank water decreased significantly final body weight and length of fish.

**Table 3: Effect of stoking density and feeding rate on growth performance of Nile tilapia (*Oreochromis niloticus*).**

Feeding rate (FR)	Overall regardless of feeding rate			FR-Level 3%			FR-Level 4%		
	T1	T2	T3	T4	T5	T6	T7	T8	T9
Stocking density	4 f/m <sup>3</sup>	8 f/m <sup>3</sup>	16 f/m <sup>3</sup>	4 f/m <sup>3</sup>	8 f/m <sup>3</sup>	16 f/m <sup>3</sup>	4 f/m <sup>3</sup>	8 f/m <sup>3</sup>	16 f/m <sup>3</sup>
Initial body weight (g)	19.61±0.07a	19.58±0.03a	19.62±0.07a	19.58±0.14a	19.63±0.07a	19.32±0.09a	19.63±0.05a	19.71±0.03a	19.53±0.09a
Final body weight (g)	173.58±4.68a	139.56±2.79b	103.25±1.46c	169.93±8.02a	139.00±5.29b	102.59±1.72c	177.23±5.65a	140.12±3.26b	103.92±2.70c
WG (g)	153.97±4.62a	119.84±2.82b	83.83±1.41c	150.35±7.88a	119.27±5.36b	83.27±1.63c	157.60±5.61a	120.41±3.26b	84.39±2.63c
FI (g)	189.21±38.80c	164.00±63.46b	124.24±75.39a	158.88±23.17c	137.59±6.85b	108.67±24.27a	219.53±21.24c	190.41±15.56b	139.81±15.6a
SGR (%/d)	2.42±0.17a	2.17±0.17b	1.85±0.17c	2.34±0.24a	2.17±0.24b	1.85±0.24c	2.44±0.24a	2.18±0.24b	1.85±0.24c
SR (%)	99.2	98.3	94.5	99.3	98.3	98.0	99.0	98.3	91.0

Means bearing different small letters are significant at (P<0.05).

T1= 4 fish/m<sup>3</sup> and feeding rate zero, T2= 8 fish/m<sup>3</sup> and feeding rate zero, T3= 16 fish/m<sup>3</sup> and feeding rate zero, T4= 4 fish/m<sup>3</sup> and feeding rate 3%, T5= 8 fish/m<sup>3</sup> and feeding rate 3%, T6= 16 fish/m<sup>3</sup> and feeding rate 3%, T7= 4 fish/m<sup>3</sup> and feeding rate 4%, T8= 8 fish/m<sup>3</sup> and feeding rate 4%, T9= 16 fish/m<sup>3</sup> and feeding rate 4%.

**Feed and protein utilization:**

All criteria studied and presented in Table 4 showed again that T6 and T1 were the best (P≤0.05) and followed by T4 (even than the T2, T3, T5, T6, T8, and T9) concerning FE, FCR, PER, and PPV in Nile tilapia. There were significant differences between T6 and T1 in data of FCR, FE, PER, and PPV. Again, T3 and T6 were the worst, compared to other treatments. This agreed with the results obtained by Amar (2009), Chakraborty and Banerjee (2010) and Kapinga *et al.* (2014). Abdel- Tawwab *et al.* (2010) found that PER and FCR decreased with increasing dietary protein content. Moreover, fish stocking density could affect the efficiency of feed utilization as the number of fish stocked in a pond increase; the amount of feed available to each fish decreases (Chang, 1988). The FCR (3.03 to 6.20) in the current study is higher than the recommended FCR of 1.5 for aquaculture. The slight differences could stem from the differences in feed sources, environmental conditions and the particular strain or species of fish used. The protein efficiency ratio (PER) decreased with increasing fish density.

**Body composition:**

Values of dry matter (DM) crude protein (CP), ether extract (EE) and ash of the body is summarized in Table 5. The results of carcass composition of Nile tilapia showed that the difference were not significant (P>0.05) in crude protein, ether extract and ash percentages, but DM differed significantly. These results agree with the finding of Kapinga *et al.* (2014). The low lipid content of high-density-reared fish confirmed the findings of Montero *et al.* (1999), who found lower lipid levels in the liver of gilthead sea bream, reared at high density. Furthermore, changes in protein and lipid

contents in fish body could be linked with changes in their synthesis and/or deposition rate in the muscles (Abdel-Tawwab *et al.*, 2006). In contrast with the present results, Gallagher (1999) did not find significant differences in moisture, protein, lipid and ash contents in the whole body of sunshine bass fed with different protein levels.

**Biochemical parameters:**

The results of protein profile and kidney and liver functions (ALT, AST, creatinine, urea and uric acid) showed significant increases in total protein and globulin and significant decrease in albumin in the treatments T3, T6, T7 and T8, but there were no significances among fish groups T1, T2, T5 and T9. These results are illustrated in Tables 6 and 7. These results were supported by Kapinga *et al.* (2014). Results in tables (6 and 7) showed the cortisol glucose, cholesterol, total protein, albumin, globulin, RBCs, ALT, AST, creatinine, urea, uric acid and WBCS count in the experimental fish of treatment (Feeding rate 4% and stocking density 4 fish/m<sup>3</sup>) were increased significantly ( $P \leq 0.05$ ) compared with the among treatment (Feeding rate zero level, 3% and stocking density 8 fish/m<sup>3</sup>). On the other side by increasing the stocking densities of fish, no significant ( $P \leq 0.05$ ) differences were recorded in all above mentioned blood parameters compared with the treatment number (1). These results were supported by Rafatnezhad *et al.*, (2008) found that hematological parameters of the great sturgeon did not change in response to change of density.

**Table 4: Effect of stoking density and feeding rate on feed utilization of Nile tilapia of Nile tilapia (*Oreochromis niloticus*).**

Feeding rate	Overall regardless of feeding rate			FR-Level 3%			FR-Level 4%		
	T1	T2	T3	T4	T5	T6	T7	T8	T9
Stocking density	4 f/m <sup>3</sup>	8 f/m <sup>3</sup>	16 f/m <sup>3</sup>	4 f/m <sup>3</sup>	8 f/m <sup>3</sup>	16 f/m <sup>3</sup>	4 f/m <sup>3</sup>	8 f/m <sup>3</sup>	16 f/m <sup>3</sup>
FCR	1.22± 0.23c	1.37± 0.54b	1.48± 0.90a	1.06± 0.04c	1.15± 0.30b	1.30± 0.03a	1.39± 0.22c	1.58± 0.26b	1.65± 0.70a
FE	0.81± 0.02a	0.73± 0.01b	0.67± 0.00c	0.95± 0.00a	0.87± 0.01b	0.77± 0.00c	0.71± 0.02a	0.63± 0.00b	0.60± 0.00c
PER	2.96± 0.07a	2.60± 0.04b	2.40± 0.01c	3.36± 0.02a	3.08± 0.03b	2.72± 0.00c	2.55± 0.06a	2.25± 0.01b	2.15± 0.01c
PPV (%)	25.75± 1.69a	11.50± 0.83b	5.34± 0.27c	22.30± 1.28a	9.92± 0.27b	4.80± 0.28c	22.30± 1.28a	9.92± 0.27b	4.80± 0.28c

Means bearing different small letters are significant at ( $P < 0.05$ ).

T1= 4 fish/m<sup>3</sup> and feeding rate zero, T2= 8 fish/m<sup>3</sup> and feeding rate zero, T3= 16 fish/m<sup>3</sup> and feeding rate zero, T4= 4 fish/m<sup>3</sup> and feeding rate 3%, T5= 8 fish/m<sup>3</sup> and feeding rate 3%, T6= 16 fish/m<sup>3</sup> and feeding rate 3%, T7= 4 fish/m<sup>3</sup> and feeding rate 4%, T8= 8 fish/m<sup>3</sup> and feeding rate 4%, T9= 16 fish/m<sup>3</sup> and feeding rate 4%.

**Table 5: Effect of stoking density and feeding rate on the chemical composition of the whole bodies of Nile tilapia (*Oreochromis niloticus*), % DM basis.**

Feeding rate		Overall regardless of feeding rate			FR-Level 3%			FR-Level 4%		
Stocking density		T1 (4 f/m <sup>3</sup> )	T2 (8 f/m <sup>3</sup> )	T3 (16 f/m <sup>3</sup> )	T4 (4 f/m <sup>3</sup> )	T5 (8 f/m <sup>3</sup> )	T6 (16 f/m <sup>3</sup> )	T7 (4 f/m <sup>3</sup> )	T8 (8 f/m <sup>3</sup> )	T9 (16 f/m <sup>3</sup> )
Parameters	Initial analysis									
Dry matter (DM)	36.97± 0.42b	33.45± 0.13b	33.55± 0.20b	34.40± 0.27a	34.47± 0.55a	33.30± 0.3a	33.40± 0.25a	34.33± 0.24a	33.80± 0.06ab	33.50± 0.15b
Crud protein (CP)	61.20± 0.21b	68.97± 0.46a	68.30± 0.51a	67.68± 0.71a	66.67± 1.09a	67.60± 0.6a	68.97± 0.87a	68.70± 0.55a	69.00± 0.61a	68.97± 0.57a
Ash	25.33± 0.07b	19.28± 0.36a	19.92± 0.17a	19.77± 0.18a	19.63± 0.30a	20.00± 0.2a	18.97± 0.66a	19.90± 0.23a	19.83± 0.28a	19.60± 0.36a
Ether extract (EE)	6.03± 0.15b	4.97± 0.12a	4.68± 0.24a	4.80± 0.16a	4.83± 0.33a	5.03± 0.38a	5.17± 0.12a	4.77± 0.12a	4.33± 0.12a	4.77± 0.15a
NFE	4.43± 0.20	4.57± 0.20a	4.52± 0.07a	4.85± 0.12a	4.93± 0.22a	4.60± 0.12a	4.57± 0.44a	4.77± 0.12a	4.43± 0.03a	4.57± 0.07a

Means bearing different small letters are significant at (P<0.05).

T1= 4 fish/m<sup>3</sup> and feeding rate zero, T2= 8 fish/m<sup>3</sup> and feeding rate zero, T3= 16 fish/m<sup>3</sup> and feeding rate zero, T4= 4 fish/m<sup>3</sup> and feeding rate 3%, T5= 8 fish/m<sup>3</sup> and feeding rate 3%, T6= 16 fish/m<sup>3</sup> and feeding rate 3%, T7= 4 fish/m<sup>3</sup> and feeding rate 4%, T8= 8 fish/m<sup>3</sup> and feeding rate 4%, T9= 16 fish/m<sup>3</sup> and feeding rate 4%.

**Table 6: Effect of stoking density and feeding rate on blood biochemical parameters of Nile tilapia (*Oreochromis niloticus*).**

Feeding rate		Overall regardless of feeding rate			FR-Level 3%			FR-Level 4%		
Stocking density		T1 (4 f/m <sup>3</sup> )	T2 (8 f/m <sup>3</sup> )	T3 (16 f/m <sup>3</sup> )	T4 (4 f/m <sup>3</sup> )	T5 (8 f/m <sup>3</sup> )	T6 (16 f/m <sup>3</sup> )	T7 (4 f/m <sup>3</sup> )	T8 (8 f/m <sup>3</sup> )	T9 (16 f/m <sup>3</sup> )
Cortisol (ug/l)	4.12± 0.07	4.15± 0.14	4.17± 0.13	4.10± 0.00	4.17± 0.13	4.20± 0.32	4.13± 0.18	4.13± 0.92	4.13± 0.17	
Glucose (mg/dl)	75.83± 1.47	76.17± 0.8	76.33± 1.05	74.33± 2.85	76.33± 0.33	77.33± 2.30	75.33± 0.67	76.00± 1.00	77.33± 0.67	
Cholesterol (mg/l)	214.33± 2.14	217.67± 3.0	214.33± 3.16	213.67± 4.91	217.00± 2.08	221.33± 2.03	211.67± 3.38	214.00± 5.69	215± 5.03	
T3 (IU/L)	2.32± 0.01	2.33± 0.03	2.32± 0.05	2.32± 0.01	2.33± 0.06	2.30± 0.09	2.31± 0.03	2.33± 0.01	2.33± 0.06	
T4 (IU/L)	1.35± 0.06	1.30± 0.08	1.38± 0.09	0.37± 0.03	1.37± 0.15	1.47± 0.13	1.33± 0.13	1.23± 0.09	1.30± 0.12	
Lysozyme (IU/L)	0.13± 0.01a	0.12± 0.01	0.12± 0.01	0.12± 0.01	0.11± 0.01	0.13± 0.01	0.13± 0.01	0.12± 0.01	0.11± 0.01	
Total protein (g/dl)	5.53± 0.09	5.55± 0.12	5.63± 0.18	5.43± 0.15	5.43± 0.20	5.60± 0.17	5.63± 0.09	5.67± 0.15	5.67± 0.37	
Albumin (g/dl)	3.53± 0.07	3.37± 0.08	5.63± 0.18	3.53± 0.13	3.23± 0.09	3.50± 0.06	3.53± 0.07	3.50± 0.06	3.37± 0.12	
Globulin (g/dl)	2.0	2.2	2.2	1.9	2.2	2.1	2.1	2.2	2.3	

T1= 4 fish/m<sup>3</sup> and feeding rate zero, T2= 8 fish/m<sup>3</sup> and feeding rate zero, T3= 16 fish/m<sup>3</sup> and feeding rate zero, T4= 4 fish/m<sup>3</sup> and feeding rate 3%, T5= 8 fish/m<sup>3</sup> and feeding rate 3%, T6= 16 fish/m<sup>3</sup> and feeding rate 3%, T7= 4 fish/m<sup>3</sup> and feeding rate 4%, T8= 8 fish/m<sup>3</sup> and feeding rate 4%, T9= 16 fish/m<sup>3</sup> and feeding rate 4%.



**Table 7: Effect of stoking density and feeding rate on blood parameters (liver and kidney functions) of Nile tilapia (*Oreochromis niloticus*)**

Feeding rate	Overall regardless of feeding rate			FR-Level 3%			FR-Level 4%		
	T1 (4 f/m <sup>3</sup> )	T2 (8 f/m <sup>3</sup> )	T3 (16 f/m <sup>3</sup> )	T4 (4 f/m <sup>3</sup> )	T5 (8 f/m <sup>3</sup> )	T6 (16 f/m <sup>3</sup> )	T7 (4 f/m <sup>3</sup> )	T8 (8 f/m <sup>3</sup> )	T9 (16 f/m <sup>3</sup> )
ALT (u/l)	65.67± 0.67	63.17± 0.75	64.3± 1.12	66.33± 0.88	63.00± 1.00	65.33± 1.76	65.00± 1.00	63.33± 1.33	63.3± 30.33
AST (u/l)	55.83± 0.60	56.67± 0.95	56.00± 1.03	55.00± 0.58	57.00± 1.53	57.33± 1.86	56.67± 0.88	56.33± 1.45	54.67± 0.33
Creatinine (mg/dl)	1.160± 0.13	0.92± 0.13	1.00± 0.13	1.12± 0.18	0.81± 0.18	1.03± 0.18	1.20± 0.18	1.02± 0.18	1.00± 0.18
Urea (mg/dl)	22.0± 1.00	22.3± 1.00	23.7± 1.00	22.3± 1.42	21.7± 1.42	23.3± 1.42	21.7± 1.42	23.1 0.42	24.0± 1.42
Uric acid (mg/dl)	2.37± 0.06	2.36± 0.06	2.30± 0.06	2.32± 0.09	2.33± 0.09	2.30± 0.09	2.41± 0.09	2.30± 0.06	2.30± 0.09

T1= 4 fish/m<sup>3</sup> and feeding rate zero, T2= 8 fish/m<sup>3</sup> and feeding rate zero, T3= 16 fish/m<sup>3</sup> and feeding rate zero, T4= 4 fish/m<sup>3</sup> and feeding rate 3%, T5= 8 fish/m<sup>3</sup> and feeding rate 3%, T6= 16 fish/m<sup>3</sup> and feeding rate 3%, T7= 4 fish/m<sup>3</sup> and feeding rate 4%, T8= 8 fish/m<sup>3</sup> and feeding rate 4%, T9= 16 fish/m<sup>3</sup> and feeding rate 4%.

## CONCLUSIONS

it was proved that Nile tilapia fingerlings reared in floating hapas (in an earthen pond) stocked at 4 fish/m<sup>3</sup> and fed a diet containing 28.5% crude protein at 4% of the biomass daily gave the best results at the end of the experiment (98 days), concerning growth performance, feed and nutrients utilization, chemical composition, and blood profile.

## REFERENCES

- Abdel-Hakim, N.F.; Amar, A. A. and Abd-Elgawad, A. S. (2008). Effect of initial stocking size and production cycle on growth performance of monosex tilapia reared in earthen ponds. 8<sup>th</sup> International Symposium on Tilapia in Aquaculture, pp: 255-269.
- Abdel-Hakim, N.F., Bakeer, M.N. and Soltan, M.A. (2002). Water Environment for Fish Culture. Deposition No. 4774.
- Abdel-Hakim, N.F., El-Nemaki, F.A., El-Gamal, A.A. and Abdel-Warith, A.A. (1995). Effect of different stocking rates on growth performance. Egypt.J.Agric. Res., 73 (3): 873-887.
- Abdel-Hakim, N.F., Hilali, I.A., Khalil, M.H. and Al-Azab, A.A. (2001). Effect of stocking density and feeding rate on performance of Nile tilapia (*Oreochromis niloticus*) reared in Tanks. Egyptian J. Nutrition and Feeds (Special Issue): 705-717.
- Abdelhamid, A.M., khalil, F.F, El Barbary, M.I., Zaki, V.H. and Hussein, H.S. (2002). Feeding Nile tilapia on Biogen to detoxify aflatoxin diets. Proc. Con. Animal, Fish Prod. Mansoura, 24- 25 Sep., pp: 207-230.

- Abdel-Tawwab, M., Ahmad, M.H., Khattab, Y. and Shalaby, A.M.E. (2010). Effect of dietary protein level, initial body weight, and the interaction on the growth, feed utilization, and physiological alterations of Nile tilapia, *Oreochromis niloticus* (L.). *Aquaculture*, 298: 267–274.
- Abdel-Tawwab, M., Khattab, Y., Ahmad, M.H. and Shalaby, A.M.E. (2006). Compensatory growth, feed utilization, whole body composition and hematological changes in starved juvenile Nile tilapia, *Oreochromis niloticus* (L.). *J Appl. Aquacult.*, 18: 17–36.
- Amar, A.A. (2009). Effect of initial weight and stocking density on growth performance of mono sex Nile tilapia reared in semi intensive system. *Egypt. J. Aquat. Biol. & Fish.*, 13 (2): 69- 80.
- A.O.A.C. (1990). Official method of Analysis, 15<sup>th</sup> Ed. Association of official Analysis of chemists, Washington D.C.
- Awad, M.H. (2015). Effect of stocking density in performance of cultured fish and productivity in earthen ponds, M Sc. Thesis, Department of Animal Production, Faculty of Agriculture, Cairo, Al-Azhar University.
- Barlarin, J.D. (1979). Africa tilapia farm shows profit potential fish farm, *Int.*, 6: 16.
- Chakraborty, S.B. and Banerjee, S. (2010). Effect of stocking density on monosex Nile tilapia growth during pond culture in Indi world Academy of science Engineering and Technology, 68: 1511-1515.
- Chang, W.Y.B. (1988). Fish production: data synthesis and model development. In: Pond dynamics/aquaculture Collaborative Research Support Program (CRSP), 6<sup>th</sup> annual administrative report. Oregon State University, Oregon, pp: 41–49.
- Coche, A.G. (1976). A general review of cage culture and its application in Africa. *FAO Tech. Conf. on Aquaculture*, Kyoto, Japan, FIR AQ/ Conf. 76/E., 72: 33.
- Duncan, D. (1955). Multiple range and multiple F-tests. *Biometrics*, 11: 1-42.
- GAFRD (2006). General Authority for Fish Resources Development. Fish Production Statistics year 2006. Ministry of Agriculture Cairo, Egypt.
- Gallagher, M.L. (1999). Growth response, tissue composition, and liver enzymes changes in juvenile sunshine bass *Morone chrysops* × *M. saxatilis*, associated with dietary protein and lipid level. *J. Appl. Aquacult.*, 9: 41–51.
- Henry, R.J. (1964). *Clinical Chemistry: principles and techniques*. Harper and Row Publications, New York.
- Jobling,S.(1983). A total protein and acid needs. In i J.E. Halve(Nutrition. Academic press: Nell york,pp.106-143.
- Kapinga, I., Mlaponi, N. and kasozi, N. (2014). .Effect of stocking density on the growth performance of sex reversed mole Nile tilapia (*Oreochromis niloticus*) under pond conditions in Tanzania. *World Journal Fish and Marine Sciences*, 6 (2): 156-161.
- Kohin, J. A. (1958). Micro electrophoretic method. *Nature*, 181: 838.
- Moav. R.; Wohlfarth, G.W.; Schroeder, G.L.; Hulata, G. and Barash, H. (1977). Intensive polyculture of fish in fresh water ponds. I.substitution of expensive foods by liquid cow manure. *Aquacult.*, 10 (1): 25-43.

- Montero, D., Izquierdo, M.S., Tort, L., Robaina, L. and Vergara, J.M. (1999). High stocking density produces crowding stress altering some physiological and biochemical parameters in gilthead seabream, *Sparus aurata*, juveniles. *Fish Physiol. Biochem.*, 20: 53–60.
- NRC (1993). Nutrient Requirement of Fish. National Academy Press. Washington, Dc, P.114.
- Pagan, F.A. (1970). Cage culture of tilapia. *FAO fish Cult. Bull.*, 3(1): 6.
- Rafatnezhad, S., Falahatkar, B. and Gilani, M.H.T. (2008). Effects of stocking density on hematological parameters, growth and fin erosion of great sturgeon (*Huso huso*) juveniles. *Aquaculture Research*, 39: 1506-1513. doi:10.1111/j.1365-2109.2008.02020.x;
- Reitman, S. and Frankel, S. (1957). Colorimetric determination of glutamic oxaloacetic and glutamic pyruvic transaminases. *Am. J. Clin. Path.*, 28: 53–56.
- SAS (2006). SAS statistical guide for personal computer, SAS Institute Inc. Cary, NC.
- Sumanasart, P. (1971). Effect of feeding mesh size and stocking of *Tilapia aurea* in cages. *Ann. Rep. Int. Centre. Aquacult. Agric. Exp. St.*, Auburn Univ. Alabama, 71: 9.
- Trinder, P. (1969). Determination of glucose concentration in the blood. *Ann. Clin. Biochem.*, 6: 24–27.
- Zannatul. F., Masum. A. and Ali. M. (2014). Influence of stocking density on growth performance and survival of monosex tilapia (*Oreochromis niloticus*) fry. (in press).

### تأثير معدلات الكثافة و التغذية على إنتاج الأحواض الترابية المستزرعة بأسماء البلطي النيلي

- محمد الشحات محمد رمضان<sup>١</sup>، نبيل فهمي عبد الحكيم<sup>٢</sup>، محمد محمد السعيد لأشيين<sup>٣</sup>،  
الدسوقي السيد محمد العزب<sup>١</sup> وأحمد محمد زغلول<sup>١</sup>
- <sup>١</sup> - قسم الإنتاج الحيواني، إنتاج الأسماك، كلية الزراعة جامعة الأزهر بالقاهرة.
- <sup>٢</sup> - قسم الإنتاج الحيواني، تغذية الحيوان، كلية الزراعة جامعة الأزهر بالقاهرة.
- <sup>٣</sup> - قسم الإنتاج الحيواني، رعاية الحيوان، كلية الزراعة جامعة الأزهر بالقاهرة.

أجريت هذه الدراسة على مدى فترة ٩٨ يوماً لدراسة تأثير مستويات كثافة ومعدلات تغذية مختلفة على إنتاج الأحواض الترابية المستزرعة بأسماء البلطي النيلي، وتم استخدام الهبات المعلقة في حوض تربي بمساحة ٤٠٠٠م<sup>٢</sup>، وكانت أبعاد كل هابة ٢\*٤\*١م. وكانت عدد المعاملات ٦ مقسمة الى ثلاث مكررات بمجموع ١٨ هابة. وكانت مستويات الكثافة التخزينية للأسماك ٤، ٨، ١٦ سمكة/م<sup>٣</sup>، فكانت أعداد الأسماك في المجموعة ١ (٣٢) والمجموعة ٢ (٦٤) والمجموعة ٣ (١٢٨) على الترتيب، وكانت معدلات التغذية (صفر، ٣، ٤ %) من الوزن الحى لجسم الأسماك. تم إتباع نظام غذائي متوازن على طول فترة التجربة باستخدام عليقة ذات بروتين خام ٢٨.٥ % . تشير النتائج إلى أن المعاملة ذات الكثافة ٤ أسماك/م<sup>٣</sup> والتي غُذيت بمعدل تغذية يومي ٤ % هي الأفضل من حيث الوزن الحى لجسم الأسماك والاستفادة الغذائية وتركيب الجسم وكذلك الحالة الفسيولوجية.

