

EFFECT OF ROOTSTOCK AND INTERSTOCK ON GROWTH, YIELD AND FRUIT QUALITY OF SOME ORANGE VARIETIES A. VEGETATIVE GROWTH, NUTRITIONAL STATUS AND YIELD

Somaia A. El-Sayed

Citrus Research Department, Hort. Res. Instit. ARC. Giza, Egypt.

Received: Feb. 22 , 2017

Accepted: Apr. 8 , 2017

ABSTRACT: *This experiment was designed as a long term study, started in 2008 till 2014 with the idea of using sour orange as interstock on Volkamer lemon rootstock (SO/VL) for budding four orange varieties, namely, Olinda valencia, Spring navel, Parent navel, and Fukumoto navel oranges. The purpose was to avoid some disadvantages of Volkamer lemon on physical and chemical fruit quality with the hope of maintaining tree growth and its productivity with good properties acceptable for local and foreign market. The obtained results showed that, Olinda valencia, Spring navel, Parent navel and Fukumoto navel oranges on Volkamer lemon rootstock showed the best tree size with higher growth parameters, while the interstock gave intermediate values with most vegetative growth parameters. On the other hand the lowest values of tested growth parameters were obtained on the scions budded on sour orange. In this respect, Olinda valencia orange gave largest tree size and strong vegetative growth parameters compared to other varieties. The highest values of chlorophyll a, b and total chlorophyll were recorded on Spring navel orange budded on Volkamer lemon rootstock. Leaf NPK analysis showed that maximum leaf nitrogen content was recorded on Olinda valencia and Parent navel oranges. Phosphorus was maximum for Spring navel orange. As for potassium, it reaches maximum values in leaves of Parent and Fukumoto navel oranges. In addition, yield in terms of weight (kg) of fruits/tree - kg/cm³ of canopy volume and kg/cm² of trunk cross section area (TCSA). were high on Volkamer lemon followed by sour orange rootstock, and intermediate on interstock. The interstock tended produce yield near to that on sour orange rootstock with better fruit properties as found in part (B) of this study.*

Key words: *Rootstock, Interstock, Budwood, Volkamer lemon, Fukumoto navel, Parent navel, Olinda valencia*

INTRODUCTION

Rootstocks selection is a major consideration in every citrus growing operation. It is fundamental to orchard success. Also, supporting the tree, the root system is responsible for absorption of water and nutrients, adapting the scion to particular soil conditions, and potentially providing tolerance to drought, salinity and some diseases (Louzada *et al.*, 1992 and Zayan *et al.*, 2004). More than twenty horticulture characters influenced by rootstock including for example tree vigour and size, nutritional status, yield and yield efficiency. In this respect, Zekri (2000) revealed that trunk cross sectional area (TCSA) and tree canopy volume of Valencia

orange grown on *C. volkameriana* were greater than those on Swingle citrumelo, Cleopatra mandarin and Milam lemon rootstocks. Dawood (2001 & 2002) found that, Valencia and Washington navel oranges on *C. volkameriana* had the largest tree size and vegetative growth as well as yield efficiency as compared with those recorded on sour orange. Also, Castle *et al* (2010) evaluated Valencia orange on 12 rootstocks for 15 years, they reported that, trees on Volkamer lemon was tallest, higher trunk cross sectional than other rootstocks.

Workers on Horticulture uses interstocks for many reasons such as avoiding scion/rootstock incompatibility, control of tree size and tolerance of some negative soil

properties as well as to prevent injury from several trunk diseases with the use of resistant interstock trunks (Krezdorn 1978, Shokrollah *et al.*, 2011, Gimeno *et al.*, 2012, Aboutalebi and Hassanzadeh 2014). Furthermore, interstocks may improve tree growth, yield and fruit quality. Castle 1992, Girardi and Filho 2006, Bakry *et al.*, 2007 and Yilmaz *et al.*, 2015, they studied the effect of interstocks on vegetative growth, leaf mineral content and amount of chlorophyll in leaf. Therefore, scions on volkamer lemon gave strong tree in size and growth as well as tolerance to environmental conditions.

The purpose of this long term study is to avoid or get rid of disadvantages of volkamer lemon on physical and chemical fruit properties , hopping to maintain tree growth and its productivity. Therefore, influences of volkamer lemon , sour orange rootstocks and interstock on growth , nutritional status and yield efficiency of four (scions) orange varieties namely, Olinda valencia, Spring navel, Parent navel and Fukumoto navel orange were studied in 2013 and 2014 seasons.

MATERIALS AND METHODS

This experiment is designed as a long term study started in 2008 with the idea of using double budding for volkamer lemon (VL) rootstock by using sour orange (SO) as interstock (SO/VL) for budding four orange varieties. In 2008 budwood from sour orange (SO) (*Citrus aurantium*) were budded on one year Volkamer lemon (VL) (*Citrus Volkameriana*) seedlings as interstock .Six months later, (VL), (SO) and (SO/VL) rootstocks were prepared at the same age and budded in 2009 with four

orange varieties, namely Olinda valencia, Spring navel, Parent navel, and Fukumoto navel orange varieties. The experiments included twelve treatments were arranged in a randomized complete block design, each treatment replicated 3 times and 3 plots for a total of 9 tree per each rootstock.

Thus, 108 budded trees (12x9) were planted in 2010 in a private orchard at El-Bustan region, El- Beheira Governorate, Egypt planted at 5x5 meter apart. The soil is sandy and the mechanical and chemical analysis was done as shown in Table (1). All agricultural practices were done as usual in the orchard.

In 2013 and 2014 seasons, samples and field data were recorded as follows:

- 1- Olinda valencia orange on sour orange (SO).
- 2- Olinda valencia orange on volkamer lemon (VL).
- 3- Olinda valencia orange on interstock (SO/VL).
- 4- Spring navel orange on sour orange (SO).
- 5- Spring navel orange on volkamer lemon (VL).
- 6- Spring navel orange on interstock (SO/VL).
- 7- Parent navel orange on sour orange (SO).
- 8- Parent navel orange on volkamer lemon (VL).
- 9- Parent navel orange on interstock (SO/VL).
- 10- Fukumoto navel orange on sour orange (SO).
- 11- Fukumoto navel orange on volkamer lemon (VL).
- 12- Fukumoto navel orange on interstock (SO/VL).

Table (1). Mechanical and chemical analysis of experimental soil.

Mechanical			Chemical		Cations (meq/l)				Anions (meq/l)		
Sand %	Silt %	Clay %	pH	Ec dS/m ⁻¹	Na ⁺	Ca ⁺⁺	Mg ⁺⁺	CO ₃ ⁻	HCO ₃ ⁻	Cl ⁻	SO ₄ ⁻⁻
77.85	6.50	15.65	8.82	0.64	2.53	1.45	0.60	-	2.23	2.10	0.25

Effect of rootstock and interstock on growth, yield and fruit quality

During 2013 and 2014 seasons the following data were recorded:

1. Vegetative growth:

1.1. Leaf parameters:

Leaves formed in spring shoot were counted in both seasons, twenty mature leaves were sampled in August from spring shoot to determining leaf area (cm²) using a leaf area meter Model Li 3100 area-meter, then total leaf area (m²) of spring shoot was calculated.

1.2. Shoot parameters:

Spring shoot formed by spring growth cycle was counted and measured as (cm) by ruler, then total growth (m) of spring shoots was calculated in both seasons.

1.3. Tree vigour:

Tree height (m) was measured from soil surface to the end of growth, trunk circumferences (cm) was measured by using stripe measurement. Canopy volume was calculated according to the following equation: $CV = 0.528 \times H \times D^2$. Whereas, H = tree height, D = tree diameter (Castle 1983).

2. Nutritional status:

2.1. Leaf chlorophyll content $\mu\text{g}/\text{cm}^2$:

Chlorophyll a and b were extracted from fresh leaves with N, N-dimethyl formamide and determined Spectrophotometrically at wave-length of 664 and 647 nm and then total chlorophyll was estimated according to the method described by Moran (1982).

2.2. Leaf NPK content:

Leaves sample was taken in September and washed with tap water followed by distilled water. Leaves were oven dried at 70°C to a constant weight. Dry weight was calculated then the dry leaves were ground and digested according to Chapman and Pratt (1961) and Jackson (1967). N, P and K. Total nitrogen % was determined by using the micro-kjeldahl method as described by Pregl (1945), Phosphorus % was determined colorimetrically as described by Murphy and Riley (1962) while, Potassium

% was determined by using flame photometer as described by Brown and Lilleland (1974).

3. Yield:

At harvest time (December in both seasons), yield of each tree was determined as weight (kg) of fruits/tree. Yield efficiency was calculated as kg/cm³ of canopy volume and kg/cm² of trunk cross section area (TCSA).

Statistical analysis:

Statistical analysis was done as analysis of variance according to Snedecor and Cochran (1990), and the least significant differences (L.S.D. at 5% level) was used to compare the mean values.

RESULTS AND DISCUSSION

1. Vegetative growth:

1.1. Leaf parameters:

Data presented in Table (2) show the effect of rootstock and interstock on orange varieties and their interaction on leaf growth parameters in both seasons. As for the effect of rootstock and interstock, it is clear that most leaf growth parameters were significantly influenced by rootstocks and interstock. Trees on volkamer lemon (VL) rootstock have had the highest values of leaves number per spring shoot, leaf area per spring shoot and total leaves area of spring shoot followed in a descending order by interstock (SO/VL) and sour orange (SO) rootstock in both seasons respectively. The differences were significant among them in both seasons. These results are in agreement with Dawood (2002) and Zayan *et al.*, (2004). In this respect, Mohamed (2011) and Hikal (2014) revealed that volkamer lemon rootstock reported the highest significant effect of leaves number per plant and leaf area of Balady lime and Washington navel orange compared to sour orange, Rangpur lime and Troyer citrange rootstocks.

Table (2). Effect of rootstock and interstock on leaf growth parameters of some orange varieties in 2013 and 2014 seasons.

Treatments		Rootstocks and interstock											
V	R	Leaves number / spring shoot				Leaf area cm ² from spring shoot				Total leaves area of spring shoots m ²			
		VL	SO	SO/VL	Mean	VL	SO	SO/VL	mean	VL	SO	SO/VL	Mean
2013													
Olinda valencia		8.33	5.67	6.33	6.78	35.02	26.50	31.10	30.87	2.92	1.51	1.98	2.14
Spring navel		7.67	5.33	5.67	6.22	24.50	20.33	22.17	22.33	1.88	1.08	1.26	1.41
Parent navel		7.33	5.33	7.33	6.66	28.80	24.80	26.89	26.83	2.11	1.33	1.97	1.80
Fukumoto navel		7.67	7.00	7.33	7.33	29.40	24.33	26.15	26.65	2.25	1.73	1.93	1.97
Mean		7.75	5.83	6.67		29.45	23.99	26.58		2.29	1.41	1.79	
L.S.D. at 5%		V = 0.82 R = 0.50 VxR = 0.97				V = 4.51 R = 1.18 VxR = 2.36				V = 0.88 R = 0.31 VxR = 0.62			
2014													
Olinda valencia		11.67	7.67	7.67	9.00	35.35	25.01	33.88	31.41	4.13	1.92	2.58	2.88
Spring navel		10.00	7.67	7.67	8.45	32.30	22.21	25.88	26.80	3.23	1.71	1.99	2.31
Parent navel		9.33	7.33	8.00	8.22	29.29	18.57	28.28	25.38	2.70	1.36	2.27	2.11
Fukumoto navel		10.00	8.33	8.33	8.89	33.71	22.87	26.06	27.55	3.40	1.92	2.19	2.50
Mean		10.25	7.75	7.92		32.66	22.17	28.53		3.37	1.73	2.26	
L.S.D. at 5%		V = 1.42 R = 0.50 VxR = 1.00				V = 4.21 R = 1.40 VxR = 4.09				V = 0.63 R = 0.27 VxR = 0.54			

SO = Sour orange, VL = Volkamer lemon, V = Variety, R = Rootstock, SO/VL = Interstock

Regarding the effect of orange varieties, the results showed that, no significant differences were found for most leaf growth parameters among orange varieties in both seasons. Anyhow, Olinda valencia orange showed the higher values of leaf growth parameters than other orange varieties in the second season. Similar results were obtained by Martinez *et al.* (1994) and Dawood (2002).

As for the effect of interaction between rootstock or interstock and scion cultivars, it is clear that, Olinda valencia orange on volkamer lemon rootstock gave highest values of leaf growth parameters followed by Fukumoto and Spring navel orange on the same rootstock, while the lowest values were obtained from Spring navel orange on sour orange rootstock followed by Parent navel orange on the same rootstock in both seasons. The interstock treatment(SO/VL) gave intermediate values with all orange varieties in both seasons. These results are

in line with those reported by Abd Alla (1999) who reported that Washington navel orange, valencia orange and Balady mandarin on volkamer lemon and Rangpur lime exhibited the most vigorous growth compared to sour orange and other rootstocks.

1.2. Shoot parameters:

Data in Table (3) showed that the tested rootstocks were significantly affected on spring shoot growth parameters of the four orange varieties in both seasons. Olinda valencia, Spring navel, Parent navel and Fukumot navel orange had the highest spring shoot number, spring shoot length and total growth of spring shoots on volkamer lemon rootstock followed by those budded on the interstock with significant differences between them in both seasons. On the other hand, the lowest values of spring shoot growth parameters are belonged to scions budded on sour orange rootstock in both seasons while the

Effect of rootstock and interstock on growth, yield and fruit quality

interstock gave intermediate values of spring shoot growth parameters. The obtained results concerning the effect of citrus rootstocks on scion shoot growth parameters go in line with those mentioned by Dawood (2001 & 2002), Zayan *et al.*, (2004) and Hikal (2014).

With regard to the effect of four orange varieties, it was evident that all spring shoot growth parameters were not significant among all orange varieties in the first season only. Olinda valencia orange had the highest values of spring shoot number, spring shoot length and total growth of spring shoots followed by Spring navel orange without significant differences between them in both seasons. Parent navel orange and Fukumoto navel orange gave the lowest values of spring shoot parameters with significant differences between them and

other varieties in the second season only. The results are in line with those obtained by Sayed and Abdel-Aziz (2010).

Also, a significant interaction effect between rootstock and interstock and scion varieties on spring shoot growth parameters was obtained. The highest values of spring shoot number, spring shoot length and total growth of spring shoots were occurred with Olinda valencia orange and Spring navel orange budded on volkamer lemon rootstock, while the lowest values was obtained from Fukumoto navel orange on sour orange rootstock. This result was more pronounced in the second season. Parent navel orange gave intermediate values in this respect. This results are in general agreement with those found by Bakry *et al.*, (2007).

Table (3). Effect of rootstock and interstock on spring shoot growth parameters of some orange varieties in 2013 and 2014 seasons.

Treatments		Rootstocks and interstock											
R	V	Spring shoot Number				Spring shoot length (cm)				Total growth of spring shoots (m)			
		VL	SO	SO/VL	Mean	VL	SO	SO/VL	mean	VL	SO	SO/VL	Mean
2013													
Olinda valencia		10.33	7.33	8.00	8.55	15.76	8.10	10.30	11.39	1.63	0.59	0.82	1.01
Spring navel		9.33	8.00	8.33	8.55	13.10	8.03	10.10	10.41	1.22	0.64	0.84	0.90
Parent navel		9.00	6.67	7.33	7.67	12.13	7.93	10.25	10.10	1.09	0.53	0.75	0.79
Fukumoto navel		6.33	6.00	6.33	6.22	9.77	8.17	8.10	8.68	0.61	0.49	0.51	0.51
Mean		8.75	7.00	7.50		12.54	8.06	9.69		1.10	0.56	0.73	
L.S.D. at 5%		V= 2.87 R=0.57 VxR= 1.14				V= 2.12 R=1.43 VxR=2.86				V= 0.23 R= 0.12 VxR= 0.24			
2014													
Olinda valencia		23.00	15.00	17.00	18.33	22.40	16.50	19.37	19.42	5.15	2.48	3.30	3.64
Spring navel		18.33	18.33	15.00	17.22	19.73	11.40	17.57	16.23	3.62	2.09	2.64	2.78
Parent navel		14.00	12.67	13.00	13.22	18.67	8.62	14.83	14.04	2.61	1.09	1.83	1.84
Fukumoto navel		13.00	12.67	20.67	15.45	16.67	9.77	13.83	13.42	2.16	1.24	2.86	2.09
Mean		17.08	14.67	16.42		19.37	11.57	16.40		3.39	1.73	2.66	
L.S.D. at 5%		V =1.71 R=1.72 VxR =0.43				V =2.32 R=1.76 VxR=2.35				V =0.55 R=0.20 VxR=0.40			

SO = Sour orange, VL = Volkamer lemon, V = Variety, R= Rootstock, SO/VL = Interstock

1.3. Tree vigour:

Data in Table (4) showed that, Volkamer lemon, sour orange and interstock were significantly effected on tree height, canopy volume and trunk circumferences of the four orange varieties in both seasons. Trees on volkamer lemon rootstock had the highest values of tree height, canopy volume and trunk circumferences followed by those on sour orange rootstock and interstock in both seasons respectively. The differences between sour orange and interstock were not significant in most cases. Similar results were obtained by Mansour *et al.*, (1993).

In according to orange varieties, data in Table (4) showed that, Olinda valencia orange have had tallest and largest canopy volume and trunk circumferences followed by Fukumoto navel orange as compared with the other varieties in both seasons. Spring navel and Parent navel oranges gave

the lowest values, and were found to be approximately at par during both seasons. These results are similar to those of Sayed and Adawy (2009).

The interaction between the two factors revealed that, Olinda valencia orange and Fukumot navel orange budded on volkamer lemon rootstock gave the highest values of tree height, canopy volume and trunk circumferences compared to other treatments in both seasons. On the other hand, Spring navel and Parent navel orange budded on sour orange or interstock had the lowest values in both seasons. The other treatments gave intermediate values for tree height, canopy volume and trunk circumferences in both seasons. Similar results about high, tree height, canopy volume and trunk circumferences of volkamer lemon were reported by Dawood, (2002) and Zayan *et al* (2004).

Table (4). Effect of rootstock and interstock on tree growth and vigour of some orange varieties in 2013 and 2014 seasons.

Treatments		Rootstocks and interstock											
R	V	Tree height (m)				Canopy volume (m ³)				Trunk circumferences (cm)			
		VL	SO	SO/VL	Mean	VL	SO	SO/VL	Mean	VL	SO	SO/VL	Mean
2013													
	Olinda valencia	2.38	1.72	1.65	1.92	2.68	2.28	2.09	2.35	15.60	6.67	6.17	9.48
	Spring navel	1.60	1.37	1.31	1.43	2.05	1.86	1.30	1.74	13.37	5.83	5.23	8.14
	Parent navel	1.78	1.38	1.36	1.51	2.20	1.96	1.49	1.88	13.83	6.17	5.50	8.50
	Fukumoto navel	2.14	1.68	1.56	1.79	2.27	2.01	1.60	1.91	15.00	6.53	5.70	9.08
	Mean	1.98	1.54	1.47		2.30	2.05	1.62		14.45	6.30	5.65	
	L.S.D. at 5%	V= 0.30 R= 0.18 VxR= 0.35				V= 0.46 R= 0.22 VxR= 0.43				V= 0.64 R= 0.52 VxR= 1.05			
2014													
	Olinda valencia	3.09	2.00	1.92	2.34	3.36	2.76	2.56	2.79	22.67	12.67	12.27	15.87
	Spring navel	2.06	1.67	1.64	1.79	2.48	2.33	1.95	2.25	15.17	9.17	8.67	11.00
	Parent navel	2.09	1.73	1.67	1.83	2.87	2.56	1.99	2.47	15.67	10.83	10.13	12.21
	Fukumoto navel	2.55	1.83	1.77	2.05	2.89	2.57	2.11	2.52	18.00	11.50	10.30	13.27
	Mean	2.45	1.81	1.75		2.90	2.56	2.15		17.88	11.04	10.34	
	L.S.D. at 5%	V= 0.54 R= 0.24 VxR= 0.48				V= 0.76 R= 0.35 VxR= 0.70				V=1.12 R=1.05 VxR=2.10			

SO = Sour orange, VL = Volkamer lemon, V = Variety, R= Rootstock, SO/VL = Interstock

Effect of rootstock and interstock on growth, yield and fruit quality

Generally, it is obvious from Tables (2, 3 and 4) that, volkamer lemon rootstock showed the best tree size and growth parameters represented by leaves number per spring shoot, leaf area per spring shoot, total leaves area of spring shoot, spring shoot number, spring shoot length, total growth of spring shoots, tree height, canopy volume and trunk circumferences. Using sour orange as Interstock gave intermediate values with most vegetative growth parameters, on the other hand the lowest values of the tested growth parameters were obtained from scions budded on sour orange. These results are similar to those obtained by Perez-Zamora *et al.*, (2002), Castle *et al.*, (2010) and Shafieizargar *et al.*, (2012). Moreover, Bakry *et al.*, (2007) they observed that Washington navel orange and Balady mandarin budded on sour orange interstock on volkamer lemon gave intermediate values of most vegetative growth parameters compared to volkamer lemon and sour orange rootstocks. Such conclusions agree with those presented by Gimeno *et al.*, (2012) who reported that Verna lemon trees grafted on valencia orange or Castellano orange interstock on sour orange rootstock had higher root, stem

and total dry weight than that on sour orange rootstock.

As for comparison among orange varieties, Olinda valencia orange gave the largest tree size and strong vegetative growth parameters. Fukumoto, Parent and Spring navel oranges gave medium tree size and growth vigour.

2. Nutritional status:

2.1. Leaf chlorophyll content $\mu\text{g}/\text{cm}^2$:

Data in Table (5) clearly showed that, chlorophyll a, b and total content was significantly increased with scions budded on volkamer lemon rootstock in both seasons. Scions on sour orange rootstock recorded the lowest values of chlorophyll a, b and total content, while those on the interstock gave intermediate values in both seasons. the differences were significant between volkamer lemon and sour orange rootstock, while were not significant between volkamer Lemon rootstock and Interstock in both seasons. These results are in line with those reported by Abd Alla (1999) and Ennab (2003) on valencia orange, Washington navel orange and Balady mandarin on different rootstocks.

Table (5). Effect of rootstock and interstock on leaf chlorophyll content ($\mu\text{g}/\text{cm}^2$) of some orange varieties in 2013 and 2014 seasons.

V	R	Rootstocks and interstock											
		Chlorophyll A				Chlorophyll B				Total Chlorophyll a + b			
		VL	SO	SO/VL	Mean	VL	SO	SO/VL	mean	VL	SO	SO/VL	Mean
2013													
Olinda valencia		44.28	42.12	45.49	43.96	22.64	20.23	22.53	21.80	66.92	62.35	68.02	65.76
Spring navel		47.16	47.06	45.62	46.61	22.84	20.75	21.92	21.84	70.00	67.81	67.54	68.45
Parent navel		45.26	41.09	43.13	43.16	22.52	19.51	21.14	21.06	67.78	60.60	64.21	64.20
Fukumoto navel		44.47	40.92	43.99	43.13	20.88	19.25	19.69	19.94	65.35	60.17	63.68	63.07
Mean		45.29	42.79	44.56		22.22	19.93	21.32		67.51	62.73	65.86	
L.S.D. at 5%		V=2.92 R = 2.56 VxR = 5.13				V = 1.14 R= 1.27 VxR= 2.55				V = 2.11 R= 2.74 VxR= 5.48			
2014													
Olinda valencia		44.00	41.85	45.36	43.74	22.80	20.71	20.85	21.45	68.17	62.56	66.21	65.65
Spring navel		48.24	46.07	42.23	45.51	24.17	20.80	23.94	22.97	72.41	66.87	66.17	68.48
Parent navel		42.15	39.66	43.30	41.70	24.28	18.49	21.49	21.42	66.43	60.37	64.79	63.86
Fukumoto navel		45.27	41.26	44.40	43.64	22.79	19.84	21.40	21.34	68.06	61.10	65.80	64.99
Mean		44.92	42.21	43.82		23.51	19.97	21.92		68.43	62.18	65.74	
L.S.D. at 5%		V = 1.30 R= 2.36 VxR= 4.72				V= 2.02 R= 0.54 VxR = 1.09				V = 1.38 R= 2.27 VxR= 4.54			

SO = Sour orange, VL = Volkamer lemon, V = Variety, R= Rootstock, SO/VL = Interstock

As for orange varieties, data in Table (5) showed that, chlorophyll a, b and its total values were higher in Spring navel orange and Olinda valencia orange than the other varieties in both seasons. The significant differences were obtained between Spring navel orange and other orange varieties. These results are similar to those of Dawood (2002) and Zayan *et al.*, (2004).

Concerning the interaction between rootstock or interstock and scions, it is clear that the interaction effect on leaf chlorophyll a, b and total content was significant in both seasons. The highest values of chlorophyll a, b and total chlorophyll were found on Spring navel orange budded on volkamer lemon rootstock, while the lowest values were obtained from Parent navel and Fukumoto navel oranges trees budded on sour orange rootstock. All orange varieties budded on the interstock gave intermediate

values of chlorophyll a, b and its total value in both seasons. Similar results were obtained by Ataweia *et al.*,(2011).

2.2. Leaf NPK content:

The results in Table (6) revealed that, volkamer lemon rootstock exhibited the highest values of leaf NPK content of scions followed by interstock and sour orange in both seasons, respectively. These results are similar with those reported by Smith *et al.*, (2004) and Barakat *et al.*, (2013). In this respect, Ahmed *et al.*, (2007) reported that, leaf NPK content of Kinnow mandarin grafted on nine rootstocks was differed significantly, maximum value of NPK were recorded on rough lemon and volkamer lemon while, the minimum were recorded on Troyer citrange and Carrizo citrange rootstocks.

Table (6). Effect of rootstock and interstock on leaf NPK content of some orange varieties in 2013 and 2014 seasons.

Treatments		Rootstocks and interstock											
V \ R	N %				P %				K %				
	VL	SO	SO/VL	Mean	VL	SO	SO/VL	mean	VL	SO	SO/VL	Mean	
2013													
Olinda valencia	1.95	1.57	1.92	1.81	0.110	0.096	0.090	0.099	1.31	1.06	1.24	1.20	
Spring navel	1.63	1.59	1.35	1.52	0.128	0.115	0.096	0.113	1.33	1.20	1.24	1.26	
Parent navel	1.98	1.52	1.94	1.81	0.105	0.084	0.099	0.096	1.98	1.24	2.02	1.75	
Fukumoto navel	1.63	1.34	1.57	1.51	0.093	0.086	0.088	0.089	1.59	1.27	1.57	1.48	
Mean	1.79	1.51	1.70		0.109	0.095	0.093		1.55	1.19	1.52		
L.S.D. at 5%	V=0.094 R= 0.046 VxR=0.094				V= 0.006 R= 0.004 VxR=0.009				V =0.046 R=0.039 VxR=0.077				
2014													
Olinda valencia	1.97	1.62	1.96	1.85	0.105	0.097	0.096	0.099	1.34	1.22	1.26	1.27	
Spring navel	1.70	1.69	1.47	1.62	0.138	0.125	0.097	0.120	1.38	1.27	1.30	1.32	
Parent navel	2.12	1.58	2.13	1.94	0.109	0.088	0.104	0.100	1.99	1.25	2.02	1.75	
Fukumoto navel	1.71	1.46	1.67	1.61	0.097	0.089	0.091	0.092	1.80	1.39	1.93	1.71	
Mean	1.88	1.59	1.81		0.112	0.099	0.097		1.63	1.28	1.63		
L.S.D. at 5%	V = 0.11 R= 0.09 VxR= 0.18				V= 0.006 R= 0.005 VxR=0.009				V=0.053 R=0.943 VxR=0.086				

SO = Sour orange, VL = Volkamer lemon, V = Variety, R= Rootstock, SO/VL = Interstock

Effect of rootstock and interstock on growth, yield and fruit quality

As for the effect of orange varieties, it is clear from Table (6) that maximum leaf nitrogen content was recorded on Olinda valencia and Parent navel oranges, and minimum on Fukumoto navel orange in both seasons. Phosphor was maximum on Spring navel orange while minimum values were recorded in Fukumoto navel orange. Olinda valencia and Parent navel orange recorded intermediate values in both seasons. As regard to potassium, it was higher on Parent and Fukumoto navel oranges while the minimum value was recorded for Olinda valencia orange in both seasons. Similar results were obtained by Toplu *et al.*, (2008) and Aboutalebi *et al.*, (2012).

The interaction between the two factors revealed that, the highest leaf nitrogen value was found on Parent navel and Olinda valencia oranges budded on Volkamer lemon rootstock, and the lowest was found for Fukumoto navel orange budded on sour orange rootstock in both seasons. Leaf phosphor content recorded higher values in Spring navel orange on volkamer lemon and on sour orange rootstocks, and the lower values were noticed in Parent and Fukumoto navel oranges budded on sour orange rootstock. This result was true in both seasons. Parent navel orange budded on interstock or volkamer lemon had the highest leaf potassium content, and the lowest values were obtained from Olinda valencia orange on sour orange rootstock. These results are in agreement with those of Labanauskas and Bitters (1974); Bakry *et al.*, (2007) and Jahromi *et al.*, (2012).

3. Yield:

Data in Table (7) showed that, yield as weight (kg/tree) of Olinda valencia, Spring navel, Parent navel and Fukumoto navel oranges was significantly higher on volkamer lemon rootstock than that on sour orange rootstock and interstock in both seasons. Moreover, orange scions on sour orange and interstock had similar yield without significant differences between them in both seasons. Similar results about high productivity of volkamer lemon were found

by Zayan *et al.*, 2004 and Al-Obeed *et al.*, 2005. Such conclusions agree with those presented by Ramin and Alirezanezhad 2005 who reported that Ruby Red and Marsh grapefruit trees on volkamer lemon rootstock had more fruit number and weight (kg) per tree than those grown on Cleopatra mandarin and sour orange rootstock. In this respect Shafieizargar *et al.*, 2012 stated that Queen orange trees grafted on volkamer lemon rootstock had larger and heavier fruits than those on Cleopatra mandarin and S wingle citrumelo rootstocks. In this respect, Gardner, 1968 reported that fruit yield of valencia and Hamlin oranges did not show a significant difference by interstock.

With regard, the effect of oranges, it is clear that, Olinda valencia orange had significantly higher yield than other varieties in both seasons. Spring navel, Parent navel and Fukumoto navel oranges had similar yield without significant differences among them in both seasons.

Olinda valencia on volkamer lemon rootstock gave the highest yield compared to other rootstocks in both seasons. This result was more pronounced in the second season. The lowest yield harvested from Fukumoto navel orange was on the interstock in both seasons. Generally, it is clear that, scions on volkamer lemon rootstock produce higher yield when compared with sour orange rootstock and interstock.

Data in Table (7) revealed that, yield efficiency as kg/cm^2 of TCSA (trunk cross sectional area cm^2) and kg/m^3 canopy volume of tree was significantly affected by all treatments in both seasons. As for the effect of rootstocks and interstock, it is clear that yield efficiency as kg/cm^2 of TCSA was the highest for scions on Sour orange rootstock in both seasons. On the other hand, scions on volkamer lemon rootstock had the lowest values of yield efficiency as kg/cm^2 of TCSA during 2013 and 2014 seasons. Interstock produced trees gave

intermediate values in this respect. Significant differences were detected among rootstocks and interstock in both seasons. Also, yield efficiency as kg/m³ canopy volume was highest for scions on volkamer lemon rootstock followed in descending order by those on sour orange rootstock and interstock in both seasons.

As for orange varieties, the results in Table (7) showed that, no significant differences were found for yield efficiency as kg/ cm² of TCSA (trunk cross sectional area cm²) and kg/m³ canopy volume of the tree among orange varieties in both seasons, excepted kg/m³ of canopy volume was variable between Olinda valencia orange and Fukumot navel orange in the second season only.

Although, there is a significant interaction effect between the two factors, but there is

no constant trend among treatments in both seasons. Generally, yield efficiency was higher on volkamer lemon followed by sour orange rootstocks, and intermediate on the interstock. In this concern Dawood (2001) and Zayan *et al.*, (2004) conducted that, accumulative yields of valencia orange and Washington navel orange were higher from trees on volkamer lemon and Rangpur lime than those on Swingle citrumelo, Cleopatra mandarin and sour orange. Also, Georgiou, 2002 reported that volkamer lemon has been reported to significantly increase accumulative yield of Clementine mandarin compared with sour orange up to 45%. Moreover, higher yield efficiency was also reported for trees showed reduction in size by the used rootstocks (Castle and Phillips, 1980 and Roose *et al.*, 1989).

Table (7). Effect of rootstock and interstock on yield efficiency of some orange varieties in 2013 and 2014 seasons.

Treatments	Rootstocks and interstock											
R \ V	kg/tree				Kg /cm ² TCSA				Kg /m ³ canopy volume			
	VL	SO	SO/VL	Mean	VL	SO	SO/VL	Mean	VL	SO	SO/VL	Mean
2013												
Olinda valencia	18.00	12.33	11.50	13.94	0.94	4.12	3.29	2.78	6.73	5.94	5.05	5.91
Spring navel	14.83	7.67	7.33	9.94	0.99	2.97	2.42	2.13	6.65	3.70	5.82	5.39
Parent navel	12.17	9.67	9.17	10.34	0.86	4.47	4.41	2.91	5.91	6.21	4.90	5.67
Fukumoto navel	11.67	8.67	7.00	9.11		3.64	2.11	2.13	5.15	5.54	3.81	4.83
Mean	14.17	9.59	8.75			3.80	2.81		6.11	5.35	4.90	
L.S.D. at 5%	V = 1.33 R= 1.19 VxR= 2.37				V =0.86 R=0.34 VxR= 0.69				V = 1.49 R= 0.76 VxR= 1.53			
2014												
Olinda valencia	20.33	16.67	16.00	17.67	0.51	1.39	1.30	1.07	6.15	6.08	6.21	6.15
Spring navel	13.00	11.17	10.50	11.56	0.51	1.24	1.04	0.93	5.45	4.90	4.16	4.84
Parent navel	12.50	9.17	8.50	10.06	0.64	1.10	0.85	0.86	4.72	5.15	3.69	4.52
Fukumoto nave	11.83	9.00	8.00	9.61		1.50	1.22	1.14	4.10	4.71	3.22	4.01
Mean	14.42	11.50	10.75			1.31	1.22		5.11	5.21	4.32	
L.S.D. at 5%	V = 1.48 R = 0.87 VxR = 1.75				V = 0.31 R=0.21 VxR= 0.42				V = 1.65 R= 0.85 VxR= 1.69			

Effect of rootstock and interstock on growth, yield and fruit quality

SO = Sour orange, VL = Volkamer lemon, V = Variety, R= Rootstock, SO/VL = Interstock

REFERENCES

- Abd Alla, S.A.E. (1999). Physiological studies on some orange varieties budded on different rootstocks. Ph.D. Thesis, Fac. Agric. Kafr El-Sheikh, Tanta University.
- Aboutalebi, A. and H. Hassanzadeh (2014). Salinity and citrus rootstocks and interstocks. *International Journal of Plant, Animal and Environmental Sciences* 4(2):654 – 672.
- Aboutalebi, A., H. Hasanzada and M.H. Farahi (2012). Evaluation the effect of rootstock type on mineral elements concentration in shoot of budded sweet orange (*Citrus sinensis* var. Valencia). *Annals of Biological Research*, 3 (9):4531 – 4534.
- Ahmed, W., M.A. Nawaz, M.A. Iqbal and M.M. Khan (2007). Effect of different of rootstocks on plant nutrient status and yield in Kinnow mandarin (*Citrus reticulata* Blanco). *Pak. J. Bot.*, 39(5):1779 – 1786.
- Al-Obeed, R.S., M.M. Harhash and M.M. Sourour (2005). Performance of Marsh grapefruit and Mexican lime trees on seven rootstocks in Saudi Arabia. *J. Adv. Agric. Res.*, (Fac. Agric. Saba Basha) 10 (1):165 – 179.
- Ataweia, A.A., F.M. El-Gindy, F.H. Ismail and F.I. Fahmy (2011). Physiological and anatomical studies on budding of citrus. *Egypt. J. App. Sci.*, 26(7):208 – 220.
- Bakry, K. A., M.A. Abd-El Rahman and M.M. Sholah (2007). Growth and mineral composition of some citrus species seedlings as influenced by rootstocks. 2- Response of citrus scions to soil type, rootstock and interstock. *Egypt J. Appl. Sci.*, 22(4B):1 – 24.
- Barakat, M.R., A.T. Mohsen, A.M. Abdel-El-Rahman and S.H. Hemeda (2013). Nutritional status and yield efficiency of Navel and Valencia orange trees as affected by used rootstocks. *J. Hort. Sci. & Ornament. Plants*, 5(2):137 – 144.
- Brown, J.D. and O. Lilleland (1974). Rapid determination of potassium and sodium in plant material and soil extracts by flame photometer. *Proc. Soc. Hort. Sci.*, 48: 341-346.
- Castle, W. (1983). Growth, yield and cold hardiness of seven year old 'Bears' lemon on twenty seven rootstocks. *Proc. Florida Sta. Hort. Soc.* 96: 23 – 25.
- Castle, W.S. (1992). Rootstock and interstock effects on the growth of young *Minneola* tangelo trees. *Proc. Florida Sta. Hort. Soc.* 105: 82 – 84.
- Castle, W.S. and R.L. Phillips (1980). Performance of March grape fruit and Valencia orange trees on 18 rootstocks in a closely spaced planting. *J. Am. Soc. Hort. Sc.*, 105:496 – 499.
- Castle, W.S., J.C. Baldwin and R.P. Muraro (2010). Performance of Valencia sweet orange trees on 12 rootstocks at two locations and an economic interpretation as a basis for rootstock selection. *HortScience*, 45(4):523 – 533.
- Chapman, H.D. and P.F. Pratt (1961). *Methods of Analysis for Soils, Plant and Waters*. Univ. of California, USA, pp. 169-170.
- Dawood, S. A. (2001). Growth, yield, fruit quality and leaf mineral content of Valencia orange trees on Sour orange and Volkamer lemon grown on slightly alkaline clay soil. *J. Agric. Res. Tanta Univ.*, 27 (4): 726 – 736.
- Dawood, S. A. (2002). Evaluation of Washington navel orange on Sour orange and Volkamer lemon grown on slightly alkaline clay soil conditions. *J. Agric. Res. Tanta Univ.*, 28 (1): 157 – 167.
- Ennab, H. A. (2003). Evaluation study on Washington navel orange cultivar budded on five rootstocks. Ph.D. Thesis, Fac. Agric. Kafr El-Sheikh, Tanta University.

- Gardner, F.E. (1968). The failure of rough lemon and sour orange interstocks to influence tree growth, yields and fruit quality of sweet orange varieties. *Proc. Fla. State Hort. Soc.*, 81:90 – 94.
- Georgiou, A. (2002). Evaluation of rootstocks for “Clementine” mandarin in Cyprus. *Scientia Hort.*, 93 : 29 – 38.
- Gimeno, V., J.P. Syvertsen, I. Simon, V. Martinez, J.M. Camara-Zapata, M. Nieves and F. Garcia-Sanchez (2012). Interstock of Valencia orange affects the flooding tolerance in Verna lemon trees. *HortScience* 47(3):403 – 409.
- Girardi, E.A. and F.A. Filho (2006). Production of interstocked Pera sweet orange nursery trees on Volkamer lemon and Swingle citrumelo rootstocks. *Sci. Agric. (Piracicaba, Braz.)*, 63(1):5 – 10.
- Hikal, A. R. F. (2014). Effect of different rootstocks on vegetative growth, fruiting, fruit quality and fruit storage on trees of Washington navel orange. *J. Plant Production, Mansoura Univ.*, 5(2):347 – 355.
- Jackson, M.L. (1967). *Soil Chemical and Analysis*. Prentice Hall of India, New Delhi, p. 498.
- Jahromi, A.A., H. Hasanzada and M.H. Farahi (2012). Effect of rootstock type and scion cultivar on citrus leaf total nitrogen. *World Applied Sciences Journal* 19 (1):140 – 143.
- Krezdorn, A.H. (1978). Interstocks for tree size control in citrus. *Proc. Fla. State Hort. Soc.* 91:50 – 52.
- Labanauskas, C.K. and W.P. Bitters (1974). Influence of rootstocks and interstocks on the macro- and micro-nutrients in Valencia orange leaves. *J. Amer. Soc. Hort. Sci.*, 99:32 – 33.
- Louzada, E.S., J.W. Grosser, F.G. Gmitter, Jr; Nielson, J.L. Chanler, X.X. Deng and N. Tusa (1992). Eight new somatic hybrid citrus rootstocks with potential for improved disease resistance. *Hort. Sci.*, 27(9):1033 -1036.
- Mansour, M.F.; A.E. Hassan and M.R. Rabeh (1993). Comparative study on leaf mineral contents and growth of navel orange scion in relation to different citrus rootstocks. *Minufiya J. Agric. Res.*, 18(1):443 – 452.
- Martinez, C., H. Lima and J. Rivas (1994). Growth and productivity of 5 types of Valencia orange on different rootstocks during the developmental phase. (*Hort. Abst.*, 64:710).
- Mohamed, M.A. (2011). Effect of some citrus rootstocks on vegetative growth and leaves mineral content of Balady lime transplants. M.Sc. Thesis, Fac. of Agric. Ain-Shams Univ., Cairo, Egypt.
- Moran, R. (1982). Formula for determination of chlorophyllous pigments extracted with N, N-dimethyl formamide. *Plant Physiol.*, 69: 1376-1381.
- Murphy, J. and J.R. Riley (1962). A modified single solution method for the determination of phosphorus in natural water. *Anal. Chem., Acta*, 27: 31-38.
- Perez-Zamora, O., V.M. Urrutia and S.B. Rodriguez (2002). Growth and yield of Valencia orange grafted on 16 citrus rootstocks growing in a calcimorphic soil and juice quality. *Articulo en Agrocienca* 36: 37 – 148.
- Pregl, F. (1945). *Quantitative organic microanalysis*, 4th ed J.A. Churchill, Ltd, London.
- Ramin, A.A. and A. Alirezanezhad (2005). Effects of citrus rootstocks on fruit yield and quality of Ruby Red and Marsh grapefruit. *Fruits*, 60 (5): 311 – 317.
- Roose, M.L., P.A. Cole, D.A. Tkin and R.S. Kupper (1989). Yield and tree size of four citrus cultivars on 21 rootstocks in California. *J. Am. Soc. Hort. Sci.*, 114:678 – 684.
- Sayed, R.A. and R.A. Abdel-Aziz (2010). Performance of some new citrus varieties under south El-Tahreer district conditions. *J. Plant Production, Mansoura Univ.*,1(2):291 – 300.
- Sayed, R.A. and S.S. Adawy (2009). New strain of Washington navel orange evaluation. *Egyptian J. Hort.*, 36(2):315 – 324.

Effect of rootstock and interstock on growth, yield and fruit quality

- Shafieizargar, A., Y. Awang, A. Juraimi and R. Othman (2012). Yield and fruit quality of Queen orange (*Citrus sinensis* (L) Osb.) grafted on different rootstocks in Iran. *Australian J. Crop Sci.*, 6(5):777 – 783.
- Shokrollah, H., T.L. Abdullah, K. Sijam and S.N. Abdullah (2011). Potential use of selected citrus rootstocks and interstocks against HLB disease in Malaysia. *Crop Protection* 30:521 – 525.
- Smith, M. W., R. G. Shaw, J. C. Chapman, J. Owen – Turner, L. Slade Lee, K. B. McRae, K. R. Jorgensen and W. V. Mungomery (2004). Long-term performance of “Ellendale” mandarin on seven commercial rootstocks in subtropical Australia. *Scientia Hort.*, 102: 75 – 89.
- Snedecor, G.W. and W.G. Cochran (1990). *Statistical methods*. 7th Ed. Iowa State Univ. Press. Ames., Iowa, USA.
- Toplu, C., M. Kaplankiran, T.H. Demirkaser and E. Yildiz (2008). The effects of citrus rootstocks on Valencia Late and Rhode Red Valencia oranges for some plant nutrient elements. *African Journal of Biotechnology* 7(24):4441 – 4445.
- Yılmaz, B., B. Cimen, M.U. Kamiloglu, M. Incesu, T. Yesiloglu and M. Yılmaz (2014). The influence of different interstock lengths of Minneola tangelo on photosynthetic parameters and fruit yield of Star Ruby grapefruit. *Turkish*
- Yılmaz, B., M.U. Kamiloglu, B. Cimen, M. Incesu, T. Yesiloglu and O. Tuzcu (2015). Effects of different interstock lengths on the yield, fruit quality and tree size of Kutdiken lemon trees in Turkey. *Journal of Global Agriculture and Ecology* 3(2):91 – 96.
- Zayan, M. A., S. M. Zeerban, H. M. Ayaad, S. A. Dawood and H. A. Ennab (2004). Evaluation study on Washington navel orange cultivar budded on five rootstocks. 1- Vegetative growth, root distribution and a ability to salt tolerance. *J. Agric. Res. Tanta Univ.*, 30 (2): 400 – 420.
- Zekri, M. (2000). Evaluation of orange trees budded on several rootstocks and planted at high density on Flatwoods soil. *Proc. Fla. State Hort. Soc.*, 113:119 – 123.

تأثير الاصل و الاصل الوسطى على النمو و المحصول وصفات الجودة لثمار بعض اصناف البرتقال.

أ- التأثير على النمو الخضري و الحالة الغذائية و المحصول.

سمية أحمد السيد

قسم الموالح - معهد بحوث البساتين - مركز البحوث الزراعية بالجيزة - مصر

الملخص العربي

اجريت هذه التجربة كدراسه طويله لأجل بدأت في 2008 - 2014 باستخدام النارج كأصل وسطى مع أصل الفولكامارينا وذلك لتفادى عيوب أصل الفولكامارينا على الخواص الطبيعه و الكيمائيه لثمار البرتقال مع أمل احتفاظه بمميزاته في قوه النمو و زياده المحصول وأجريت هذه التجربة بمزرعه خاصه بمنطقه البستان بحافظه البحيره بهدف دراسه تأثير أصل النارج كأصل وسطى بين أصل الفولكامارينا و الصنف المطعم عليه من أربعة أصناف من البرتقال (البرتقال الصيفى أوليندا - البرتقال ابو سره اسبرينج - البرتقال ابوسره بيرينت - البرتقال ابو سره فوكومتو) وذلك على النمو الخضري والحاله الغذائية للأشجار والمحصول والكفاءه المحصوليه . وقد اظهرت النتائج الاتى :

- 1- أعطت كل الأصناف المطعومه على أصل الفولكامارينا اعلى القيم لكل من حجم الشجره - و معظم قياسات النمو , وأعطى الأصل الوسطى قيم وسطيه لهذه القياسات وسجل أصل النارج أقل القيم كما سجل البرتقال الصيفى أوليندا أعلى قيم لحجم الشجره وقياسات النمو الخضري عند مقارنته بالأصناف الأخرى
- 2- زادت قيم كل من كلوروفيل أ , ب , الكلوروفيل الكلى فى الأصناف المطعومه على أصل الفولكامارينا ولم يكن هناك فروق معنويه بينه و بين الأصناف على الأصل الوسطى وكانت أعلى القيم على صنف ابوسره اسبرينج المطعوم على أصل الفولكامارينا.
- 3- احتوت اوراق الأصناف المطعومه على أصل الفولكامارينا أعلى قيم من NPK يليه الأصناف التى على الأصل الوسطى ثم النارج فى كلا الموسمين واعطت الاصناف اولندا الصيفى وابوسره بيرينت على اصل الفولكامارينا أعلى القيم مقارنه بالاصناف الأخرى .
- 4- أعطت الاصناف على اصل الفولكامارينا اعلى قيم للمحصول ممثل بالكيلو جرام لكل شجره بفروق معنويه بينه وبين الاصول الأخرى يليه أصل النارج ثم الاصل الوسطى بدون فروق معنويه بينهما فى كلا الموسمين كما سجل صنف البرتقال الصيفى اولندا أعلى القيم للمحصول مقارنه بالاصناف الأخرى بفرق معنويه فى معظم الحالات.
- 5- اظهرت النتائج ان الأصناف المطعومه على أصل النارج أعطت أعلى كفاءه محصوليه ممثله بكيلوجرام/سم² من مساحه مقطع الجذع (Kg /cm² of TCSA) بينما اعطت الأصناف على اصل الفولكامارينا اعلى قيم للكفاءه المحصوليه الممثله بالكيلو جرام لكل متر مكعب من حجم الشجره (Kg/m³ /canopy volume) يليه اصل النارج ثم الأصل الوسطى فى كلا الموسمين بدون فروق معنويه بينهما ولم يكن هناك فروق معنويه بين الأصناف. تبين أن استخدام النارج كأصل وسطى لم ينقص من محصول الأشجار المطعومه عليه نقصا معنويا وكان المحصول قريبا من الناتج عن أصل النارج بمفرده.

Effect of rootstock and interstock on growth, yield and fruit quality