

## Antimicrobial Activity of Some Plant Essential Oils Compared with Propolis

EL- Refai, A. A.\*; Amalika D. EL-Dahshan\*; M. Abou-Taleb\*\* and Fify R. Anees\*\*

\* Food Industries Dept., Fac. of Agric., Mansoura Univ., Egypt.

\*\*Fish Processing and Technology Laboratory, National Institute of Oceanography and Fisheries, Academy of Scientific Research, Cairo, Egypt.



### ABSTRACT

Antimicrobial activity of some plant essential oils namely, rosemary (*Rosmarinus officinalis*), ginger (*Zingiber officinale Roscoe*), red pepper (*Capsicum annum L.*), marjoram (*Origanum marjorana L.*) oils and propolis were studied. About 24 different phenolic compounds were identified by HPLC which have the ability to act as antimicrobial agents. Results indicated that all examined oils and propolis were varied in their content of phenolic compounds. These oils and propolis were evaluated for their antimicrobial activities against four pathogenic bacteria strains (*Staph aureus*, *Listeria monocytogenes*, *Salmonella sp* and *E. coli*), two strains of fungi (*A. flavus* and *A. niger*) and yeast strain (*Rhodotorula sp.*). Results indicated that marjoram oil had inhibitory effect against all tested strains of bacteria, fungi and yeast. Ginger and rosemary oils affected strains of *Salmonella sp* and *Listeria monocytogenes*. Also, propolis had inhibitory effect against all tested strains of bacteria, fungi and yeast. In contrast, the red pepper essential oil showed no inhibition effect against any tested strain. Obtained results suggest the possibility to use the studied plant oils and propolis to control pathogens for food preservation.

**Keywords:** Antimicrobial, essential oils, rosemary, ginger, marjoram, red pepper and propolis.

### INTRODUCTION

Food-borne diseases are considering a great problem all over world, even in advanced nations (Mead *et al.*, 1999). An assortment of microorganisms also causes food deterioration which is one of the most significant concerns of the food technology. A variety of pathogenic microorganisms such as *Escherichia coli*, *staphylococcus aureus* and *Listeria monocytogenes* have been reported as the causal of food-borne disease and food deterioration (Deak and Beuchat 1996 and Betts *et al.*, 1999).

In last years a great excess in the number of registered cases of food-borne disease has been observed, with 82041 registered cases in 1995 (Monitor 1996). So, there is a great attention in methods to stop this upward trend and decrease the occurrence of food poisoning. An important part of research is the improvement of new and developed procedure of food preservation. Plant extracts, including their essential oils are well established that these substances have antimicrobial effect against of bacteria, moulds and yeasts (Pai and Paltt 1995). However, it is only comparatively newly that great concern has been offered to possibility of application as food preservative substances (Smith-Plmer *et al.*, 1998).

In food technology, the shelf life of food products is generally extended by the supplement of antibacterial chemical substances to these products. However, there is a necessity for natural preservative additives, which do not change the sensory characteristics of food products. Several of the essential oils are considered very good candidate for achievement that object to prevent food deterioration (Schelz *et al.*, 2010).

Rosemary (*Rosmarinus officinalis L.*) originally grows in southern Europe. Its herb and oil are commonly used as spice and flavoring agents in food processing for its desirable flavor, high antioxidant activity and lately as antimicrobial agent (Ouattara *et al.*, 1997; Lo *et al.*, 2002). It is a complex botanical, containing over 240 medical and nutritional active compounds. Rosemary volatile oil is characterized by

containing a high percentage of terpenoid oxygenated compounds like L-penene and 1.8-cineole as the major compounds, which are mainly responsible for their aromatic profiles (Celiktas, *et al.*, 2007 and El-Bastawesy, *et al.*, 2008).

Ginger (*Zingiber officinale*, *Zingiberaceae*) has widely been consumed as spices and food preservative. It is added to food products in a form of essential oils and various extracts (Yu *et al.*, 2007).

Capsicum (red pepper) has its beginning since the beginning of civilizations. It is a part of human diet since 7500 BC (Bosland, 1996). Dorantes *et al.*, (2000) have reported the inhibitory effect of pathogens by capsicum annum extract. Also, the inhibition of *salmonella typhimurium* and *Pseudomonas aeruginosa* inoculated in minced beef meat by extract of *capsicum annum* bell pepper has been recorded (Careaga *et al.*, 2003).

Among several essential oils that may be useful as antimicrobial agents, marjoram oil (*Origanum majorana L.*) may have the greatest potential for use in industrial applications (Daferera *et al.*, 2000; Ezzeddine *et al.*, 2001).

Propolis is a rubbery and resinous substance produced by worker bees from buds of flowers, coat of trees and leave of plants. Bees used the enzymatic secretion-enriched substance to coat hive walls to secure a very clean environment. Propolis extract as a natural product have been used both inwardly and outwardly for thousands of the years as one of the cure factors in conventional medicine. The biological characteristics of propolis such as antiviral, antibacterial, antifungal and other properties have attracted the researcher's attention (Simone-Finstrom and Spivak, 2020). The precise chemical composition of propolis is complex influenced by the phytogeographical origin, the season of collection, and the type of bees foraging (Murad *et al.*, 2002; Bankova, 2005 and Silici and Kutluca, 2005).

In general, propolis contains a variety of chemical compounds such as polyphenols (flavonoid aglycones, phenolic acids, and their esters, phenolic aldehydes, alcohols, and ketones), terpenoids, steroids,

amino acids, and inorganic compounds (Moreno *et al.*, 2000 and Osman, 2012).

The aim of this research is testing the antimicrobial activity of four plant essential oils against four prevalent food-borne pathogens (*Salmonella sp*, *Listeria monocytogenes*, *Staphylococcus aureus* and *Escherichia coli*), two strains of fungi (*A. flavus* and *A. niger*) and yeast strain (*Rhodotorula sp.*).

## MATERIALS AND METHODS

### Materials:

#### Essential oils:

Four plant essential oils namely; rosemary, ginger, marjoram and red papper oils were obtained from Medical Plant and Agriculture Seeds Haraz Company. Cairo, Egypt.

#### Propolis:

Propolis was obtained from Egyptian Company for Production and Marketing of honey bee products. Cairo, Egypt.

#### Microorganisms:

Four pathogenic bacterial strains namely, *Staphylococcus aureus*, *Listeria monocytogenes*, *Salmonella sp* and *E. coli* (O157:H7), two fungi strains (*A. flavus* and *A. niger*) and one yeast strain (*Rhodotorula sp*) were obtained from Food Hygiene Inspection Lab., Damietta Seaport. These strains were chosen for their economic importance regarding the human health.

#### Media:

Tryptone soya agar (Oxoid, 2006, CM0131). Sabouraud dextrose agar (Oxoid, 2006, CM0041).

#### Methods:

#### Chemical analysis:

#### Extraction and Identification of phenolic compounds:

Extraction of phenolic compounds of rosemary (*Rosmarinus officinalis*), ginger (*Zinger officinale* Roscoe), red pepper (*Capsicum annum L.*) marjoram (*Origanum marjorana L.*) oils and propolis, was carried out according to the method described by Wojdylo *et al.*, (2007), then determined according to the method described by Waskmundzka *et al.*, (2007), which

calculated as mg Gallic acid/100g of dry weight material. Phenolic compounds of studied oils were identified using High Performance Liquid Chromatography (HPLC), "HP1050" in Center Laboratory of Food Technology Research Institute, Giza, Egypt.

#### Preparation of propolis extract:

Ethanolic extract of propolis (EEP) which used in holes (in the plate diffusion method) was obtained by diluting 25 g crude propolis in 100 ml of 70% ethanol, and extracted at room temperature. After three days the extract was filtered (Whatman paper No. 1) and kept at refrigerator temperature at 4 °C.

#### Microbiological experiments:

#### Assessment of antimicrobial activities:

The plate diffusion method was used, holes with a cork borer were pushed in medium inoculated with a standard incolumn of 0.5 ml cell suspension of bacteria using vortex mixer (No 502550, Taiwan) under aseptic conditions. These cultivation media were tryptone soya agar for bacteria and sabouraud dextrose agar for both fungi and yeast. Each hole was filled with 10µl of the oil samples and propolis then left one hour to allow diffusion, the plates of *Salmonella sp*, *E. coli* and *Staphylococcus aureus* bacteria were incubated at 37 °C for 24-48 h. The plates of *Listeria monocytogenes* were incubated at 30 °C for 24-48 h. While, plates of fungi strains (*A. flavus* and *A. niger*) and yeast strain (*Rhodotorula sp*) were incubated at 25 °C for 4-5 days. At the end of incubation period, the inhibition zones of microbial growth were measured by ruler (as mm) and recorded (Bagamboula *et al.*, 2003).

## RESULTS AND DISCUSSION

#### Extraction and identification of phenolic compounds in plant essential oils and propolis:

Data in Table (1) show the identification of different phenolic compounds of investigated plants oils. The obtained results indicate that phenolic acids were the most abundant compounds in all examined oils.

**Table (1): Identification of phenolic compounds in plant oils compared with propolis (mg/100gm)**

Phenolic compounds (mg/100gm)	Essential oils					propolis
	Rosemary	Ginger	Red pepper	Marjoram		
Gallic acid	1.56	0.60	0.61	0.60		ND
Chlorogenic	0.90	ND	1.05	0.32		5.52
Pyrogallol	ND	ND	3.32	ND		39.75
Protocatechuic	ND	ND	0.05	ND		1.72
Catechol	ND	0.19	ND	ND		21.07
Catechin	0.77	0.78	0.83	1.74		ND
Ferulic	1.05	4.17	0.41	ND		3.74
Caffeic	ND	0.36	0.53	1.01		ND
Vanillic	ND	ND	2.31	ND		1.46
P.Cumaric	ND	0.06	0.42	ND		ND
Cholchecien	ND	1.02	ND	ND		ND
Caffeine	ND	ND	1.79	ND		ND
Salicylic	2.65	1.09	ND	54.1		ND
Chrysin	1.05	0.32	0.07	ND		287.28
Cumarin	ND	ND	ND	ND		2.03
Cinnamic	ND	ND	ND	ND		142.59
Benzoic	ND	ND	ND	ND		5.45

ND: Non Detected

Results of HPLC indicate that fourteen phenolic compounds were separated, six of them were identified salicylic acid was the major phenolic compound in rosemary (2.65 mg/100gm) followed by gallic (1.56 mg/100gm) then both ferulic and chrysin (1.05 mg/100gm). More ever, chlorogenic and catechin also were detected (0.90 and 0.77 mg/100gm).

In ginger oil the main phenolic compound was ferulic acid (4.17 mg/100gm), followed by salicylic (1.09 mg/100gm) and cholchecien (1.02 mg/100gm), while there were lower amounts of catechin (0.78 mg/100gm), gallic (0.60 mg/100gm), caffeic (0.36 mg/100gm), chrysin (0.32 mg/100gm), catechol (0.19 mg/100gm) and p.cumaric (0.06 mg/100gm).

The phenolic compounds of red pepper oil were pyrogallol (3.32 mg/100gm), vanillic (2.31 mg/100gm), caffeine (1.79 mg/100gm), chlorogenic (1.05 mg/100gm), catechin (0.83 mg/100gm), gallic (0.61 mg/100gm), caffeic (0.53 mg/100gm), p.cumaric (0.42 mg/100gm), ferulic (0.41 mg/100gm), chrysin (0.07 mg/100gm) and protocatechuic (0.05 mg/100gm).

Marjoram oil is used also as natural flavoring agent, nearly five phenolic compound were identified. Data in the same table indicate that salicylic was the predominant phenolic compound (54.1 mg/100gm) followed by catechin (1.74 mg/100gm) and caffeic (1.01 mg/100gm). However, there were lower amounts of gallic and chlorogenic (0.60 and 0.32 mg/100gm, respectively).

The phenolic compounds of propolis are shown in Table (1). According to the results of HPLC analysis, the propolis sample contains about 10 identified phenolic compounds. Among these phenolic compounds chrysin (287.28 mg/100gm) followed by cinnamic (142.59 mg/100gm), pyrogallol (39.75 mg/100gm) and

catechol (21.07). However, there were lower amount of protocatechuic (1.72 mg/100gm), chlorogenic (5.52 mg/100gm) and vanillic (1.46 mg/100gm).

**Antimicrobial activity of plant essential oils and propolis:**

The antimicrobial effect of the tested plant essential oils and propolis was assessed against *Staph aureus*, *Listeria monocytogenes*, *Salmonella sp* and *E. coli*, *A. flavus*, *A. niger* and *Rhodotorula sp* by diffusion method. Results are presented in Table (2). The results show the wide variation in the antimicrobial effect of the antimicrobial effect of the plant essential oils. From the microbial examination, it could be reported that ginger and rosemary essential oils had an effect against tested strains of *Salmonella sp* and *Listeria monocytogenes*.

The essential oil of red pepper as revealed in the present study showed no effects against gram-negative, gram-positive bacteria, fungi and yeast (no inhibition zone) comparing with those of other oils such as marjoram oil and propolis. These results are similar with those reported by Dorman and Deans (2000) and in contrary with those reported by Dorantes *et al.*, (2000) and Careaga *et al.*, (2003), which they found that there was an inhibition effect caused by the five capsainoids. It was observed that m-cumaric and cinnamic acids are responsible for the inhibitory effect of the bacteria. It can be seen that the inhibition effect against *Salmonella typhimurium* is peripheral but *Bacillus cereus* and *Staphylococcus aureus* were the most sensitive to these substances. Both of bacteria showing an inhibition zone of 10 and 9.8 mm respectively, for m-cumaric acid and 8.2 and 6 mm respectively, for cinnamic acid. Also, there was strong inhibition of *Listeria monocytogenes* by m-cumaric acid (6.2 mm) and cinnamic acid (5mm).

**Table (2): Antimicrobial activity of some plant oils and propolis (inhibition zone/mm)**

Essential oils and propolis (10 µl)	Examined Microorganisms						
	Gram negative bacteria (G -)		Gram positive bacteria (G+)		Fungi		Yeast
	<i>Salmonella sp</i>	<i>E.coli</i>	<i>Staph aureus</i>	<i>Listeria monocytogenes</i>	<i>A. flavus</i>	<i>A. niger</i>	<i>Rhodotorula sp</i>
Rosemary oil	11.50	NA	NA	15.00	NA	NA	NA
Ginger oil	10.00	NA	NA	19.50	NA	NA	NA
Red pepper oil	NA	NA	NA	NA	NA	NA	NA
Marjoram oil	15.75	15.00	12.5	16.75	13.50	13.50	17.5
Propolis	9.50	9.81	14.3	10.20	15.40	15.60	16.20

NA= non active

G -= gram negative cells

G += gram positive cells

Marjoram essential oil resulted in zones of 15.75, 15, 12.5 and 16.75 mm against gram negative bacteria *salmonella sp* and *E. coli* , gram positive bacteria *Staph aureus*, *Listeria monocytogenes*, respectively, and 13.5 , 13.5 and 17.5 against *A. flavus*, *A. niger* and *Rhodotorula sp*. So, such results indicate that marjoram oil had the superior compared with other oils. These results may be due to that marjoram oil was rich in some phenolic compounds namely salicylic, caffeic, catechin and gallic acid. Dwivedi (1990) reported that complete growth inhibition of *F. oxysporum f.sp.psidii*

at 0.1% and 0.15% concentration of phenol and 0.15% concentration of salicylic acid due to strong toxicity of these compound against pathogen. Also, gallic acid has been shown to possess antibacterial activity against human pathogens (*Staphylococcus aureus*, *Corynebacterium accolans*), a plant pathogen (Fogliani *et al.*, 2005). These results are in accordance with those given by Govaris *et al.*, (2010) and El-Gammal and Omar (2012), which they mentioned that antimicrobial effect of marjoram oil constitutes were found to be phenols, alcohols, aldehydes, ketones, ethers and

hydrocarbons can increase the antimicrobial activity and their bacteriostatic effect depending on their effectiveness concentration.

Marjoram essential oil has been tested as a natural preservative substance in food processing. Antimicrobial effects of marjoram essential oil occurs via cell membrane destroy, with envelop disruption, formation of bubble and lack of cytoplasmic substance. These changes correlate with the ability of hydrocarbons to react with hydrophobic structures, like bacterial membranes. These observations are based on examination of *E. coli* and *S. aureus* by transmission electron microscopy, which allow to studying the possible bacterial ultrastructural modifications Becerril *et al.*, (2007).

Shan *et al.*, (2007) studied a series of dietary spices and herbal medicines in order to confirm their antibacterial effects against pathogenic bacteria, of which infections are responsible for food toxicity. The results showed that phenolic compounds significantly participated to the antibacterial activity of the investigated herbs. This antibacterial activity may be due to the enzyme inhibition by the more oxidized phenolic compounds probably via interaction sulfhydryl compounds or via more non-specific reaction with the protein. Caffeic acid and cinnamic acid, pyrogallol and catechol were shown to be toxic to microorganisms (Cowan, 1999).

Results also, indicated that marjoram oil have also high levels of antifungal activity, it has been demonstrated that antimicrobial effect of volatile oils structural and functional damages to the bacterial cell membrane, it also, indicated that the optimum range of hydrophobicity is involved the toxicity of volatile oils (Goni *et al.*, 2009).

A lot of investigations reporting the antibacterial effect of volatile oils against pathogenic bacteria accept that volatile oils are comparatively more active against Gram (+ve) than Gram (-ve) bacteria (Viuda-Martos *et al.*, 2008). The cell structure of Gram (-ve) bacteria is constituted essentially with lipopolysaccharides. This structure averts the collection of oils on the membrane of bacterial cell (Bezic *et al.*, 2003).

Also, the results of propolis effect as antimicrobial agent were shown in table (2). These results revealed that propolis had strong antibacterial activity against Gram (+ve) than Gram (-ve) bacteria. The largest inhibition zone was recorded against *Staphylococcus aureus* (by 14.3 mm). From these results it was observed that propolis results were nearly in accordance with that obtained by (Park *et al.*, 2005), who reported that Hungarian propolis had greater inhibitory effect on growth of resistant Gram (+ve) bacteria and fungi than Gram (-ve) bacteria. Abd El Hady and Hegazi (2002) reported that *Staphylococcus aureus* were sensitive to Propolis which is in agreement with the results of the present study. Furthermore, the result of disc diffusion methods of propolis revealed that, *Staphylococcus aureus* was highly sensitive with 13 mm as zone of inhibition. In contrary with our results, the antibacterial activity of propolis was generally investigated on *Escherichia coli* by Hendi *et*

*al.*, (2011). The results confirmed a very inhibitory activity of propolis on the growth of *Escherichia coli* and these results were in complete accordance with the results of Sforcin *et al.*, (2000), Drago *et al.*, (2000). As a basic rule, an extract substance is conceder active against both bacteria and fungi if the zone of inhibition was greater than 6 mm (Muhammad and Muhammad, 2005). From otherwise, different researchers (Katircio and Nazime 2006; Yaghoubi. *et al.*, 2007) have reported that propolis antibacterial activity is due to a number of phenolic compounds, mainly flavonoids, phenolic acids and their esters. Also, Kosalec *et al.*, (2004) reported that, propolis and some of its cinnamic acid derivatives and flavonoids were responsible for uncoupling the energy transuding cytoplasmic membrane inhibiting bacterial motility, which might contribute to the antibacterial action.

Finally it could be concluded that the effective essential oils and propolis could be a natural source of reducing the total contamination level of foods beside on their effect against food pathogen microorganisms.

The obtained results suggest that, marjoram oil and propolis have a strong antimicrobial activity.

## REFERENCES

- Abd El Hady, F. K. and Hegazi, A. G. (2002). Egyptian Propolis: 2. chemical composition, antiviral and antimicrobial activities of east Nile Delta propolis. *Z. Ntaureforsch.* 57: 386 -394.
- Bagamboula, C. F.; Uyttendaele, M. and Debevere, J. (2003). Antimicrobial effect of spices and herbs on *Shigella sonnei* and *shigilla flexneri*. *J. Food Protection*, 66(4): 668-673.
- Bankova, V. (2005). Recent trends and important developments in propolis research. *Evid Based Compl Altern Med.*, 2: 29-32.
- Becerril, R.; Gomez-Lus, R.; Goni, P.; Lopez, P., and Nerin, C. (2007). Combination of analytical and microbiological techniques to study the antimicrobial activity of a new active food packaging containing cinnamon or oregano against *E. coli* and *S. aureus* *Anal. Bioanal. Chem.*, 388: 1003-1011.
- Betts, G. D.; Linton, P. and Betteridge, R. J. (1999). Food spoilage yeasts: effects of pH, NaCl and temperature on growth. *Food Control*, 10, 27-33.
- Bezic, N.; Skocibusic, M. V. and Radonic, A. (2003). Composition and antimicrobial activity of *Achillea clavennae* L. essential oil. *Phototherapy Research*, 17, 1037-1040.
- Bosland, P. W. (1996). Capsicums: Innovative uses of an ancient crop. In: J. Janick (Ed.), *Progress in New Crops*. ASHS Press; Arlington, VA. p 479-487.
- Careaga, M.; Fernandez, E.; Dorantes, L.; Mota, L.; Jaramillo, M. E. and Hernandez-Sanchez H. (2003). Antibacterial activity of Capsicum extract against *Salmonella/ typhimurium* and *Pseudomonas aeruginosa* inoculated in raw beef meat. *Int. J. Food Microbiol.* 83(3): 331-335.

- Celiktas, O. Y.; Bedir, E. and Sukan, F. V. (2007). In vitro antioxidant activities of *Rosemarinus officinalis* extracts treated with supercritical carbon dioxide. *Food Chem.*, 101(4): 1457-1464.
- Cowan, M. M. (1999). Plant products as antimicrobial agents. *Clinical Microbiology Reviews*, 12: 564-582.
- Daferera, D. J.; Ziogas, B. N. and Polissiou, M. G., (2000). GC-MS analysis of essential oils from some Greek aromatic plants and their fungitoxicity on *Penicillium digitatum*. *J. Agric. Food Chem.* 48, 2576-2581.
- Deak, T. and Beuchat, L. R. (1996). Handbook of Food Spoilage; CRC Press: New York, USA.
- Dorantes, L.; Colmenero, R.; Hernandez, H.; Mota, L.; Jaramillo, M. E.; Fernandez, E. and Solano, C. (2000). Inhibition of growth of some foodborne pathogenic bacteria by *Capsicum annum* extracts. *Int. J. Food Microbiol.* 57(1-2): 125-128.
- Dorman, H. J. D. and Deans, S. G. (2000). Antimicrobial agents from plants: antibacterial activity of plant volatile oils. *J. App. Microbiol.* 88 308-316.
- Drago, L.; Mombelli, B.; De Vecchi, E.; Fassina, M. C.; Tocalli, L.; and Gismondo, M. R. (2000). In vitro antimicrobial activity of propolis dry extract. *J. Chemother.* 12 (5), 390 - 395.
- Editor: Deprasasad Chattopadhyay.
- Dwivedi, S. K. (1990). Antifungal activity of some phenolic compounds on *Fusarium oxysporum f.sp.psidii* causing guava wilt. *Hindustan Antibiotic Bull.* 32(1&2): 33-35.
- El-Bastawesy, Amal M., Ragaa, H. and El-Refai, A. A. (2008). Chemical and biological evaluation of rosemary (*Rosmarinus officinalis* L.) leaves volatile oil and its methanolic extract. 4<sup>th</sup> Arab. Mans. Conf. of Food & Dairy Sci. & Tec. November, 81-100.
- El-Gammal, Rania E. and Omar, Sabreen A. (2012). Antioxidant and antibacterial activity of some herbs volatile oils and their ethanolic extracts. *J. Food and Dairy Sci., Mansoura Univ.*, 3(1):54-61.
- Ezzeddine, N. B.; Abdelkefi, M. M.; Ben-Aissa, R. and Chaabouni, M. M. (2001). Antibacterial screening of *origanum majorana* L. oil from Tunisia. *J. Essent. Oil Res.* 13, 295-297.
- Fogliani, B.; Raharivelomanana, P.; Bianchini, J. P.; Madjebi, S. B. and Hnawia, R., (2005). Bioactive ellagitannins from *Cunonia macrophylla*, an endemic Cunoniaceae from New Caledonia. *Phytochem.*, 66, 241-247.
- Goni, P., Lopez, P., Sanchez, C., Gomez-Lus, R., Becerril, R. and Nerin, C. (2009). Antimicrobial activity in vapour phase of combination of cinnamon and clove essential oils. *Food Chemistry*, 116, 982-989.
- Hendi Nada, K. K.; Habeeb, S. Naher and Alaa H. Al-Charrakh (2011). In vitro antibacterial and antifungal activity of Iraqi propolis. *Journal of Medicinal Plants Research* Vol. 5(20), pp. 5058-5066.
- Katircio, H. and Nazime, M. (2006). Antimicrobial activity and chemical compositions of Turkish propolis from different regions. *Afr J Biotechnol.* 5: 1151-1153.
- Kosalec, I.; Bakmaz, M.; Pepeejnjak, S. and Kneževi, S. V. (2004). Quantitative analysis of the flavonoids in raw propolis from northern Croatia. *Acta Pharmaceut.* 54: 65-72.
- Lo, A. H.; Liang, Y. C.; Lin-Shiau, S. Y.; Ho, C. T. and Lin, J. K. (2002). Carnosol, an antioxidant in rosemary, suppresses inducible nitric oxide synthase through down-regulating nuclear factor- $\kappa$ B in mouse macrophages. *Carcinogenesis*, 23: 983-991.
- Mead, P. S.; Slutsker, L.; Dietz, V.; McCiag, L. F.; Breese, J. S.; Shapiro, C.; Griffin, P. M. and Tauxe, R. V. (1999). Food related illness and death in the United States. *Emerg. Infect. Dis.* 5, 607-625.
- Muhammad, H. and Muhammad, S. (2005). The use of *Lawsonia inermis* Linn. (Henna) in the management of burn wound infections. *Afr. J Biotechnol.* 4(9): 934-937.
- Monitor (1996). Population and Health. National Statistics. Government Statistical Service. London: Office for National Statistics.
- Moreno, M. I.; Isla, M. I.; Sampietro, A. R. and Vattuone, M. A. (2000). Comparison of the free radical-scavenging activity of propolis from several regions of Argentina. *Journal of Ethnopharmacology*, 71: 109-114.
- Murad, J. M.; Calvi, S. A.; Soares, A. M.; Bankovea, V. and Sfrocin, J. M. (2002). Effect of propolis from Brazil and Bulgaria on fungicidal activity of macrophages against *paracoccidioides brasiliensis*. *Journal of Ethnopharmacology*, 79: 331-334.
- Osman, M. F. (2012). Chemical composition and antimicrobial activity of some Egyptian propolis collected from different regions in Kafrelsheikh governorate. 6<sup>th</sup> Arab. Mans. Con. Food & Dairy Sci. November. 141-149.
- Ouattara, B.; Simard, R. E.; Holley, R. A.; Piette, G. J. P. and Begin, A. (1997). Antibacterial activity of selected fatty acids and essential oils against six meat spoilage organisms. *Int. J. Food Microbiol.*, 37:155-162.
- Pai, S. T. and Platt, M. W. (1995). Antifungal effects of *Allium sativum* (garlic) extract against the *Aspergillus* species involved in otomycosis. *Letters in Applied Microbiology* 20, 14-18.
- Park, Y. K.; Alencar, S. M. and Aguiar, C. L. (2005). Botanical Origin and Chemical Composition of Brazilian Propolis. *J. Agric. Food Chem.* 50: 2502-2506.
- Schelz, Z.; Hohmann, J. and Molnar, J. (2010). Recent advances in research of antimicrobial effects of essential oils and plant derived compounds on bacteria. *Ethnomedicine: A Source of Complementary Therapeutics*: 179-201 ISBN: 978-81-308-0390-6.

- Sforcin, J. M.; Fernandes, A.; Lopes, C. A. M.; Bankova, V. and Funari, S. R. C. (2000). Seasonal effect on Brazilian propolis antibacterial activity. *Journal of Ethnopharmacology*. Volume 73, Issues 1–2, Pages 243–249.
- Shan, B.; Cai, Y. Z.; Brooks, J. D. and Corke, H. (2007). *Int. J. Food. Microbiol.* In Press.
- Silici, S. and Kutluca, S. (2005). Chemical composition and antibacterial activity of propolis collected by three different races of honeybees in the same region. *J Ethnopharmacol.*, 99: 69-73.
- Simone-Finstrom, M. and Spivak, M. (2010). Propolis and bee health: the natural history and significance of resin use by honey bees. *Apidologie*, 41: 295-311.
- Smith-Plamer, A.; Stewart, J. and Fyfe, L. (1998). Antimicrobial properties of plant essential oils and essences against five important food-borne pathogens. *Letters in Applied Microbiol.* 26, 118-122.
- Viuda-Martos, M.; Ruiz-Navajas, Y.; Fernández-López, J. and Perez-Álvarez, J. A. (2008). Antibacterial activity of lemon (*Citrus lemon L.*), mandarin (*Citrus reticulata L.*), grapefruit (*Citrus paradise L.*) and orange (*Citrus sinensis L.*) essential oils. *Journal of food safety*, 28, 567-576.
- Waskmundzka, M.; Wianowska, D.; Szewczyk, K. and Oniszczyk, A. (2007). Effect of sample preparation methods on HPLC quantitation of some phenolic acids in plant materials. *Acta Chromatographic (19)*: 227-237.
- Wojdylo A.; Oszmianski J. and Czemerys R. (2007). Antioxidant activity and phenolic compounds in 32 selected herbs. *Food Chemistry*, 105: 490-494.
- Yaghoubi, S. M. J.; Ghorbani, G. R.; Soleimani, Zad, S. and Satari, R. (2007). Antimicrobial activity of Iranian propolis and its chemical composition. *DARU*, 15: 45-48.
- Yu, H. S.; Lee, S. Y. and Jang, C. G. (2007). Involvement of 5-HT1A and GABAA receptors in the anxiolytic-like effects of *Cinnamomum cassia* in mice. *Pharmacol. Biochem. Behav.*, 87: 164-170.

**الخواص المضادة للنشاط الميكروبي لبعض الزيوت النباتية العطرية بالمقارنة مع البروبوليس**  
**أحمد عبدالعزيز الرفاعي\*، أماليكا درويش الدهشان\*، محمد أبو طالب\*\* و فيفي راغب أنيس محمود\*\***  
**\*قسم الصناعات الغذائية - كلية الزراعة - جامعة المنصورة - مصر.**  
**\*\*معمل تكنولوجيا تصنيع الأسماك - المعهد القومي لعلم البحار والمصايد - أكاديمية البحث العلمي - مصر.**

باستخدام جهاز الفصل الكروماتوجرافي (HPLC) تم تقدير المواد الفينولية للزيوت المستخلصة من بعض النباتات وهي حصى اللبان، الزنجبيل، الفلفل الأحمر، البردقوش و البروبوليس وكانت حوالي ٢٤ مركب والتي لها القدرة أن تعمل كعوامل مضادة للميكروبات. وأظهرت النتائج ان الزيوت التي تم تحليلها وكذلك البروبوليس كانت مختلفة في محتواه من المركبات الفينولية. كما تم تقدير النشاط الميكروبي لهذه الزيوت النباتية الطيارة وكذلك النشاط الميكروبي للبروبوليس على أربعة سلالات من البكتيريا (*A. flavus*, *A. niger*) و سلالتين من الفطر (*Staph aureus*, *Listeria monocytogenes*, *Salmonella sp* and *E.coli*) وسلالة من الخميرة (*Rhodotorula sp*) وقد أظهرت النتائج ان كلا من زيت البردقوش أعطى أعلى فاعلية تجاه جميع سلالات البكتيريا والفطريات والخميرة المختبرة تلاه زيت الزن+9جيبيل وحصى اللبان حيث كان له تأثير على بكتريا *Listeria monocytogenes*, *Salmonella sp* كما أظهر البروبوليس تأثير واضح على جميع السلالات الميكروبية المختبرة وعلى العكس فلم يظهر الفلفل الأحمر أي تأثير على أي من السلالات المختبرة سواء كانت البكتيريا أو الفطريات أو الخميرة لذلك نقترح من النتائج التي تم الوصول إليها إمكانية استخدام الزيوت النباتية الطيارة محل الدراسة التي أظهرت تأثير على السلالات المختبرة وكذلك البروبوليس في مجال حفظ الأغذية والحد من نمو الميكروبات الممرضة للإنسان.