

## Effect of Some Fertilization and Micro-Nutrients Treatments on Growth and Chemical Constituents of *Echinacea purpurea* plant

Ghata, Y.A.A. \* and Wafaa H. Abdallah\*\*

\* Horticulture Department, Faculty of Agriculture, Moshtohor, Benha University, Egypt.

\*\* Medicinal and Aromatic Plants Department, Desert Research Center, Egypt.



### ABSTRACT

Two field experiments were carried out at special Farm at El Khatatba region Minofia Governorate , Egypt and in the Laboratory of Horticulture Department, Faculty of Agriculture at Moshtohor, Benha Univ., during 2014 and 2015 seasons to evaluate the effect of some fertilization treatments i.e. control (compost at 10 ton/fed.), 100 % full chemical fertilizer dose, 75% chemical fertilizer (NPK) dose + compost at 10 ton/fed. + bio fertilizer (nitrobein + phosphorein) , 50% chemical fertilizer (NPK) dose + organic fertilizer (compost) + bio fertilizer (nitrobein + phosphorein) and 25% chemical fertilizer (NPK) dose + organic fertilizer (compost) + bio fertilizer(nitrobein + phosphorein) and some micro-nutrients i.e. Zn and B each at 100 ppm as well as their combinations on growth, total caffeic acid derivatives and total alkalamides as well as chemical constituents of *Echinacea purpurea* plants. Results showed that different tested treatments of fertilization and micro-nutrients treatments as well as their combination led to significant increase of the studied growth parameters i.e. plant height, number of branches, fresh and dry weights of herb/plant, number of suckers/plant, number of flowering heads/plant, fresh and dry weights of flowering heads/plant as compared with control plants in both seasons. Additionally, the enhanced growth of *Echinacea purpurea* plants due to the different treatments was accompanied by pronounced increase in leaves N, P and K content of treated plants in both seasons. Furthermore, total caffeic acid derivatives and total alkalamides were increased by all fertilization and micro-nutrients treatments as well as their combinations, especially the combined treatment between 75% chemical fertilizer dose + compost at 10 ton/fed. + bio fertilizer and Zn or B each at 100 ppm. Conclusively, from the obtained results, it is preferable to fertilize *Echinacea purpurea* plants with 75% chemical fertilizer (NPK) + 10 ton compost/fed. + bio fertilizer(nitrobein + phosphorein) in combination with some micro-nutrients (Zn or B each at 100 ppm) as foliar spray to enhance plant growth and chemical composition which led finally to a safe product of high quality suitable for exportation and safe on human health

**Keywords.** *Echinacea purpurea*, chemical, organic and bio fertilization, micro-nutrients, growth , total caffeic acid derivatives and alkalamides contents.

**Corresponding Author:** E-mail: yasser.abdelaty @ fagr.bu.edu.eg

### INTRODUCTION

Medicinal and aromatic plants are use by 80% of global population for their medicinal therapeutic effects as reported by WHO (2008). Many of these plants synthesize substances that are useful to the maintenance of health in humans and other animals. These include aromatic substances, most of which are phenols or their oxygen-substituted derivatives such as tannins. Others contain alkaloids, glycosides, saponins and many secondary metabolites (Naguib, 2011)

*Echinacea purpurea* (L.) Moench is an herbaceous perennial and a member of the *Asteraceae* family. Commonly called purple coneflower, it has a natural range extending from Michigan, Ohio, Illinois, and Iowa, to southeastern United States and west to Texas. *E. purpurea* grows at a rate of twelve to eighteen inches a year to a mature height of two to four feet. The leaves are ovate to lanceolate and the flowers are cone-shaped disks adorned with deep pink to purple ray flowers. Flowers bloom from June to August (Hendawy, 2000).

*Echinacea* is an herbal medicine, which is used by native Americans for enhancing the human immune system. In Europe and North America, they widely used the *Echinacea purpurea* as the herbal medicine for most of the remedies. Due to the increased market demand, economic value and its potential benefits to human health make increased cultivation of *Echinacea*. Beside, *Echinacea purpurea* is used for both ornamental and phytochemical in Europe, United States and Australia. Three groups of phytochemicals are determined such as caffeic acid derivatives, polysaccharides and lipophilic alkalamides, which are responsible for the genus medicinal properties. Most of Americans and European

have been used medicinal preparation of *Echinacea purpurea* for remedy of many diseases, including colds, toothaches, snake bites, headache and wound infections. It is the most effective antioxidant and it has immunoenhancing effects. *Echinacea* roots are used to treat blood poisoning, snake poisoning, skin disease, syphilis and rabbits. *Echinacea purpurea* herb is also used to treat chronic infections of respiratory tract and lower urinary tract (viral and bacterial origin). The polysaccharide from *Echinacea purpurea* is used to kill bacteria such as staphylococci. Arabinogalactan, a high molecular weight purified polysaccharide from plant cell cultures of *Echinacea purpurea* has potent to activate macrophage cytotoxicity actions against tumor cells and micro organisms. The main caffeic acid derivative (caftaric acid, chlorogenic acid and echinacoside) has been functionally linked to antiinflammatory and wound healing properties of *Echinacea* when applied topically. Caffeic acid derivatives are very effective antioxidants in free radical generation systems. Groups of phenolic compounds and alkalamides, which have demonstrated antiviral and antifungal properties, respectively (Kumar and Ramaiah, 2011).

Recently, unconventional efforts are used to minimize the amounts of chemical fertilizers which applied to medicinal and aromatic plants in order to reduce production cost and environmental pollution without reduction of yield. Therefore, the trend now is using the bio and organic fertilizers. Bio-fertilizers are reasonably safer to the environment than chemical fertilizers and play an important role in decreasing the use of chemical fertilizers. Consequently, it causes a

reduction in environmental pollution. Bio fertilizers are microbial inoculants consisting of living cells of micro-organism like bacteria, algae and fungi alone or in combination which may help in increasing crop productivity. Bio fertilizers can influence plant growth directly through the production of phytohormones such as gibberellins, cytokinins and IAA that act as growth regulators and indirectly through nitrogen fixation and production of bio-control agents against soil-borne phytopathogens and consequently increase formation of metabolites which encourage the plant vegetative growth and enhance the meristematic activity of tissues to produce more growth (Glick, 2003 and Ahmed and Kibret, 2014). The effect of bio-fertilizers on vegetative growth, yield and oil productivity in several studied was revealed by Badran and Safwat (2004) on fennel. Ismail (2007) on dragonhead plants, Amran (2013) on *Pelargonium graveolens* plants and El-Khyat (2013) on *Rosmarinus officinalis*.

Organic fertilizers are obtained from animal sources such as animal manure or plant sources like green manure. Continuous usage of inorganic fertilizer affects soil structure. Hence, organic manures can serve as alternative to mineral fertilizers for improving soil structure (Shahram and Ordoorkhani, 2011) and microbial biomass (Suresh *et al.* 2004). The addition of organic fertilizers to agricultural soils has beneficial effects on crop development and yields by improving soil physical and biological properties (Zheljazkov and Warman, 2004). Organic and bio fertilizers in comparison of the chemical fertilizers have lower nutrient content and are slow release but they are as effective as chemical fertilizers over longer periods of use (Naguib, 2011 and Mohamed *et al.*, 2012).

Micro-nutrients, especially Zn and B act either as metal components of various enzymes or as functional, structural, or regulatory cofactors. Thus, they are associated with saccharide metabolism, photosynthesis and protein synthesis (Marschner, 1997). Many investigators reported the stimulating effect of applied micronutrients as foliar spray on growth and flowering of different medicinal and aromatic plants; i.e. El-Khyat (2013) on *Rosmarinus officinalis* and Amran (2013) on *Pelargonium graveolens*. They indicated that foliar application of Fe, Zn and Mn improved the growth and chemical composition of the plants. Also, Gomaa and Mady (2008) found that boron foliar spray at 75 ppm increased growth parameter and oil productivity of chamomile plants. Therefore the purpose of this study was to evaluate the benefits of supplementing organic and bio fertilizers with chemical fertilizer in presence of Zn and B on growth and chemical composition of *Echinacea purpurea* plants and to minimize consuming of chemical fertilizers.

## MATERIALS AND METHODS

Two field experiments were carried out at Special Farm at El Khatatba region Minofia Governorate, Egypt and in the Laboratory of Horticulture Department, Faculty of Agriculture at Moshtohor, Benha Univ., during 2014 and 2015 seasons to evaluate the effect of some fertilization

treatments and some micro-nutrients i.e. Zn and B each at 100 ppm as well as their combinations on growth and chemical constituents of *Echinacea purpurea* plants.

*Echinacea purpurea* seedlings (7-9 height with 4-5 leaves) were obtained from Floriculture Farm, Horticulture Department, Faculty of Agriculture, Benha Univ. The seedlings were transplanted in sandy soil on mid February of both seasons in beds (1x1 m) containing two rows (50 cm inbetween) and each row contained two hills (50 cm apart). The soil was directly irrigated to provide suitable moisture for growth. All the traditional cultural practices for growing *Echinacea purpurea* plants were followed as recommended in this region.

Physical and chemical analyses of the used soil are presented in Table (a). Mechanical and chemical soil analyses were determined according to Black *et al.*, (1982).

**Table a. Mechanical and chemical analyses of the used soil.**

Physical analysis	Chemical analysis				
	Cations (meq/l)		Anions (meq/l)		
Coarse sand	56.2%	Ca <sup>++</sup>	1.34	CO <sub>3</sub> <sup>--</sup>	Zero
Fine sand	29.8%	Mg <sup>++</sup>	0.81	HCO <sub>3</sub> <sup>-</sup>	2.23
Silt	6.2%	Na <sup>+</sup>	1.94	Cl <sup>-</sup>	1.48
Clay	7.8 %	K <sup>+</sup>	0.12	SO <sub>4</sub> <sup>--</sup>	0.78
Texture class	Sandy	Soil pH		7.98	
		EC		0.43dS m <sup>-1</sup>	
		Organic matter		0.81 g kg <sup>-1</sup>	
		Available N		16.2 mg kg <sup>-1</sup>	
		Available P		6.93 mg kg <sup>-1</sup>	
		Available K		98 mg kg <sup>-1</sup>	

## Bio-fertilizer treatments

*Echinacea purpurea* seedlings were inoculated with a mixture of nitrobein + phosphorein contained efficient strains of nitrogen fixing bacteria namely, *Azotobacter chroococcum* + phosphate dissolving bacteria (*Bacillus megaterium* var phosphaticum) which supplied by the Department of Microbiology, Agric. Res. Center, Giza was used in this study as biological activators. The strains were characterized by a good ability to infect its specific host plant and by its high efficiency in N-fixation and phosphate solubilizing. The roots of *Echinacea purpurea* seedlings were washed with water, thereafter the roots were soaked in cell suspension of the mixture of nitrobein and phosphorein (1ml contains 10<sup>8</sup> viable cell) for 30 min. Gum arabic (16 %) was added as an adhesive agent prior to soaking the roots. The inoculated roots were air dried at room temperature for one hour before planting. Another two applications were applied (1kg/fed) as an aqueous solution, the first one was applied just before irrigation after 60 days from planting date, whereas the second one was done after 90 days from planting date to increase the power ability of bacteria.

## Organic fertilizer treatments

Organic fertilizer i.e. compost containing plant sources and cattle manure at the rate of 10 ton/fed., was thoroughly mixed with the soil before planting, the chemical properties of the tested compost are presented in Table (b).

**Table b. Chemical properties of the used compost:**

Parameters	Ec dS.m <sup>-1</sup> (1:5)	pH (1:5)	Total C %	Total N %	Total P %	Total K %	Total Fe (ppm)	Total Zn (ppm)	C:N ratio
Values	2.81	6.69	24.21	1.34	0.77	1.62	1368	347	18:1

**Chemical fertilizer treatments**

The plants were fertilized with full chemical fertilizer dose as a recommended dose; where ammonium nitrate (33.5% N) was added at the rate of 100Kg N/fed., calcium superphosphate (15.5 % P<sub>2</sub>O<sub>5</sub>) was added at the rate of 150 Kg and potassium sulphate (48.5 % K<sub>2</sub>O) at the rate of 150 Kg /fed. The amount of N and K fertilizers were divided into six equal portions as side dressing at 30, 45, 60, 75, 90 and 110 days after planting date of both seasons. However, the amount of P-fertilizer was added to the soil before seedlings plantins during soil preparation.

**Micro-nutrients treatments**

In the two seasons, the foliage was sprayed four times during the growth period with Zn and B each at concentration of 100 ppm at 60, 90, 120 and 150 days after planting date of both seasons. A surfactant (Tween 20) at a concentration of 0.01% was added to all tested solution treatments including the control.

**Experiment layout**

This experiment was set up in a split plot design with three replicates. The main plot was employed by five fertilization treatments i.e. control (compost at 10 ton/fed,(T1) ), full chemical fertilizer dose ( T2), 75% chemical fertilizer dose + compost at 10 ton/fed. + bio fertilizer (nitroben + phosphorein) (T3), 50% chemical fertilizer dose + organic fertilizer (compost) + bio fertilizer (T4) and 25% chemical fertilizer dose + organic fertilizer + bio fertilizer (T5). Whereas, the sub plot was devoted to three micro-nutrients sprays i.e. control (tap water), Zn at 100 ppm and B at 100 ppm.

**Data recorded**

In both seasons, at harvest time (Mid - September) the following measurements were conducted as follow; plant height (cm.), number of branches/plant, fresh and dry weights of herb (leaves and stems)/plant (g), number of suckers/plant, number of flowering heads/plant, fresh and dry weights of flowering heads/plant (g). In addition, Total nitrogen, phosphorus and potassium were determined in *Echinacea purpurea* leaves at the flowering stage, according to the methods described by Horneck and Miller (1998), Sandell (1950) and Horneck and Hanson (1998), respectively.

Total alkalamides were quantitatively determined in *Echinacea purpurea* aerial parts (flowering heads, leaves and stems) in the second season (2015) using High Performance Liquid Chromato-graphy (HPLC) according to Bauer and Remiger (1989). Total caffeic acid derivatives content in the plant organs (roots, herb and flowering heads) of *Echinacea purpurea* were determined as chicoric acid using spectrophotometer according to A.O.A.C (1980).

**Statistical analysis:**

The obtained data in both seasons of study were subjected to analysis of variance as a factorial

experiment in split plot design. L.S.D. method was used to differentiate between means according to Snedecor and Cochran (1989).

**RESULTS AND DISCUSSION**

**Effect of some fertilization and micro-nutrients treatments as well as their combination on growth and chemical composition of *Echinacea purpurea* plant.**

**Vegetative growth parameters:**

**Plant height (cm)**

Data presented in Table (1) indicated that all tested fertilization treatments significantly increased the plant height of *Echinacea purpurea* as compared with control in both seasons, with superior for T2 treatment (100% chemical fertilizer), followed by T3 treatment (75% chemical fertuilizer+10 ton compost/fed+bio) in the two seasons. Moreover, all tested treatments of micro-nutrients i.e., Zn and B each at 100 ppm succeeded in increasing plant height, especially Zn treatment as compared with control in the two seasons. However, most tested combinations of fertilization and micro-nutrients induced significant increments in plant height in the first and second seasons as compared with control. However, the highest value of plant height (91.3 and 82.4cm) were scored by the combined treatment of T2 and Zn at 100 ppm in the first and second seasons, respectively.

**Number of branches /plant**

Table (1) reveals that number of branches /plant was increased due to fertilization treatments as compared with control in the two seasons. However, the highest number of branches per plant was registered by T3, followed by T2 treatment, with non significant deferences between them in both seasons. Moreover, all tested applications of micro-nutrients showed significant increaments in this regard. Anyway, the highest number of branches/plant was gained by B at 100 ppm in both seasons as compared with control. Regarding the interaction effect between fertilization and micro-nutrients, it was found that all combinations of fertilization and micro-nutrients led to increase the number of branches per plant in both seasons. However, the highest number of branches per plant (9.16 and 8.61) was scored by the interaction between T3 and B at 100 ppm in the first and second seasons, respectively.

**Fresh and dry weights of herb (g) / plant.**

Table (2) shows that fresh and dry weights of herb per plant were significantly increased by all teted fertilizations and micro-nutrients treatments in both seasons. However, T3 treatment succeeded in producing the heaviest fresh and dry weights of herb per plant in both seasons. Moreover, all treatments of micro-nutrients statistically increased the fresh and dry weights of herb per plant, especially the treatment of B at 100 ppm in the two seasons. As for the interaction

effect between fertilization and micro-nutrients, it was observed that all resulted combinations of fertilization and micro-nutrients increased the fresh and dry weights of herb per plant as compared with control in both

seasons. In all, the heaviest fresh and dry weights of herb per plant in both seasons were recorded by the combined treatment between T3 and B at 100 ppm in both seasons.

**Table 1. Effect of some fertilization and micro-nutrients treatments on plant height number of branches /plant of *Echinacea purpurea* plants during 2014 and 2015 seasons.**

Parameters Fertilization	First season (2014)							
	Micro-nutrients Control	Plant height (cm)			Number of branches /plant			
		Zn at 100 ppm	B at 100 ppm	Mean	Control	Zn at 100 ppm	B at 100 ppm	Mean
T1: Control (compost at 10 ton/fed.)	68.3	74.6	71.9	71.6	5.64	6.26	7.39	6.43
T2: (chemical NPK)	86.2	91.3	89.3	88.9	7.84	8.16	8.94	8.31
T3:(75%NPK+ 10 ton compost/fed.)+bio	84.9	89.4	88.1	87.5	7.89	8.24	9.16	8.43
T4:(50%NPK+ 10 ton compost/fed.)+bio	79.8	84.1	81.7	81.9	7.16	8.16	8.43	7.91
T5:(25%NPK+ 10 ton compost/fed.)+bio	74.6	79.5	76.4	76.8	6.88	7.83	7.94	7.55
Mean	78.8	83.8	81.5		7.08	7.73	8.37	
L.S.D at 0.05 for fertilization			3.12				0.32	
Micro-nutrients interaction			2.74				0.29	
			6.61				0.69	
Second season (2015)								
T1: Control (compost at 10 ton/fed.)	65.6	68.9	67.1	67.2	5.14	6.72	6.94	6.26
T2: (chemical NPK)	78.4	82.4	81.0	80.6	8.12	8.24	8.36	8.24
T3:(75%NPK+ 10 ton compost/fed.)+bio	76.7	81.2	78.3	78.7	8.26	8.41	8.61	8.42
T4:(50%NPK+ 10 ton compost/fed.)+bio	73.1	76.3	74.9	74.8	7.64	7.91	8.21	7.92
T5:(25%NPK+ 10 ton compost/fed.)+bio	69.8	72.4	71.3	71.2	7.20	7.31	7.61	7.37
Mean	72.7	76.2	74.5		7.27	7.71	7.94	
L.S.D at 0.05 for fertilization			2.94				0.31	
Micro-nutrients interaction			2.63				0.27	
			6.32				0.72	

**Table 2: Effect of some fertilization and micro-nutrients treatments on fresh weight of herb/plant (g) and dry weight of herb/plant (g) of *Echinacea purpurea* plants during 2014 and 2015 seasons.**

Parameters Fertilization	First season (2014)							
	Micro-nutrients Control	Fresh weight of herb/plant (g)			Dry weight of herb/plant (g)			
		Zn at 100 ppm	B at 100 ppm	Mean	Control	Zn at 100 ppm	B at 100 ppm	Mean
T1: Control (compost at 10 ton/fed.)	841	992	1184	1006	92.5	109.0	131.0	110.8
T2: (chemical NPK)	1248	1296	1468	1337	149.6	155.5	176.2	160.4
T3:(75%NPK+ 10 ton compost/fed.)+bio	1262	1312	1483	1352	151.4	157.4	177.9	162.2
T4:(50%NPK+ 10 ton compost/fed.)+bio	1152	1296	1365	1271	132.4	149.0	156.9	146.1
T5:(25%NPK+ 10 ton compost/fed.)+bio	1088	1251	1286	1208	121.8	140.1	144.0	135.3
Mean	1118	1230	1357		129.5	142.2	157.2	
L.S.D at 0.05 for fertilization			84.6				24.6	
Micro-nutrients interaction			74.7				21.6	
			178.5				51.9	
Second season (2015)								
T1: Control (compost at 10 ton/fed.)	719	1072	1124	972	79.1	117.9	123.6	106.9
T2: (chemical NPK)	1218	1312	1354	1295	146.1	157.4	162.4	155.3
T3:(75%NPK+ 10 ton compost/fed.)+bio	1229	1344	1361	1311	147.4	161.2	163.2	157.3
T4:(50%NPK+ 10 ton compost/fed.)+bio	1130	1264	1328	1241	129.9	145.3	152.7	142.6
T5:(25%NPK+ 10 ton compost/fed.)+bio	1071	1168	1231	1157	119.9	130.8	137.8	129.5
Mean	1073	1232	1280		124.5	142.5	147.9	
L.S.D at 0.05 for fertilization			73.1				18.9	
Micro-nutrients interaction			64.3				16.6	
			154.3				39.8	

**Number of suckers/plant**

Table (3) declares that all tested fertilization treatments succeeded in increasing number of suckers/plant as compared with control in both seasons.

Moreover, all micro-nutrients treatments significantly increased number of suckers per plant, particularly B treatment in the two seasons. As for the interaction effect between fertilization and micro-nutrients

treatments, it was observed that all interactions of fertilization and micro-nutrients treatments increased the number of suckers per plant as compared with control in both seasons. In general, the highest number of suckers/plant (8.43 and 8.62) were recorded by the combined treatment between T3 and B at 100 ppm in the first and second seasons, respectively.

The aforementioned results of tested fertilization treatment are in agreement with those obtained by Kandeel (2004) on *Ocimum basilicum*, Niakan *et al.* (2004) on *Mentha piperita*, El-Maadawy (2007) on *Amaranthus tricolor*, El-Maadawy and Moursy (2007) on jojoba, Gomaa and Youssef (2007a) on fennel, Badran *et al.* (2007) on cumin, Gomaa and Youssef (2007b) on lovage, El-Shora (2009) on *Mentha piperita*, Abou El-Ghait *et al.* (2012) on indian fennel, Gendy *et al.* (2012) on roselle plants, Gendy *et al.* (2013) on guar plants, Mohamed *et al.* (2012) on *Stevia rebaudiana*, Amran (2013) on *Pelargonium graveolens* and El-Khyat (2013) on *Rosmarinus officinalis*. Whereas, the abovementioned results of micro-nutrients are nearly similar to those obtained by Hendawy (2000) on *Echinacea purpurea*, Kuntal *et al.*, (2005) on *Stevia rebaudiana*, Gomaa (2008) on *Hibiscus sabdariffa*, Youssef (2009) on rosemary plant, Nasiri *et al.*, (2010) on chamomile plant, Ajay *et al.*, (2010) on *Mentha arvensis* L., Said-Al Ahl and Mahmoud (2010) on sweet basil, Amuamuha *et al.*, (2012) on marigold plant., Khalid (2012) on anise plant, Shilpa and Dhumal (2012) on *Cassia angustifolia*, Saeid Zehtab *et al.*, (2012) on Psyllium plant, Amran (2013) on *Pelargonium graveolens* and El-Khyat (2013) on *Rosmarinus officinalis*.

**Flowering growth parameters:**

Tables (3) and (4) illustrate that all tested fertilization treatments resulted in significant

increments in number, fresh and dry weights of flowering heads/plant, with superiority for T3 treatment as compared with control in both seasons. Also, all tested micro-nutrients statistically increased the number, fresh and dry weights of flowering heads/plant, especially those received B treatment in both seasons.

As for the interaction effect between fertilization and micro-nutrients, it was found that all resulted interactions of fertilization and micro-nutrients increased the values of the number, fresh and dry weights of flowering heads/plant. However, the highest number of flowering heads/plant (23.60 and 25.30), the heaviest fresh weight of flowering heads/plant (542.8 and 581.7 g) and the heaviest dry weight of flowering heads/plant (74.2 and 79.1 g) were recorded by T3 treatment combined with B at 100 ppm, in the first and second seasons, respectively.

The aforementioned results of fertilization treatment are in parallel with those obtained by El-Maadawy (2007) on *Amaranthus tricolor*, El-Maadawy and Moursy (2007) on jojoba, Gomaa and Youssef (2007a) on fennel, Badran *et al.* (2007) on cumin, Gomaa and Youssef (2007b) on lovage, Abou El-Ghait *et al.* (2012) on indian fennel, Gendy *et al.* (2012) on roselle plants, Gendy *et al.* (2013) on guar plants, Mohamed *et al.* (2012) on *Stevia rebaudiana* and Amran (2013) on *Pelargonium graveolens*. While, the aforementioned results of micro-nutrients are in parallel with those obtained by Hendawy (2000) on *Echinacea purpurea*, Gomaa (2008) on *Hibiscus sabdariffa*, Nasiri *et al.*, (2010) on chamomile plant, Amuamuha *et al.*, (2012) on marigold plant., Khalid (2012) on anise plant, Shilpa and Dhumal (2012) on *Cassia angustifolia*, and Saeid Zehtab *et al.*, (2012) on Psyllium plant.

**Table 3: Effect of some fertilization and micro-nutrients treatments on number of suckers and flowering heads / plant of *Echinacea purpurea* plants during 2014 and 2015 seasons.**

Parameters	First season (2014)								
	Number of suckers/plant				Number of flowering heads/plant				
	Micro-nutrients	Cont rol	Zn at 100 ppm	B at 100 ppm	Mean	Control	Zn at 100 ppm	B at 100 ppm	Mean
<b>Fertilization</b>									
T1: Control (compost at 10 ton/fed.)		4.21	4.62	4.92	4.58	11.24	12.31	15.17	12.90
T2: (chemical NPK)		7.26	7.84	8.39	7.83	18.17	19.26	22.92	20.13
T3:(75%NPK+10ton compost/fed.)+bio		7.39	7.89	8.43	7.90	18.29	19.21	23.60	20.36
T4:(50%NPK+10 ton compost/fed.)+bio		5.36	6.14	6.81	6.10	15.36	17.31	20.11	17.59
T5:(25%NPK+10 ton compost/fed.)+bio		4.94	5.26	5.84	5.34	13.98	15.62	18.21	15.93
Mean		5.83	6.35	6.87		15.40	16.74	20.00	
L.S.D at 0.05 for	fertilization			0.45				1.23	
	Micro-nutrients interaction			0.39				1.08	
				0.94				2.59	
		Second season (2015)							
T1: Control (compost at 10 ton/fed.)		4.82	5.04	5.86	5.24	13.46	14.50	16.94	14.97
T2: (chemical NPK)		7.84	8.19	8.57	8.20	19.46	21.27	24.91	21.88
T3:(75%NPK+10 ton compost/fed.)+bio		7.94	8.23	8.62	8.26	19.36	21.36	25.30	22.00
T4:(50%NPK+10 ton compost/fed.)+bio		6.92	6.98	7.30	7.06	17.21	19.24	21.46	19.30
T5:(25%NPK+10 ton compost/fed.)+bio		5.64	5.93	6.74	6.10	15.92	17.21	19.20	17.44
Mean		6.63	6.87	7.41		17.08	18.72	21.56	
L.S.D at 0.05 for	fertilization			0.34				1.44	
	Micro-nutrients interaction			0.29				1.26	
				0.71				3.00	

**Table 4: Effect of some fertilization and micro-nutrients treatments on fresh and dry weights of flowering heads of *Echinacea purpurea* plants during 2014 and 2015 seasons.**

Parameters	First season (2014)							
	Fresh weight of flowering heads/plant (g)				Dry weight of flowering heads/plant (g)			
	Control	Zn at 100 ppm	B at 100 ppm	Mean	Control	Zn at 100 ppm	B at 100 ppm	Mean
T1: Control (compost at 10 ton/fed.)	224.2	258.3	319.3	267.3	29.1	34.1	43.1	35.4
T2: (chemical NPK)	380.6	422.1	503.6	435.4	51.3	56.9	67.9	58.7
T3:(75%NPK+ 10 ton compost/fed.)+bio	399.7	416	542.8	452.8	54.6	56.9	74.2	61.9
T4:(50%NPK+ 10 ton compost/fed.)+bio	321.4	363.9	442.3	375.9	43.1	48.6	59.2	50.3
T5:(25%NPK+ 10 ton compost/fed.)+bio	280.7	312.8	401.7	331.7	37.2	41.5	53.3	44.0
Mean	321.3	354.6	441.9		43.1	47.6	59.5	
L.S.D at 0.05 for fertilization			25.3			8.6		
Micro-nutrients interaction			22.2			7.5		
			53.4			18.1		
Second season (2015)								
T1: Control (compost at 10 ton/fed.)	281.6	319.4	372.7	324.6	36.8	41.7	48.7	42.4
T2: (chemical NPK)	426.3	466.8	572.6	488.6	57.5	62.9	77.2	65.9
T3:(75%NPK+ 10 ton compost/fed.)+bio	443.4	489.4	581.7	504.8	60.2	66.5	79.1	68.6
T4:(50%NPK+ 10 ton compost/fed.)+bio	378.0	422.3	471.5	423.9	50.6	56.4	63.1	56.7
T5:(25%NPK+ 10 ton compost/fed.)+bio	349.7	378.5	422.6	383.6	46.1	49.8	55.7	50.5
Mean	375.8	415.3	484.2		50.2	55.5	64.8	
L.S.D at 0.05 for fertilization			38.4			6.2		
Micro-nutrients interaction			33.8			5.4		
			81.1			13.1		

**Leaf chemical composition:**

Table (5) declares that all tested treatments fertilization and micro-nutrients increased leaf N, P and K content of *Echinacea purpurea* leaves in both seasons. In this concern, the richest leaf N, P and K content was recorded by T2 and B at 100 ppm treatments as well as their combinations in both seasons as compared with control and the rest treatments. This may be due to the combined effects of both fertilization treatments which induces cell division and enlargement, in addition to the effects of the tested micro-nutrients which supply the plant with the required nutrients necessary for healthy growth.

**Total alkaloids (%)**

Table (6) demonstrates that all tested fertilization and micro-nutrients treatments as well as their interactions succeeded in increasing total alkaloids (%) in the above ground organs (leaves, stems and flowering heads) of *Echinacea purpurea* plant. However, the highest value of total alkaloids (0.082%) was recorded by T3 treatment combined with Zn at 100 ppm, followed by the combined treatment between T3 and B at 100 ppm treatments as it gave 0.079 against to 0.034% for control.

**Total caffeic acid derivatives (%)**

It is clear from Table (7) that the flowering heads of *Echinacea purpurea* were the richest plant part with total caffeic acid derivatives as they contained 0.75 and 0.83%, followed by the herb (0.43 and 0.50%), then the roots (0.29 and 0.31%) in the first and second seasons, respectively. However, T3 treatment showed to be the most effective one in inducing the highest values of total caffeic acid derivatives of *Echinacea purpurea* plants (flowering heads, herb and roots), in both

seasons. Also, all tested micro-nutrients sprays increased total caffeic acid derivatives, especially Zn at 100 ppm in case of flowering heads, herb parts and root parts in both seasons. In general the highest values of total caffeic acids derivatives in the flowering heads (0.92 and 0.94%), in the herb (0.59 and 0.64%) and in the roots (0.43 and 0.41%) were recorded by the combined treatment between T3 treatment and Zn at 100 ppm, in the first and second seasons, respectively.

The aforementioned results of fertilization concerning chemical constituents are in parallel with those obtained by El-khayat (2001) on roselle plants, Kandeel (2004) on *Ocimum basilicum*, Niakan *et al.* (2004) on *Mentha piperita*, El-Maadawy (2007) on *Amaranthus tricolor*, El-Maadawy and Moursy (2007) on jobba, Gomaa and Youssef (2007a) on fennel plant, Badran *et al.*, (2007) on cumin plant, Gomaa and Youssef (2007b) on lovage plants, El-Shora (2009) on *Mentha piperita*, Abou El-Ghait *et al.* (2012) on indian fennel plant, Gendy *et al.* (2012) on roselle plants, Mohamed *et al.* (2012) on *Stevia rebaudiana*, Amran (2013) on *Pelargonium graveolens* and El-Khyat (2013) on *Rosmarinus officinalis*.

The aforementioned results of micro-nutrients are in conformity with those obtained by Hendawy (2000) on *Echinacea purpurea*, Gomaa (2008) on *Hibiscus sabdariffa*, Youssef (2009) on rosemary plant, Nasiri *et al.*, (2010) on chamomile plant, Ajay *et al.*, (2010) on *Mentha arvensis* L., Amuamuha *et al.* (2012) on marigold plant., Shilpa and Dhupal (2012) on *Cassia angustifolia*, Saeid Zehtab *et al.*, (2012) on *Psyllium* plant, Amran (2013) on *Pelargonium graveolens* and El-Khyat (2013) on *Rosmarinus officinalis*.

**Table 5: Effect of some fertilization and micro-nutrients treatments on leaf nitrogen , phosphorus and potassium contents of *Echinacea purpurea* plants during 2014 and 2015 seasons.**

Parameters	First season (2014)											
	Con.	N% Zn at 100 ppm	B at 100 ppm	Mean	Con.	P% Zn at 100 ppm	B at 100 ppm	Mean	Con.	K% Zn at 100 ppm	B at 100 ppm	Mean
T1:Control (compost at 10 ton/fed.)	1.86	1.92	1.98	1.92	0.216	0.221	0.219	0.219	1.26	1.29	1.34	1.30
T2: (chemical NPK)	2.42	2.49	2.51	2.47	0.237	0.252	0.250	0.246	1.48	1.57	1.68	1.58
T3:(75%NPK+10 ton compost/fed.)+bio	2.46	2.41	2.48	2.45	0.234	0.249	0.248	0.244	1.46	1.54	1.64	1.55
T4:(50%NPK+10 ton compost/fed.)+bio	2.19	2.34	2.41	2.31	0.229	0.236	0.234	0.233	1.39	1.42	1.53	1.45
T5:(25%NPK+10 ton compost/fed.)+bio	1.94	1.99	2.17	2.03	0.221	0.228	0.226	0.225	1.32	1.36	1.42	1.37
Mean	2.17	2.23	2.31		0.227	0.237	0.235		1.38	1.44	1.52	
L.S.D fertilization at 0.05 Micro-nutrients for interaction		0.14 0.12 0.29				0.011 0.009 0.023				0.11 0.09 0.23		
	Second season (2015)											
T1:Control (compost at 10 ton/fed.)	1.79	1.84	1.94	1.86	0.221	0.232	0.229	0.227	1.31	1.33	1.39	1.34
T2: (chemical NPK)	2.30	2.39	2.49	2.39	0.251	0.281	0.273	0.268	1.61	1.64	1.73	1.66
T3:(75%NPK +10 ton compost/fed.) + bio	2.31	2.36	2.46	2.38	0.253	0.283	0.270	0.269	1.58	1.61	1.71	1.63
T4:(50%NPK +10 ton compost/fed.)+bio	2.13	2.18	2.21	2.17	0.247	0.261	0.258	0.255	1.47	1.56	1.62	1.55
T5:(25%NPK +10 ton compost/fed.) + bio	1.91	1.92	2.08	1.97	0.236	0.254	0.251	0.247	1.39	1.42	1.49	1.43
Mean	2.09	2.14	2.24		0.241	0.262	0.256		1.47	1.51	1.59	
L.S.D fertilization at 0.05 micro-nutrients for interaction		0.16 0.14 0.33				0.012 0.010 0.025				0.09 0.07 0.18		

**Table 6: Effect of some fertilization and micro-nutrients treatments on total alkamides (%) of *Echinacea purpurea* during 2015 season.**

Micro-nutrients Fertilization	Control (0.0 ppm)	Zn at 100 ppm	B at 100 ppm
T1: Control (compost at 10 ton/fed.)	0.034	0.046	0.041
T2: (chemical NPK)	0.051	0.071	0.067
T3:(75%NPK+10 ton compost/fed.)+bio	0.058	0.082	0.079
T4:(50%NPK+10 ton compost/fed.)+bio	0.049	0.064	0.062
T5:(25%NPK+10 ton compost/fed.)+bio	0.043	0.053	0.051

**Table 7: Effect of some fertilization and micro-nutrients treatments on total caffeic acid derivatives (g/100g dry weight) of *Echinacea purpurea* plants during 2014 and 2015 seasons.**

Season Organ	First season (2014)											
	Flowering heads				Herb				Roots			
Micronutrients Fertilization	Con	Zn at 100 ppm	B at 100 ppm	Mean	Con.	Zn at 100 ppm	B at 100 ppm	Mean	Con.	Zn at 100 ppm	B at 100 ppm	Mean
T1:Control (compost at 10 ton/fed.)	0.62	0.68	0.65	0.63	0.31	0.39	0.36	0.35	0.19	0.25	0.21	0.21
T2: (chemical NPK)	0.71	0.80	0.79	0.76	0.40	0.48	0.44	0.44	0.28	0.32	0.31	0.30
T3:(75%NPK+10ton compost/fed.)+bio	0.84	0.92	0.88	0.88	0.49	0.59	0.56	0.54	0.34	0.43	0.39	0.38
T4:(50%NPK+10ton compost/fed.)+bio	0.73	0.81	0.80	0.78	0.41	0.49	0.46	0.45	0.29	0.34	0.32	0.31
T5:(25%NPK+10ton compost/fed.)+bio	0.69	0.74	0.71	0.71	0.37	0.42	0.40	0.39	0.24	0.29	0.26	0.26
Mean	0.71	0.78	0.76	0.75	0.39	0.47	0.44	0.43	0.26	0.32	0.29	0.29
L.S.D fertilization at 0.05 micro-nutrients for interaction		0.022 0.019 0.045				0.018 0.015 0.041				0.014 0.012 0.031		
	Second season (2015)											
T1:Control (compost at 10 ton/fed.)	0.69	0.78	0.73	0.73	0.36	0.43	0.41	0.40	0.21	0.26	0.23	0.23
T2: (chemical NPK)	0.81	0.89	0.88	0.86	0.48	0.57	0.52	0.52	0.28	0.37	0.33	0.32
T3:(75%NPK+10ton compost/fed.)+bio	0.84	0.94	0.91	0.89	0.51	0.64	0.61	0.58	0.31	0.41	0.39	0.37
T4:(50%NPK+10ton compost/fed.)+bio	0.79	0.91	0.88	0.86	0.49	0.58	0.53	0.53	0.29	0.38	0.34	0.33
T5:(25%NPK+10ton compost/fed.)+bio	0.76	0.86	0.81	0.81	0.42	0.51	0.49	0.47	0.26	0.31	0.28	0.28
Mean	0.77	0.87	0.84	0.83	0.45	0.54	0.51	0.50	0.27	0.34	0.31	0.31
L.S.D fertilization at 0.05 micro-nutrients for interaction		0.032 0.028 0.064				0.021 0.018 0.045				0.011 0.009 0.025		

The obtained results of this study may be due to the role of fertilization and micro-nutrients in growth and development of the plants; where the use of N-fixing bacteria (nitrobein) as a bio-fertilizer product containing nitrogen fixing bacteria, e.g. Azotobacter and Azospirillum was found to have not only the ability to fix nitrogen but also to release certain phytohormones of cytokinins, gibberellins and auxins which could enhance plant growth through absorption of nutrients and so on enhancing photosynthesis process (Hegde *et al.*, 1999). Microorganisms used as bio-fertilizers may affect the integrity of growing plants by one mechanism or more such as nitrogen fixation production of growth promoting substances or organic acids, enhancing nutrients uptake or protection against plant pathogens (Hawaka, 2000) Also, N-fixers synthesize stimulatory compounds such as, gibberellins, cytokinins and IAA. They act as growth regulators, which increased the surface area per unit of root length and were responsible for root hair branching with an eventual increase in the uptake of nutrients from the soil (Sperenat, 1990 and Dadarwal *et al.*, 1997). Besides, the use of Phosphate dissolving bacteria (phosphorein) as a bio-fertilizer product containing very active phosphatase dissolving bacteria has proved its efficiency in enhancing different aspects of growth and development of many plant species including medicinal and aromatic ones. Establishment of a strong root system is related to the level of available phosphate in the soil. Phosphate dissolvers or vesicular arbuscular mycorrhizae and silica bacteria are capable of converting tricalcium phosphate to monocalcium phosphate ready for plant nutrition. Phosphate also increased mineral uptake and water use efficiency (Hawaka, 2000). Moreover, when organic manures (compost) added as fertilizer, it led to decrease soil pH which in turn increasing solubility of nutrients for plant uptake, in some cases organic materials may act as low release fertilizer. Recently, on the way of sustainable agriculture with minimum effects, the use of organic manures (compost or chicken manure, ..etc) as natural soil amendments is recommended to replace the soluble chemical fertilizers. They improve the structure of weak-structured sandy soils and increase their water holding capacity. Also, they improve soil fertility, and stimulate root development, induce active biological conditions and enhancing activities of micro-organisms especially those involved in mineralization (Suresh *et al.*, 2004). Furthermore, to interpret and evaluate the effect of chemical fertilization concerned in this study, on augmenting the different tested vegetative growth parameters, yield component parameters and chemical constituents of *Echinacea purpurea* plants. It is important to refer to the physiological roles of nitrogen, phosphorus and potassium in plant growth and development. Such three macronutrient elements are the common elements usually included in fertilizers. Plant supplement with these macronutrients in form of fertilizers is necessary because the soil is usually in deficient of them due to plant removal leaching or they are not readily available for plants. Therefore, such addition of well balanced NPK fertilization quantities

insured production of high productivity and chemical constituents of *Echinacea purpurea* plants.

The role of NPK fertilization in promoting vegetative growth characters, enhancing yield component parameters and increasing growth, as well as stimulating the chemical constituents content of *Echinacea purpurea* plants could be explained by recognizing their fundamental involvement in the very large number of enzymatic reaction that depend on NPK fertilization. NPK reflected directly on increasing the content of total carbohydrates, total sugars and total free amino acids as well as NPK % in the leaves were indirectly the cause for enhancing the augmenting of all other vegetative growth traits and chemical constituents of *Echinacea purpurea* plants (Cooke, 1982).

For adequate plant growth and production, micronutrients are needed in small quantities in balance of macronutrients;. However, their deficiencies cause a great disturbance in the physiological and metabolic processes in the plant. Plants normally take up nutrients from soils through their roots although nutrients can be supplied to plants as fertilizers by foliar sprays (Baloch *et al.*, 2008). Zn and B acts either as metal components of various enzymes or as functional, structural, or regulatory cofactors. Thus, it is associated with saccharide metabolism, photosynthesis, and protein synthesis (Marschner, 1997). The positive effects of the tested micro-nutrients may be due to the role of Zinc which is consider one of the essential microelements for growth and flowering of plants (Chandler, 1982). Zinc is an important micronutrient that is closely involved in the metabolism of RNA and ribosomal content in plant cells, leading to stimulation of carbohydrates, proteins and the DNA formation. It is also, required for the synthesis of tryptophan, a precursor of IAA which acts as a growth promoting substance. Zinc has three functions: catalytic, cocatalytic (coactive) and structural (Amberger, 1974). Moreover, boron (B) is one of the important micronutrients, which has basic role in stabilizing certain constituents of cell walls structure and function and activity of plasma membrane, enhancement of cell division, tissue differentiation. Thus, boron could be directly associated with cell growth (Goldbach *et al.*, 1990). Also, boron has been involved in metabolism of nucleic acid, carbohydrate, protein, auxin and phenol. Moreover, boron has been role in sugar translocation, nucleic acids synthesis and pollen tube growth. Also, boron plays a key role in higher plants by facilitating the short- and long- distance transport of sugar via the formation of borate- sugar complexes. However, such a proposal is unacceptable because, the prevalent sugar transport in the phloem forms only weak complexes with boron, and in the mechanisms of phloem loading of sucrose boron is not involved (Marschner, 1997, Goldbach and Wimmer, 2007 and Ganie *et al.*, 2013). Besides, more than 90% of the boron is found in cell walls. Its functions are also related to cell wall synthesis, lignifications and maintenance of cell wall structure (Hansch and Mendel, 2009). In addition, Ganie *et al.* (2013) reported that application of boron increase net photosynthetic rate which may be



attributed to the increase in chlorophylls content of leaves. Furthermore, application of boron increased the activity of catalase and glutathione reductase, which act as antioxidants thus saving the electron transport mechanism of plant from getting oxidized by free radicals like superoxide radicals, singlet oxygen radicals (Wojcik *et al.*, 2008).

Therefore, sufficient amount of these nutrients in the plant is necessary for normal growth, in order to obtain satisfactory yield (Yassen *et al.*, 2010). So, micronutrients such as zinc and boron have important roles in growth and chemical composition of *Echinacea purpurea* plant.

Conclusively, from the obtained results, it is preferable to fertilize *Echinacea purpurea* plants with 75% of chemical fertilizer (NPK) + 10 ton compost/fed. + bio fertilizer (nitrobein + phosphorein) in combination with some micro-nutrients (Zn or B each at 100 ppm) as foliar spray to enhance plant growth and chemical composition which led finally to a safe product of high quality suitable for exportation and safe on human health.

## REFERENCES

- A.O.A.C.(1980): Official Methods of Analysis,12Th Ed. Association of Official Analysis Chemists :Washington , D.C., U.S.A.
- About El-Ghait, E.M.; A.O. Goma; A.S.M. Youssef; E.M. Atia and W.H. Abd-Allah (2012). Effect of sowing dates, bio, organic and chemical fertilization treatments on growth and production of Indian fennel under North Sinai conditions. Bull. Fac., Cairo Univ., 63: 52-68.
- Ahmed, M. and M. Kibret ( 2014): Mechanisms and applications of plant growth promoting rhizobacteria: Current perspective. J. King saud Univ. Sci., 26 : 1-20.
- Ajay, K.; H.K. Patro and K.Walanand (2010). Effect of zinc and sulphur on herb, oil yield and quality of Menthol mint (*Mentha arvensis* L.) var. Kosi. J. Chem. Pharm. Res., 2(4):642-648.
- Amberger, A. (1974): Micro-nutrients, dynamics in the soil and function in plant metabolism. Proc. Egypt, Bot. Soc. Workshop, 1, Cairo .
- Amran, K.A.A. (2013): Physiological studies on *Pelargonium graveolens* L plant. Ph.D. Thesis, Fac. of Agric., Moshtohor, Benha. Univ.
- Amuamuha, L.; A. Pirzad and H. Hadi (2012). Effect of varying concentrations and time of Nanoiron foliar application on the yield and essential oil of pot marigold. Intl. Res. J. Appl. Basic. Sci. Vol., 3 (10), 2085-2090.
- Badran, F.S. and M.S. Safwat (2004): Response of fennel plants to organic and biofertilizer in replacement of chemical fertilization. Proc. 2<sup>nd</sup> International Conf. of organic Agric., Cairo, Egypt, March (2004).
- Badran, F.S.; M.K. Aly; E.A. Hassan and S.G. Shalatek (2007). Effect of organic and bio-fertilization treatments on cumin plants. The third conference of sustainable Agricultural Development, 12-14 November 2007.
- Baloch, O.B.; O.I. Chacar and M.N.Tareen (2008). Effect of foliar application of macro and micronutrients on production of green chilies (*Capicum annum* L.). J Agric Tech 4(2): 177- 184.
- Bauer, R. and P. Remiger (1989). TLC and HPLC analysis of alkamides of Echinacea drug planta Medica,55:367-371.
- Black, C.A.; D.O. Evans; LE. Ensminger; J.L. White; F.E. Clark and R.C. Dinauer (1982). Methods of soil analysis. part 2. Chemical and microbiological properties. 2nd Ed. Soil Sci.,Soc. of Am. Inc. Publ., Madison, Wisconsin, U. S.A.
- Chandler, H. (1982): Zinc As Nutrient for Plants. Bot. Hag. 98, 625-646.
- Cooke, G.W. (1982): Fertilizing for Maximum Yield. Third Edition Granada Publishing limited.
- Dadarwal, L.R.; L.S. Yadav and S.S. Sindhu (1997). Bio-fertilizer Production Technology: Prospects In. Biotechnological approaches: In. Soil Microorganisms for sustainable crop production. Pp: 323- 337. Scientific publisher, Jodhpur, India (C.F. Proceeding of training course on Bio-organic farming systems for sustainable Agriculture. July, 1997, Cairo, Egypt).
- El-Khayat, A.S.M. (2001). Physiological effects of tryptophan, thiamin and ascorbic acid on *Hibiscus sabdariffa*, L. plant. The Fifth Arabian Horticulture Conference, Ismailia, Egypt, March 24-28, Vol.(11) 251-264.
- El-Khyat, L.A.S. (2013). Effect of chemical and bio fertilizer on growth and chemical composition of rosemary plants. M.Sc. Thesis Fac. Agric. Moshtohor, Benha Univ.
- El-Maadawy, E.I. (2007). Response of summer annual flowering plants to chemical, organic and bio-fertilization treatments 1- Joseph's coat (*Amaranthus tricolor*, L.) plants. J. Product. & Dev.,12 (1): 133-152.
- El-Maadawy, E.I. and Kh.S. Moursy (2007). Bio-fertilizers as a partial alternative to chemical NPKL fertilization of jojoba (*Simmondsia chinensis*, L.) plants grown in different soil types. J. Product. & Dev.,12(1): 211-236.
- El-Shora, S.E.E. (2009): Physiological studies on *Mentha Spp.* (fertilization – post harvest treatments). M.Sc. Thesis, Fac. Of Agric. Benha Univ.
- Ganie, M.A.; M.A. Farida, Akhter; M.A. Bhat; A.R. Malik and T.A. Bhat (2013): Boron- a critical nutrient element for plant growth and productivity with reference to temperate fruits. Curr. Sci.,104: 76- 85.
- Gendy, A.S.H.; H.A.H. Said-Al Ahl and Abeer A. Mahmoud (2012): Growth, productivity and chemical constituents of roselle (*Hibiscus sabdariffa* L.) plants as influenced by cattle manure and biofertilizers treatments. Australian J. of Basic and Applied Science, 6(5): 1-12.
- Gendy, A.S.H., H.A.H. Said-Al Ahl; Abeer A. Mahmoud and Hanaa F.Y. Mohamed (2013). Effect of nitrogen sources, bio-fertilizers and their interaction on the growth, seed yield and chemical composition of guar plants. Life Science Journal, 10(3): 389-402.
- Glick, B.R. (2003). Plant growth promoting bacteria. In: Glick, B.R. and J.J. Pasternak (eds.), Molecular Biology-Principles and Applications of Recombinant DNA, pp: 436–54. ASM Press, Washington DC, USA.
- Goldbach, H.E.; D. Hartmann and T. Rotzer (1990). Boron is required for the stimulation of the ferricyanide-induced proton release by auxins in suspension-cultured cells of *Daucus carota* and *Lycopersicon esculentum*. Physiol. Plant., 80: 114-118.
- Goldbach, H.E. and M.A. Wimmer (2007). Boron in plants and animals: Is there a role beyond cell-wall structure? J. Plant Nutr. Soil Sci., 170: 39-48.

- Gomaa, A.O. (2008): Effect of tryptophan, Fe, Zn, and Mn foliar application on growth, productivity and chemical composition of roselle plants. *J. Biol. Chem. Environ. Sci.*, Vol. 3(1): 771-790.
- Gomaa, A.O. and A.S.M. Youssef (2007a). Bio-fertilizers as a partial alternative to chemical N.P.K. fertilization and its influence on the productivity of fennel plants (*Foeniculum vulgare*, Miller). The Third Conf. of Sustain. Agric. and Develop. Fac. of Agric., Fayoum Univ., 12-14 Nov.
- Gomaa, A.O. and A.S.M. Youssef (2007b). Influence of chemical, organic and bio-fertilizer application on growth and productivity of lovage plant (*Levisticum officinale*, Koch). *Egypt J. of Appl. Sci.*, 22 (IIB): 492-520
- Gomaa, A.O. and M.A. Mady (2008). Response of chamomile plants to foliar spray with boron and some antioxidants. The 4<sup>th</sup> Scientific Conference of the Agricultural & Biological Research Division.
- Hansch, R. and R.R. Mendel (2009). Physiological functions of mineral micronutrients (Cu, Zn, Mn, Fe, Ni, Mo, B, Cl). *Curr. Opin in Plant Biol.*, 12: 259-266.
- Hawaka, F.I.A. (2000). Effect of using single and composite inoculation with *Azospirillum brasilense*, *Bacillus megatherium* var. *Phosphaticum* and *Glomus macrocarpus* for improving growth of *Zea mays*. *J. Agric. Sci. Mansoura, Egypt*, 32(12): 239-252.
- Hegde, D.M.; B.S. Dwivedi and S.S. Sudhakar Babu (1999). Biofertilizers for cereal production in India. A review. *Ind. J. Agric. Res.*, 69(2): 73-83.
- Hendawy, S.F. (2000). Physiological and chemical studies on *Echinacea purpurea* L., plant. Ph.D. Theses, Fac., Agric., Moshtohor, Zaga. Univ. Egypt.
- Horneck, D.A. and D. Hanson (1998). Determination of potassium and sodium by flame Emission spectrophotometry. In hand book of reference methods for plant analysis, e.d. Kolra, Y.P. (e.d.) 153-155.
- Horneck, D.A. and R.O. Miller (1998). Determination of total nitrogen in plant tissue. In hand book of reference methods for plant analysis, (e.d) Kolra, Y.P. 73.
- Ismail, S.E. (2007). Comparison study between bio and mineral fertilization of dragonhead plant on yield and volatile oil. Ph. D. Thesis, Fac. Agric. Moshtohor Benha Univ. Egypt.
- Kandeel, Y.M.R. (2004). Effect of bio, organic and chemical fertilization on growth, essential oil productivity and chemical composition of *Ocimum basilicum*, L. plant. *Annals of Agric. Sci., Moshtohor*, Vol. 42 (3): 1253-1270.
- Khalid, A. K. (2012). Effect of NP and foliar spray on growth and chemical compositions of some medicinal Apiaceae plants grow in arid regions in Egypt. *Journal of Soil Science and Plant Nutrition*, 12 (3) 617-632.
- Kumar, K.M. and S. Ramaiah (2011). Pharmacological importance of *Echinacea purpurea*. *International Journal of Pharma and Bio Sciences*. Vol. 2(4): 304-314.
- Kuntal, D.R.; T.N. Shivananda and P. Sur (2005). Interaction between phosphorus and zinc on the biomass yield and yield attributes of the medicinal plant *Stevia rebaudiana*. *The Scientific World Journal*, 5, 390-395.
- Marschner, H. (1997). *Mineral Nutrition of Higher Plants*. 2<sup>nd</sup> ed. San Diego: Academic Press, 379-396.
- Mohamed, S.M.; E. M. Abou El-Ghait; A.S.M. Youssef; A.M.M. Khalil and K. E. Attia (2012). Effect of irrigation rate and some fertilization treatments on vegetative growth and chemical composition of *Stevia rebaudiana*. *Annals of Agric. Sci., Moshtohor*, 50(4): 435-446.
- Naguib, N.Y.M. (2011): Organic vs chemical fertilization of medicinal plants: a concise review of researches. *Adv. Environ. Biol.*, 5(2): 394-400.
- Nasiri Y.; S.Z. Salmasi; S. Nasrullahzadeh; N. Najafi and K. Ghassemi-Golezani (2010). Effects of foliar application of micronutrients (Fe and Zn) on flower yield and essential oil of chamomile (*Matricaria chamomilla* L.). *Journal of Medicinal Plants Research* Vol. 4(17), pp. 1733-1737.
- Niakan, M.; R.A. Khavarynejad and M.B. Rezaee (2004). Effects of different rates of NPK fertilizer on the leaf fresh weight, dry weight, leaf area and oil content of *Mentha piperita*, L. *Iranian Journal of Medicinal and Aromatic Plants Research*, 20 (2): 131-148.
- Saeid Zehab, S.; S. Behrouznajhad and K. Ghassemi-Golezani (2012). Effects of foliar application of Fe and Zn on seed yield and mucilage content of Psyllium at different stages of maturity. *International Conference on Environment, Agriculture and Food Sciences (ICEAFS'2012)* August 11-12, 2012 Phuket (Thailand).
- Said-Al Ahl, H.A.H. and Aber A. Mahmoud (2010). Effect of zinc and / or iron foliar application on growth and essential oil of sweet basil (*Ocimum basilicum* L.) under salt stress. *Ozean Journal of Applied Sciences* 3(1), pp. 97-111.
- Sandell, R. (1950). *Colorimetric determination of traces of metal* 2nd Ed. Inter since. Pub. Inc. New. York
- Shahram S. and K. Ordookhani (2011). Organic and bio fertilizers as a good substitute for inorganic fertilizers in medicinal plants farming. *Australian Journal of Basic and Applied Sciences*, 5(12): 1330-1333.
- Shilpa M. S. and K. N. Dhupal (2012): Influence of foliar applications of micronutrients on photosynthetic pigments and organic constituents of medicinal plant *Cassia angustifolia* Vahl. *Annals of Biological Research*, 3 (1): 520-526.
- Snedecor, G.W. and W.G. Cochran (1989). *Statistical methods*. 6<sup>th</sup> Ed. The Iowa state Univ. Press, Ames., Iowa. U.S.A.
- Sperenat, M. (1990): *Nitrogen fixing organisms*. P.5. Chapman and hall London.
- Suresh, K.D., G. Sneha; K.K. Krishn and C.M. Mool (2004). Microbial biomass carbon and microbial activities of soils receiving chemical fertilizers and organic amendments. *Arch. Agron. Soil Sci.*, 50: 641-647.
- Wojcik, P.; M. Wojcik and K. Klamkowski (2008). Response of apple trees to boron fertilization under conditions of low soil boron availability. *Sci. Hort.*, 116: 58-64.
- World Health Organization (WHO) (2008). "Traditional medicine" Fact sheet number: 134 (December)". <http://www.who.int/mediacentre/factsheets/fs134/en/>.
- Yassen A.A.; E.A.A. El-Nour and S. Shedeed (2010). Response of wheat to foliar spray with urea and micronutrients. *J Amer Sci* 6(9): 14-22.
- Youssef, A.S.M. (2009): Effect of some amino acids and mineral nutrients treatments on growth and productivity of rosemary plant (*Rosmarinus officinalis*, L.). *Annals of Agric. Sc., Moshtohor*, Vol. 47(1): 133-148.
- Zheljzakov, V.D. and P.R. Warman (2004). Source-separated municipal solid waste compost application to Swiss chard and basil. *J. Environ. Qual.*, 33: 542-552.

تأثير بعض معاملات التسميد ومعاملات التغذية بالعناصر الصغرى علي النمو والمحتوي الكيماوي لنبات الاشينيسيا  
ياسر عبد الفتاح عبد العاطي غطاس\*، وفاء حامد عبد الله\*\*  
\* قسم البساتين - كلية الزراعة - مشتهر - جامعه بنها - مصر .  
\*\* قسم النباتات الطبية والعطرية - مركز بحوث الصحراء - مصر .

اجريت تجربة حقلية بمزرعة خاصة بمنطقة الخطاطبة - محافظة المنوفية - مصر ومعمل قسم البساتين - كلية الزراعة بمشتهر - جامعه بنها لمدته عامين خلال موسمي ٢٠١٤ & ٢٠١٥ بهدف دراسة تأثير بعض معاملات التسميد المختلفة وقد تم استخدام سماد الكمبوست بمعدل ١٠ طن / للفدان كمقارنة وجرعه كامله من السماد المعدني المكون من النتروجين والفسفور والبوتاسيوم منفرده، (١٠٠%) وثلاث ارباع جرعه من السماد المعدني المكون من النتروجين والفسفور والبوتاسيوم + التسميد الحيوي المكون من النتروجين + الفوسفورين + الكمبوست بمعدل ١٠ طن / فدان كما تم استخدام نصف جرعه من السماد المعدني المكون من النتروجين والفسفور والبوتاسيوم + التسميد الحيوي المكون من النتروجين والفسفور + التسميد المعدني باستخدام النتروجين والفسفور والبوتاسيوم + التسميد الحيوي المكون من النتروجين والفسفور + التسميد العضوي (الكمبوست) بالاضافه الي استخدام ربع جرعه مع استخدام معاملات الرش بالعناصر الصغرى ممثله في عنصري الزنك والبورون بمعدل ١٠٠ جزء في المليون والتدخلات المختلفه بينهم وتأثيره علي النمو ومشتقات حامض الكافيك والكاميدات الكليه والمحتوي الكيماوي لنباتات الاشينيسيا. وقد اوضحت النتائج ان المعاملات المختلفه من التسميد ومعاملات الرش بالعناصر الصغرى بالاضافه الي التدخل بينهم ادت الي زياده معنويه كبيره في معاملات النمو المختلفه تحت الدراسه كارتفاع النبات ، عدد الافرع ، الوزن الطازج والجاف للعشب، وعدد الخلفات، وعدد الرؤوس الزهرية والوزن الطازج والجاف لها مقارنة بالكنترول في كلا الموسمين بالاضافه الي ذلك فان الزيادة في النمو الحادته والمتحصل عليها لنباتات الاشينيسيا باستخدام المعاملات المختلفه كانت مصحوبه بزياده واضحه في محتوى الاوراق من النتروجين والفسفور والبوتاسيوم للنباتات المعامله في كلا الموسمين . وعلاوه علي ذلك فان مشتقات حامض الكافيك والكاميدات الكليه قد زادت زياده واضحه باستخدام معاملات التسميد المختلفه ومعاملات الرش بالعناصر الصغرى فضلا عن التدخلات المختلفه بينهم وخاصة عند استخدام معاملة التدخل بين ثلاث ارباع جرعه من السماد المعدني باستخدام النتروجين والفسفور والبوتاسيوم + الكمبوست بمعدل ١٠ طن / فدان + التسميد الحيوي المكون من النتروجين + الفوسفورين والزنك والبورون بتركيز ١٠٠ جزء في المليون. وبناء علي نتائج هذه الدراسه ، فانه يفضل تسميد نباتات الاشينيسيا بثلاث ارباع جرعه من التسميد المعدني النتروجين والفسفور والبوتاسيوم (٧٥%) + ١٠ طن كمبوست / للفدان + التسميد الحيوي المكون من النتروجين + الفوسفورين ، مدعما بالرش بعنصري الزنك او البورون عند تركيز ١٠٠ جزء في المليون لزياده نمو النبات والمحتوي الكيماوي والذي يؤدي في النهايه الي الحصول علي منتج ءامن ذات جوده عاليه مناسب للتصدير وامن علي صحة الانسان.