

## **MANAGEMENT OF KHABRA BEETLE ADULTS, *Trogoderma granarium* (EVERTS) ON WHEAT GRAIN USING SOME SAFE PHYSICAL AND CHEMICAL MEASURES OF CONTROL UNDER LABORATORY CONDITIONS**

**Ibrahim, Sahar I. A.**

**Pesticides Dept., Faculty of Agriculture, Kafr-El-Shiekh University**

### **ABSTRACT**

○Laboratory experiments was conducted to evaluate the adverse action of six plant powders (red pepper, nigella, kidney bean, green peas, cumin and clove), three chemical insecticides (pirimiphos methyl, spinosad and lufenuron) and four environmental temperatures (10,15,25 and 40 °C) against the insect species under study, *Trogoderma granarium* using an incubator. Two types of packages, glass jars and plastic bags were also used for investigating the effect of package kind on the tested insect under the four tested temperatures

The capacity of packaging were studied to prevent tested insect from infesting wheat grain, two assay methods were used, package of one layer and package of two layers. Percentage of damage was estimated. The results revealed that the two layer bags greatly reduced the damage of wheat grain by *T. granarium* adults compared to that of one layer only. Results demonstrated that the high mortality percentages of *T. granarium* adults were found in polyethylene bags after 3 and 7 days of exposure periods at 40, 25, 15 and 10 °C respectively. The highest mortality was found with 10 °C followed by 40, 15 and 25 °C.

The effect of three tested factors together (kind of package, temperatures and insecticides) was higher than that of the two factors of package and temperatures only. The obtained data also revealed that pirimiphos-methyl had the highest effect on *T. granarium* adults compared to spinosad and lufenuron. Lufenuron produced the least effect on the tested insect species among the tested insecticides.

### **INTRODUCTION**

Cereal crops are subjected to attack by the khapra beetle (*Trogoderma granarium* Everts) which is one of the most serious insect pests in tropical and subtropical regions of Asia and Africa (Salunkhe *et al.*, 1985 and Viljoen, 1990). Synthetic insecticides have been used since the 1950s to control stored-products insects (Subramanyam and Hagstrum 1995). Pirimiphos-methyl (Actellic) and malathion were used as a potential protectants for grain and their products against stored product insects

Resistance to traditional insecticides used in stored-product treatments as well as the public's growing demand for residue free products has led researchers to evaluate alternative methods for the control of insects that infest stored products. Cheap and effective method are needed for reducing *T. granarium* damage. Damaged grain has reduced nutritional value, low germination percentage, reduced weight and lowered market value.

Protection of stored products generally involves mixing grains with protectants made up of plant materials.

Fresh, dry or processed plant materials can be applied as insecticides or to repel the pest insects. Plant powders can have a protective effect on the seeds based on several mechanisms. In addition, to direct toxic effects, the plant material may produce odours that repel or confuse the adult beetles, which could prevent invasion or cause emigration from the treated stock. Many plants have been tested in laboratories for their toxic effects on storage beetles, but few of them were tested for their repellent effect. (Okonkwo and Ewete, 2000, Fields *et al*, 2001, Ofuya and Salami, 2002, Umoetok and Gerd, 2003, Abdullahi and Muhammad, 2004, Emeasor *et al*, 2005, Echezona, 2006 and Selase and Getu (2009) .

Spinosad, a broad-spectrum insecticide of low mammalian toxicity, is based on metabolites of the actinomycete *Saccharopolyspora spinosa* Mertz and Yao (1990). Spinosad, the first active ingredient in the naturalyte class of insect control product was introduced by Dow Agrisciences for control of Lepidopterous pests in cotton in 1997 under the trade name of tracer, spinosad is a naturally occurring mixture of two active components, spinosyn A and spinosyn D (Salgado, 1998).

Lufenuron is a benzoylphenyl-urea (BPU) class of insecticides, which acts as a chitin synthesis inhibitor (CSI). Dean *et al.* (1998), CSI have gained significant popularity due to their low mammalian toxicity and absence of mutagenic and teratogenic effects on warm-blooded animals (Mondal and Parween, 2000)

Physical control measures such as high temperature or packing materials are needed. Since most stored product insects cannot tolerate extreme temperature, heating and cooling are logical approaches to insect control. The insect become inactive and eventually die at a temperature below 12C°. Lowt emperature is also important in maintaining seed viability (Zewar, 1993, Ayad and Alyousef, 1994, Ali *et al*, 1997, Dohino *et al* (1999), Lale and Vidal, 2003, Arthur, 2006, Burges, 2008 and Xial *et al* (2009).

In recent years, Plastic film also protects food from changes in composition (Fernandez, 2000).

**Consequently, this study aims to investigate the following points:**

- The toxic activity of nine compounds: one chemical insecticide organophosphorus (pirimiphos-methyl), one biocide [tracer (spinosad)], one insect growth regulator [match (Lufenuron)], and six botanical plant dusts (red pepper, nigella, green peas,cumin, clove and kidney bean).
- Evaluation of efficacy of temperature against the tested insect.
- To evaluate the combined effect of temperature with packages
- To evaluate some of packing materials.
- To evaluate the different packaging manners.

## MATERIALS AND METHODS

### Insects used:

#### **(*Trogoderma granarium*) :**

*T. granarium* adults tested in this study were continuously reared free of any insecticidal contamination for several years at 30 + 2°C and 70 + 5 R.H at Department of Stored Product Pests, Plant Protection Research Institute, Sakha Agriculture Research Station. The subculture was maintained under the same conditions, 200-400 adults from the previous culture were added in 850 ml glass jars containing 400 gm of crushed and whole wheat grain (*Triticum aestivum* L.) (variety of Sakha 69 with initial moisture content of 14%) as a culture medium. The jars were covered with muslin cloth to be tightly closed and was ranged by rubber bands. Newly emerged adults (1-2 weeks old) were used in the experiments

### Chemicals used:

#### **Pirimiphos-methyl (actellic, 50%EC).**

**Chemical name:** O-2 diethyl amino-6-methyl-pyrimidin-4-yl-o,o-dimethyl phosphorothioate.

#### **Spinosad (tracer,24%SC):**

**Chemical name:** Mixture of spinosyn A and spinosyn D. Spinosyn A: (2R, 3as, 5bs, 9s, 13s, 14R, 16bR)-2-(6-deoxy-2,3,3,4, Tri-O-methyl- $\alpha$ -L-Mannopyranosyloxy)-13-(4-dimethylamino-2,3,3a, 2, 3, 4, 6-tetra-deoxy-B-D-erythro-pyranosyloxy)-9-ethyl-2,3,3,a 5a, 5b, 6, 7, 9, 10, 11, 12, 13, 14, 15, 16a,16b-hexadecahydro-14-methyl-1H-8-oxacyclododeca[b]as-indacene-7, 15-dione.

Spinosyn D: (2s, 3aR, 5as, 5bs, 9s, 13s, 14R, 16as, 16bR)-2-(6-deoxy-2,3,4-tri-o-methyl- $\alpha$ -L-Mannopyranosyloxy)13-(4-dimethylamino-2,3,4,6-tetra-deoxy- $\beta$ -D-erythro-pyranosyloxy)-9-ethyl-2,3 3a, 5a, 5b, 6, 7, 9, 10, 11, 12, 13, 14, 15, 16a, 16b-hexadecahydro-4, 14-dimethyl-1H-8-oxacyclododeca[b]as-indacene-7, 15-dioine.

#### **Lufenuron®( match 5% EC)**

**Chemical name:** (RS)-1-[2,5-dichloro-4-(1,1,2,3,3,3-hexafluoro-propoxy) phenyl]-3-(2-6, difluorobenzoyl) urea.

### Botanical dusts:

Six plant materials were used as dusts against stored product insect (*T.granarium*).The target plant part was dried and finally ground into a fine powder in an electrical blender for five minutes. The powder was thoroughly sieved (300 mech) for obtaining the dust. Mixing of media bioassay method was used.10 gm of whole wheat grain was mixed with the desired amount of a plant powder in 1/4Kg glass jar and ten unsexed insects of *T. granarium* adults were introduced in glass jar which contains the treated wheat grain and tightly covered.% correct mortality was calculate according control treatment.

**Effect of temperature on the *T. granarium***

A laboratory experiment was conducted to evaluate the adverse action of four environmental temperatures (10, 15, 25 and 40°C) against *T. granarium*, where the temperatures were maintained using an incubator. Two types of packages, glass jars and plastic bags were also used for investigating the effect of package kind on the tested insects under the four tested temperatures. Ten adult of *T. granarium* were introduced into glass jar or plastic bag. Glass jars or plastic bags which contain 10 g of whole wheat (untreated) and 10 adult insects of *T. granarium* were kept in an incubator at 10, 15, 25 and 40°C) for 3 and 7 days. mortality Percentages were recorded and corrected by Abbott's formula (1925).

**Combined effect of temperature with used insecticides on *T. granarium***

For each tested insecticide (Pirimiphos methyl, spinosad and Lufenuron), the suitable required series of concentrations were prepared by adding the desired amount of each material in distilled water, then 10 g of sterilized wheat grain were treated with the above mentioned concentrations. Ten adult of *T. granarium* (1-2 weeks old) were separately introduced into either glass jars or plastic bags each contain the treated wheat grain . Both glass jars or plastic bags were kept in an incubator at different rates of temperature, 10, 15, 25 and 40°C. Treatments which have no insecticide were used as control. Three replicates of each concentration were used for each incubated period (3 and 7 days). After each incubated period, samples of treatments were withdrawn and percent of corrected mortality was calculated .LC<sub>50</sub>'s were estimated and used for comparison between the potential activities of the tested toxic agents at the different degrees of temperature.

**Effect of packaging on % damage infestation:**

To prevent invasion of insect pests into stored grains and their products, the capacity of packaging were studied to prevent insects from infesting food products.

Four plant powders (red pepper, nigella, kidney bean and green peas)were selected based on their efficacy to carry out further studies in combination with the backing materials.

Two assay methods i.e. the first, was the common packaging method where the wall of package is one layer. The second, was uncommon method where the wall of package contains two layers, the internal face of the two layers was treated with the selected plant powders. Untreated internal faces of package material was used for comparison.

**A. The first method: (package of one layer):**

In this experiment polyethelene package was used. ½ kg polyethylene bags contain treated wheat grain with the powders of the tested plants (LC<sub>50</sub> of each) were used. For check treatment, bags containing untreated wheat grain were used. The all treatments were transferred into glass jars (50 x 50 x 30 cm), number of 300 adult insects of *T. granarium* was introduced into the used glass jars and covered with lids. After 45 days of treatment, the bags were inspected and the percent of damage infestation was estimated.

**The second method assay (package of two layers):**

In this uncommon method, packages of the polyethelene consist of two layers were used. The internal side faces of the two layers were treated with the LC<sub>50</sub> of each plant powder which prior produced in the experiment of toxicity. ½ kg polyethylene bags which have treated wall of two layers contain untreated wheat grains were used. For control treatment, another packages (have wall of two layers) free of the plant powder between the two layers of package contain untreated wheat grain were used. The treated bags or control replicated three times. The all treatments were introduced into glass jars (50 x 50 x 30) contain 300 adult insects of *T. granarium*. The jars were covered with their lids, 45 days post-treatment, the bags were opened and wheat grains were inspected to determine the damage percentage using the following equation:

$$\% \text{ Damage} = \frac{\text{Number of infested wheat grain}}{\text{Number of total wheat grain}} \times 100$$

**RESULTS AND DISCUSSION**

**Efficacy of the tested chemical insecticides and plant powders against *T. granarium* adults**

Based on results obtained in Tables (1and2) red pepper powder with LC<sub>50</sub> of 0.13% w/w had the highest effect on *T. granarium* adults followed by green peas, kidney bean, nigella, cumin and clove, while pirimiphos-methyl was the strongest insecticides compared to spinosad and match with LC<sub>50</sub> of 2.49, 19.23 and 82.47 ug/g wheat grain, respectively.

**Table (1): Toxicity of the tested plant powders to *T. granarium* exposed to treated media (after 3 days).**

Plant powders	LC <sub>50</sub> w/w%	Slope value	C.L		Toxicity index
		S.V	Lower	Upper	
Red pepper	0.13	0.93	0.03	0.23	100
Cumin	0.6	0.24	0.36	0.90	21.67
Clove	0.85	1.59	0.70	1.04	15.29
Nigella	0.36	9.70	0.14	0.55	36.11
Green peas	0.18	1.01	0.07	0.28	72.22
Kidney bean	0.24	1.04	0.12	0.35	54.17

**Table (2): Toxicity of the tested insecticides to *T. granarium* exposed to treated media (after 24 hours).**

Toxicants	LC <sub>50</sub> µg/g	Slope value	C.L		Toxicity index
		S.V	Lower	Upper	
Pirimiphos-methyl	2.49	2.31	2.13	2.87	100
Spinosad	19.23	1.40	10.34	27.1	12.95
Lufenuron	82.47	1.58	63.47	100.76	3.02

**Efficacy of the tested chemical insecticides and plant powders against *T. granarium* adults.**

From the present results it is observed that all dry plant powders achieved good protection for wheat grain against adults of *T. granarium*. Plant products are seem to be an effective means of (appropriate technology) suitable to small hold farmers for protecting stored grain from insect damage. There is now overwhelming evidence that many plant species exert diverse biological effects on insects, i.e killing, attracting, repelling, feeding dekering, growth inhibiting and sterilizing effects (Abbassy, 1981 & 1982, Saleh, 1984 & 1986 and Meisner Nemny, 1992 and Ursula *et al* (2009).

Paul (2006) studied the effect of yellow field pea (*Pisum sativum*) fractions that were mainly protein (50%), fibre (90%) as starch (85%) were obtained from a commerce hial pea mill and mixed with wheat kernels or wheat flow against nine stored grain beetles. He found that the all fractions tested had toxic action against the all tested insect beetles for 8 months at 30 C<sup>o</sup> , 70 % r. h.

Spinosad is a (soft) chemical in the sense that it degrades rapidly in the environment and has minimal impact on non-target organisms (Thompson *et al* 2000). Although spinosad breaks down quickly in sunlight, limited published data suggest that spinosad in stored grain will be stable and loss of efficacy will be negligible (Fang *et al* 2002).

Subsequently, spinosad many exploited for insect control of stored product insects and is likely to be a safe alternative of chemical insecticide, pirimiphos-methyl. The high lighted in this study is that concerned in use powders of kidney bean and green peas which demonstrated great detrimental effect on *T. granarium* adults, these it is may be successful adoption for control of stored grain insects .

**Effect of packaging process on adults of *T. granarium*:**

Results obtained showed that polyethylene treatment damage ratio ranged from 3.0% (green peas) to 8.0 %(red pepper) compared with that of untreated which had 13.0% damage.

Based on the infestation percentage green peas was the most effective plant powder followed by nigella, kidney bean and red pepper either with one layer or two layer polyethelene bags (Table 3).

**Table (3): Infestation percentage of *T. granarium* adults after 45 days of exposure to treated wheat grain.**

Plant Powders	Treatments			
	Treated		Untreated	
	One layer	Two layers	One layer	Two layers
	Polyethylene	Polyethylene	Polyethylene	Polyethylene
Red pepper	8.0	6.0	13.0	12.0
Nigella	3.0	2.0	13.0	12.0
Kidney bean	8.0	3.0	13.0	12.0
Green peas	3.0	2.0	13.0	12.0

The treatment of package only had level of damage higher than that of in combination with plant powders. In conclusion, bags of two layers were the better treatment in reducing the damage level of adults of *T. granarium* if compared with that of customary bags which has one layer only. The sacs which has a wall of two layers contain plant powder between their inside faces had the highest effect in minimizing the damage level compared with that of no plant powders. The use of the second type (of two layers with plant powders) was found to be a rational mean in threat cease of stored grain insects, in addition, this method of application reduce the hazard of insecticidal contamination or toxicant materials which do not direct reach to stored products.

The level of damage adults of *T. granarium* with polyethylene bags may depends on the chemical composition of polyethylene, the numbers of holes in mm<sup>2</sup> and the number of wall layers of bag (Highland 1991)

These advantages of two layer sacs which contain plant powders in their internal face are likely to be an important factor in integrated pest management programs. Insect pests may be prevented from entering large or small quantities of stored grain by sealing or packaging.. It will probably depends on several factors including pest pressure, size of holes in the barrier, and tolerance of infestation. Four small bulks of up to a few kilograms, packaging and sealing are widely used to prevent invasion of insect pests into cereal grain and products. There has been extensive work on the capacity of packaging to prevent insects from infesting food products and there are several reviews on this topics (Wohlgemuth, 1979; Newton, 1988; Highland, 1978, 1981 and 1991)

Packaging provides a physical barrier that prevents or impedes infestation by insects. There are three factors that determine if a commodity will be infested: insect species, packaging material, and commodity packaged. Insects vary in their capacity to penetrate packaging, *Trogoderma variable* (Ballion) are all capable of penetrating intact packages. *R. dominica* is also capable of penetrating but is rarely found in packaged foods. Other stored-product insects (e.g. *T. castaneum*, *T. confusum*, and *O. surinamensis*) must have an opening to enter most packaged foods (Highland, 1991). Holes larger than 2 mm<sup>2</sup> will allow most stored-product adult insects to enter packages, whereas holes smaller than 0.3 mm<sup>2</sup> will prevent entry (Cline and Highland, 1981). Packaging materials differ in their capacity to prevent penetration. Listed in order of decreasing ease of penetration they are: cellophane, polyethylene, paper, polyvinyl chloride, aluminum foil, polyester and polycarbonate (Cline, 1978). Insecticides can be inserted into the packaging materials to render them more resistant insects. Multiwall packages are necessary to prevent the migration of the pesticide into the commodity.

The widespread protection practice at the farmer level in West and Central Africa regions remains that of mixing grains with locally available natural substances such as wood-ash or dried plant leaves (Chiranjeevi, 1991; Rajapakse and Van Emden, 1997) which they consider less hazardous and of no cost.

**Effect of temperature and its combination with the tested chemicals in different storage equipments against adults *T. granarium*:**

The principle factors that lead to the proliferation of insects in stored grain are temperature and humidity. In general, the condition suitable for most insect pests is around 30°C, with a relative humidity of 40 to 80 percent. Above 40°C, all species will eventually be killed, while reproduction ceases at temperature below 20°C, with dormancy induced at temperature below 10°C.. Alternatively, brief, high-temperature treatment of grain has been found to disinfest all stages of *S. granarius* in wheat (20 minutes at 70°C) (Zewar, 1993) and other storage insects (two minutes at 55°C (Lapp *et al.*, 1986).

Therefore, laboratory treatments were set to investigate the efficiency of together three factors for suppressing the damage of the tested insect species. The studied factors were, four various temperatures (10, 15, 25, 40°C), two package types (polyethylene, glass jars) and three insecticides (pirimiphos methyl, spinosad and lufenuron ). Two treatments were conducted: in the first treatment, *T. granarium* adults were exposed to untreated media either in glass jars or in polyethylene bags under the four tested temperatures. The second treatment was exposing the *T. granarium* adults to treated media by the tested insecticides either in polyethylene bags or in glass jars under the four tested temperatures. Results of the first treatment in Table (4) demonstrated that the high percent of mortalities (83.3, 100), (53.3, 73.3), (76.7, 90) and (90, 93.3) of *T. granarium* adults were found in polyethylene bags after 3 and 7 days of exposure periods at 40, 25, 15 and 10°C, respectively. The highest mortality was found with 10°C followed by 40°C, 15°C and 25°C especially with 3 days post treatment either with package type or with the tested insect species.. Results in Table (4) showed that the mortality percentages increased with the increasing of exposure period with the all tested temperatures, and the two types of package. While, the highest effect of polyethylene bags on the tested insects may be due to the different properties between glass and polyethylene, in addition, the effect of package volume where the volume of polyethylene bags was smaller than that of glass jars during the experiment, this means that the amount of oxygen in polyethylene bags was less than that of glass jars

**Table (4):Mortality percentage of *T. granarium* at indicated temperatures after 3 and 7 days of treatment (untreated media)**

Type of package	40°C		25°C		15°C		10°C	
	3 days	7 days	3 days	7 days	3 days	7 days	3 days	7 days
Glass jars	36.7	73.3	13.3	16.7	46.7	70	46.7	66.7
Polyethylene	83.3	100	53.3	73.3	76.7	90	90	93.3

Results in Tables (5-6) revealed that the effect of the three factors together (temperature, package and insecticides) was higher than that of the two factors of package and temperature only. Data obtained also revealed that pirimiphos-methyl had the highest effect on *T. granarium* compared to



spinosad and match Also. Also, the results demonstrated that the toxic effect increased with the increasing of concentrations at the different levels of temperature

*T. granarium* (Everts) is a common pest of grain and other stored products in the tropics and subtropics. It has been distributed worldwide by trade but requires hot conditions for rapid breeding. Moderately low temperatures slow down development and induce diapause, so that cooling grain is an effective, ecofriendly method to control of *T. granarium* (Borges and Burrell, 1964).

**Table (5): Comparative and combined effect of temperatures, insecticides and glass jars on *T. granarium* adults mortality after 24 h of treatment**

Tested parameters	Conc.	40°C	25°C	15°C	10°C
T+g+p	10 ppm	96.7	86.7	86.7	96.7
	5 ppm	90.0	76.7	83.3	93.3
	2.5 ppm	80.0	60.0	76.7	86.7
	1.25 ppm	63.3	26.7	46.7	70.0
T + g + S	150 ppm	83.3	73.3	76.7	86.7
	100 ppm	73.3	60.0	66.7	80.0
	50 ppm	50.0	36.7	43.3	63.3
	25 ppm	33.3	20.0	20.0	43.3
T + g + M	400 ppm	83.3	63.3	80.0	93.3
	200 ppm	76.7	53.3	60.0	83.3
	100 ppm	60.0	36.7	40.0	66.7
	50 ppm	40.0	20.0	46.7	46.7
Control		0	0	0	0

T = Temperature, G =glass jar, P = Pirimiphos-methyl, S = Spinosad, M = Match

**Table (6): Comparative and combined effect of temperatures, insecticides and polyethylene on *T. granarium* adults after 24 h of treatment.**

Tested parameters	Conc.	40°C	25°C	15°C	10°C
T+Po+p	10 ppm	96.7	86.7	93.3	100.0
	5 ppm	93.3	70.0	83.3	96.4
	2.5 ppm	76.7	63.3	73.3	86.7
	1.25 ppm	60.0	36.7	53.3	76.7
T + Po + S	150 ppm	90.0	66.7	83.3	93.3
	100 ppm	86.7	56.7	73.3	86.7
	50 ppm	63.3	43.3	53.3	66.7
	25 ppm	43.3	26.7	33.3	46.7
T + Po + M	400 ppm	86.7	70.0	76.7	93.3
	200 ppm	76.7	53.3	60.0	86.7
	100 ppm	63.3	36.7	43.3	76.7
	50 ppm	46.7	20.0	26.7	63.3
Control		0	0	0	0

T = Temperature, Po=Polyethylene bags, P = Pirimiphos-methyl, S = Spinosad, M = Match.

Burges (2008), determined the low temperature limit for breeding of the khapra beetle *T. granarium* at constant temperatures in the laboratory. The length of female adult life, oviposition, egg incubation period and duration of larval plus pupal development were observed at 25, 22.5, 20 and 17.2°C, respectively at 80% R.H. The results show that breeding is slow at 25°C, very slow at 22.5°C and population decline at (20°C and below). Thus cooling of products to 20°C and retention at or below this temperature is a sound, safe, environmental friendly method for controlling of this pest. He also claimed that mortality (sex unknown) increased from 30.3% at 24.9°C to 100% at 17.2°C. These findings are in agreement with our results where the highest mortality of *T. granarium* achieved by 10°C.

In fact, 15°C is now the maximum storage temp. recommended by the Home-Grown Cereals Authority, to ensure control of other pest species (Burges, 2008). A technique that has been used successfully for many years against stored-product pest is the use of extreme temperatures (Fields, 1992). Use of elevated temperatures in heat treatments has long been recognized as an effective strategy for managing stored product insects associated with food-processing facilities (Dean *et al.*, 1998). Nakakita and Ikenaga (1997) studied the low temperatures to control insect pests in storage rice in Japan. They investigated the action of low temperature on the physiology of *S. oryzae* and *S. zeamais* the major insect pests on rice. They found that oxygen consumption by the adults of both species was reduced as temperature was decreased. Both species exhibited a biphasic decline in respiratory rate as the temperature was reduced, giving sharp fall in respiratory response between 30 and 20°C and a moderate fall between 15 and 5°C. Population increase of *S. oryzae* was completely inhibited at 15°C, while a small number of *S. zeamais* emerged. Xial *et al.* (2009) examined the effect of four constant temperature (20, 24, 28 and 32°C) on the physiological properties (survival, development time, sex ratio, fecundity and longevity of females) in the laboratory at 85% RH. The total development time, longevity shortened as temperature increased.

## REFERENCES

- Abbassy, M.A. (1981). Insecticidal and synergistic volatile oils isolated from certain food and medicinal plants. Proc. 4<sup>th</sup> Arab. Pesticide Conf. Tanta Univ., IIIA: 409-414.
- Abbassy, M.A. (1982). Naturally occurring chemicals for pest control. III. Insecticidal and synergistic alkaloid isolated from *Schinus terbinthifolius* Reddi Med. Fac. LandBouww, Rijksuniv. Gent. 4712: 617-626.
- Abbott, W.W. (1925). A method of computing the effectiveness of an insecticide. J. Econl. Entomol., 18: 265-267.
- Abdullahi, Y.M. and S. Muhammad (2004). Assessment of the toxic potentials of some plant powders on survival and development of *Callosobruchus maculatus*. African Journal of Biotechnology. Vol. 3(1): 60-62.
- Abdullahi, Y.M. and S. Muhammad (2004). Assessment of the toxic potentials of some plant powders on survival and development of *Callosobruchus maculatus*. African Journal of Biotechnology. Vol. 3(1): 60-62.

- Ali, M.F.; E.F.M. Abdel-Reheem; and H.A. Abdel-Rahman (1997). Effect of temperature extremes on the survival and biology of the carpet beetle, *Attagenus fasciatus* (Thunberg) (Coleoptera: Dermestidae). J. Stored Prod. Res. 33(2):147-146.
- Arthur, F.H. (2006). Initial and delayed mortality of late-instar larvae, pupae and adults of *Tribolium castaneum* (Coleoptera: Tenebrionidae) exposed at variable temperatures and time intervals. J. of Stored Prod. Res. V. 42: 1-7.
- Ayad, F.A. and E.F. Alyousef (1994). Effect of temperature on the efficiency of some insecticides against the cowpea weevil *Callosobruchus maculatus* (Fab.) (Coleoptera Bruchidae). Bulletin of the Entomological Society of Egypt, Economic Series. 14: 329-385.
- Burges, H.D. (2008). Development of the khaprabeetle, *Trogoderma granarium*, in the lower part of its temperature range. J. of Stored Prod. Res.. 44: 32--35.
- Burges, H.D. and N.J. Burrell, (1964). Cooling bulk grain in the British climate to control storage insects and to improve keeping quality. J. Sci. Food and Agric., 15: 32-500.
- Chiranjeevi, C.H. (1991). Efficacy of some indigenous plant materials and ashes on the percentage of damaged grains, percentage of protection and viability of green gram seed infested by pulse beetle, *Callosobruchus chinensis* (L.). Bulletin of Grain Technology, 29: 84-88.
- Cline, L.D. (1978). Clinging and climbing ability of larvae of eleven species of stored product insects on nine-flexible packaging materials and glass. J. Econ. Entomol., 71: 689-691
- Cline, L.D. and H.A. Highland (1981). Minimum size of holes allowing passage of adults of stored-product coleopteran. J. Georgia Entomol. Soc. 16: 525-531.
- Dean, S.R.; R.W. Meola; S.M. Meola; H.S. Bhattar and R. Schenker (1998). Mode of action of lufenuron on larval cat flea: (Siphonaptera: Pulicidae). J. Med. Entomol. 35(5): 720-724.
- Dohino, T.; S. Masaki; I. Matsuoaka; M. Tanno and T. Takano (1999). Low temperature as an alternative to fumigation for disinfecting stored products. Res. Bull. of the Plant Protection. Service, Japan, No.35. 5-17.
- Echezona, B.C. (2006). Selection of pepper cultivars (*Capsicum* spp.) for the control of bruchids *Callosobruchus maculatus* (F.) on stored cowpea (*Vigna unguiculata* (L.) Walp) seeds, African. J. of Biotechnology. 5(8): 624-628.
- Emeasor, K.C.; R.O. Ogbuji; S.O. Emosairue (2005). Insecticidal activity of some seed powders against *Callosobruchus maculatus* (F.) (Coleoptera: Bruchidae) on stored cowpea. Zeitschrift-fur-Pflanzenkrankheiten-Und-Pflanzenschutz. Vol. 112(1): 80-87.
- Fang, L.; S. Bhadriraju and D. Sean (2002). Persistence and efficacy of spinosad residues in farm stored wheat. Journal of Economic Entomology ISSN. 0022-0493, Vol. 95(5): 1102-1109.
- Fernandez, M. (2000). Review: active food packaging. Food Science and Technology International 6: 97-108.
- Fields (2006). Effect of *Pisum sativum* fractions on the mortality and progeny production of nine stored-grain beetles. J. Stored products Research 42, 86-89

**Ibrahim, Sahar I. A.**

- Fields, P.G. (1992). The control of stored-product insects and mites with extreme temperatures. *J. Stored Prod. Res.* 28: 89-118.
- Fields, P.G.; Y.S. Xie, and X. Hou (2001). Repellent effect of pea (*Pisum sativum*) factions against stored-product insects. *J. Stored Prod. Res.* 31: 359-370.
- Highland, H.A. (1978). Insect resistance of food packages. A Review. *J. Food Process. Preserv.* 1: 123-130.
- Highland, H.A. (1981). Resistant barriers for stored-product insects. pp. 41-46 in *CRC Handbook of Transportation and Marketing in Agriculture*. E.E. Finney (Ed.) 1, CRC Press, Boca Raton, FL.
- Highland, H.A. (1991). Protecting Packages against insects. pp. 345-350. In: *Ecology and Management of Food-Industry Pests*, J.R. Corgham (ed.) Assoc. Official Anal. Chem. Arlington, VA.
- Lale, N.E.S. and S. Vidal (2003). Effect of constant temperature and humidity on oviposition and development of *Callosobruchus maculatus* (F.) and *Callosobruchus subinnotatus* (Pic.) on bambara groundnut *Vignan subterranean* (L.) Verdcourt. *J. of Stored Prod. Res.* 39(5): 459-470.
- Lapp, H.M.; F.J. Madrid and L.B. Smith (1986). A continuous thermal treatment to eradicate insects in stored wheat. Paper 86-3008. *Am. Soc. Agric. Eng., St. Joseph., MI*, 14 pp.
- Meisner, J. and N.E. Nemny (1992). The effect of margosan-O on the development of the Egyptian cotton leafworm, . *J. of Stored Prod* 28:119-129
- Mertz, P.P. and R.C. Yao (1990). *Sacchropolyspora spinos* sp. nov. isolated from soil collected in a sugar rum still. *Int. J. Sust. Bacteriol.* 40: 34-39.
- Mondal, K.A.M. and S. Parween (2000). Insect growth regulators and their potential in the management of stored product insect pests. *Int. Pest Management Rev.* 5: 255.
- Nakakital, H. and H. Ikenaga (1997). Action of low temperature on physiology of *Sitophilus zeamais* motschulsky and *Sitophilus oryzae* (L.) (Coleoptera: Curculionidae) in rice storage. *J. of Stored Prod. Res.* V 33: 31-38.
- Newton, J. (1988). Insects and packaging a review. *Int. Biodet.* 24: 175-187.
- Ofuya, T.I. and A.Salami (2002). Laboratory evaluation of different powders from *Dennetia tripetala* Bok as protectant against damage to stored seeds cowpea by *Callosobruchus maculatus* (F.). *J. of Sustainable-Agriculture and the Environment.* 4(1): 36-41.
- Ofuya, T.I. and A.Salami (2002). Laboratory evaluation of different powders from *Dennetia tripetala* Bok as protectant against damage to stored seeds cowpea by *Callosobruchus maculatus* (F.). *J. of Sustainable-Agriculture and the Environment.* 4(1): 36-41.
- Okankwo, E.V. and F.K. Ewete (2000). Comparative evaluation of *Dennetika trippetala* seed powder, pirimiphos-methyl and aluminum phosphide tablets against *Callosobrudhus maculatus*Fabricius (Coleopotera: Bruchidae) and their organoleptic effects on stored cowpeas in Nigeria. *Moor. J. of Agric. Res.* Vol. 1(1): 48-53.

- Okankwo, E.V. and F.K. Ewete (2000). Comparative evaluation of *Dennettika trippetala* seed powder, pirimiphos-methyl and aluminum phosphide tablets against *Callosobruchus maculatus* Fabricius (Coleoptera: Bruchidae) and their organoleptic effects on stored cowpeas in Nigeria. *Moor. J. of Agric. Res.* Vol. 1(1): 48-53.
- Paul G. Fields (2006). Effect of *Pisum sativum* fractions on the mortality and progeny production of nine stored-grain beetles. *J. Stored products Research* 42, 86-89 .
- Rajapakse, R. and H.F. Van Emden (1997). Potential of four vegetable oils and ten botanical powders for reducing infestation of cowpea by *Callosobruchus maculatus*, *C. chinensis* and *C. rhodesianus*. *J. Stored Prod. Res.*, 33: 59-68.
- Saleh, M.A. (1984). An insecticidal diacetylene from *Artemisia monosperma*. *Phytochem.*, 23(11): 2497-2498.
- Saleh, M.A. (1986). A desert plant from Egypt, *Anabasis setifera* an efficient natural factory of caryacrol and thymol. *J. Agr. Food Chem.*, 34(2): 192-194
- Salgado, V.L. (1998). Studies on the mode of action of spinosad insect symptoms and physiological correlates. *Pesticide Biochemistry and Physiology*, 60: 91-102.
- Salunkhe, D.K.; J.K. Chiavan and S.S. Kadam (1985). *Postharvest biotechnology of cereals*. CRC press, Boca Raton, Philippines.
- Selase, A.G. and E. Getu (2009). Evaluation of botanical plants powders against *Zabrotes subfasciatus* (Boheman) (Coleoptera: Bruchidae) in stored haricot beans under laboratory condition. *African Journal of Agricultural Research* 4(10): 1073-1079.
- Subramanyam, B. and Hagstrum, D.W., (1995). Resistance measurement and management. In : Subramanyam, B., Hagstrum, D.W. (Eds), *Integrated management of Insects in Stored Products*. Marcel Dekker Inc., New York, pp. 331 – 398.
- Thompson, G.D.; R. Dutton and T.C. Spraks (2000). Spinosad a case study: an example from a natural products discovery programme. *Pest. Manag. Sci.* 56: 696-702.
- Umoeok, S.B.A. and M.B. Gerd (2003). Comparative efficacy of *Acarus calamus* powder and two synthetic insecticides for control of three major insect pests of stored cereal grains. *Global. J. of Agric. Sci.*, Vol. 2(2): 94-97.
- Ursula V. Paul, Juma S. Lossini, Peter J. Edwards and Angelika Helbeck (2009). Effectiveness of products from four locally grown plants for the management of *Acanthoscelides oryzae* (Say) and *Zabrotes subfasciatus* (Boheman) (both Coleoptera : Bruchidae) in stored beans under laboratory and field conditions in Northern Tanzania. *J. Stored products Research*, 45, 2 97-1
- Viljoen, J.H. (1990). The occurrence of *Trogoderma* (Coleoptera: Dermestidae) and related species in southern Africa with special reference to *T. granarium* and its potential to become established. *J. Stored Prod. Research*, 26: 13-51.
- Wohlgemuth, R. (1979). Protection of stored foodstuffs against insect infestation by packaging. *Chem. Ind.*, 5: 330-334

- Xial, B.; D. Luo; Z. Zou and Z. Zhu (2009). Effect of temperature on the life cycle of *Aleuroglyphus ovatus* (Acari: Acaridae) at four constant temperatures. J. of Stored Prod. Res. V. 45: 190-194.
- Zewar, M.M. (1993). The use of high temperature for disinfecting wheat from *Sitophilus granarius* (L.) and cowpea *Callosobruchus maculatus* (F.) Egypt. J. Agric. Res., 71(3): 671-678

**ادارة حشرة خنفساء الصعید علي حبوب القمح باستخدام بعض الطرق الطبيعية والكيمائية  
الأمنة تحت الظروف المعملية  
سحر ابراهيم أحمد ابراهيم  
قسم المبيدات – كلية الزراعة – جامعة كفر الشيخ**

أجريت تجارب معملية لتقييم فاعلية أربعة درجات حرارة بينية (١٠ و١٥ و٢٥ و٤٠ درجة مئوية) ضد الحشرة موضع الدراسة وهي خنفساء الصعید باستخدام الحضان واستخدمت في الدراسة نوعين من العبوات (البرطمانات الزجاجية والأكياس البلاستيكية) لدراسة التأثير المشترك علي الحشرة موضع الدراسة تحت الأربعة درجات الحرارة المدروسة .  
تم دراسة مدي قدرة عملية التغليف لحماية حبوب القمح من الاصابة بالحشرات باستخدام طريقتين من طرق التقييم وهما  
١- التغليف في طبقة واحدة  
٢- أو التغليف في طبقتين وتم تقدير النسبة المئوية للاصابة في القمح بالحشرات الكاملة من خنفساء الصعید في الحالتين وقد أشارت النتائج الي الأتي:  
١- مستوي الاصابة تأثر بطريقة التغليف  
٢- أن النسبة العالية للموت من خنفساء الصعید وجدت في أكياس البولي اثيلين بعد ٣ و٧ أيام من التعرض علي ٤٠ و٢٥ و١٥ و١٠ درجة مئوية علي التوالي  
٣- ظهرت أعلى نسبة موت عند درجة حرارة ١٠ تليها ٤٠ و١٥ ثم ٢٥ درجة مئوية وأشارت النتائج الي أن تأثير ثلاث عوامل معا (العبوات ودرجات الحرارة والمبيدات) أكثر من تأثير عاملي التغليف والحرارة فقط-  
وأشارت الي أن البريميغوس ميثيل كان الأشد تأثيرا علي خنفساء الصعید بالمقارنة بمبيدي السباينوساد و الماتش وأظهر الماتش أقل تأثير علي الحشرة بالمقارنة بالمبيدات الأخرى المدروسة.

**قام بتحكيم البحث**

كلية الزراعة – جامعة المنصورة  
كلية الزراعة – جامعة كفر الشيخ

أ.د / علي عبد الهادي  
أ.د / عطيه يوسف قريظم