

SOME PHYSICAL AND CHEMICAL PROPERTIES OF TRADITIONAL HONEYS IN NEW DESERT REGIONS IN EGYPT

Fathy, H.M.¹; E.I. Haggag²; M.I. sanad³ and M.R.M. Abd EL-Dayem²

1 Dept. Plant Protection, Fac. , Agric., Mansoura University, Egypt.

2 Beekeeping Research Department, Plant Protection Res Institute (PPRI), Agric. Res. Center, Giza, Egypt.

3 Dept. Agricultural Chemistry, Fac., Agric., Mansoura University, Egypt.

ABSTRACT

Ten citrus and clover honeys samples as traditional honeys were collected from new desert regions in Egypt. Eight parameters were measured in clouding moisture, total soluble solid (T.S.S.), PH, Electric conductivity (EC), Free acidity, Lactone, Total acidity and hydroxy methyl furfural (HMF)

The results indicated that moisture was parallel and the average values in citrus and clover honeys were 20.33 and 19.25%, respectively. The average values of EC, T.S.S., pH, Free acidity, lacton, total acidity and HMF in citrus and clover honey samples were 80 and 77.5, 79.67 and 80.75, 3.92 and 3.83, 16.64 and 14.44, 8.85 and 9.7, 25.85 and 24.14, 1.96 and 1.32, respectively.

INTRODUCTION

Honey standardization is an important object to study. Every country in all over the world has its special honey standard, depending on the different environmental factors existing in each country. Surveys of floral honey composition have established that the three major components are fructose, glucose, and water (Doner, 1977). Honey is the sweet substance produced by honey bee from the flowers or from secretions of living parts of plants, which honey bee collected, transform and combine with specific substances of their own, store and leave in honey comb to ripen and mature. (Codex Alimentarius 1998). Chemically, honey is quite complex. it comprised primarily sugars; however, it contains many other potentially biologically active components, such as antioxidants, which display antimutagenic activity (Wang *et al.*, 2002). Recently, standardization of the honey depends on major nine tests have to be determined. They were reducing sugar content, sucrose content, fructose; glucose ratio, moisture content, water insoluble solids content, ash, acidity, diastase activity and hydroxy methyl furfural (H.M.F.) content. These tests need abig effort to study all honey types in the country regions.

The aim of this study were determination of physical and chemical properties of traditional honeys (citrus and clover) in new desert regions in Egypt. This work was carried out during they year 2010 and 2011 in Bee Research Department Plant protection Research Institute.

MATERIALS AND METHODS

Ten traditional honeys sample (citrus and clover) were collected from new desert regions (North Sinai and Ismailia and New Valley and Al Menofia and Al Dakahlia) as in table 1

Table 1: Honey types and their different locations

Sample No.	Honey types	Regions	Locations
1	Citrus (<i>Citrus spp.</i>)	North Sinai	Shiekh Zowied
2		North Sinai	Rafah
3		Ismailia	Ismailia
4		Ismailia	Fayed
5		Ismailia	Qantara East
6		Al Menofia	Ashmoun
7	Clover (<i>Trifolium alexandrinum</i>)	New Valley	Al Dakhlah
8		Ismailia	Ismailia
9		Al Menofia	Ashmoun
10		Al Dakahlia	Qlabasho

The physical properties, the moisture (%) was measured according to (A.O.A.C., 1995), the total soluble solids (T.S.S.) was measured according to (A.O.A.C., 1980), the include electrical conductivity (EC) was measured according to (vorwohl, 1964).

In relation to chemical properties, pH, free acidity, lacton and total acidity based on the method of (white *et al.*, 1962). The hydroxy methyl furfural (HMF) was measured according to (AOAC 980.23. 1980).

RESULTS AND DISCUSSION

Physical and chemical properties of Citrus honeys:-

Data in table (2) show that citrus honey moisture ranged between 19 - 21.5(%) with mean value of 20.33(%). Concerning the electrical conductivity (EC) the citrus honey ranged between 60-110(ppm) with a mean value of 80(ppm), where as total soluble solids (T.S.S.) ranged between 78.5- 81(%) with a mean value of 79.67%. Also in table (2), the chemical properties of Citrus honey sample indicated that, the pH value ranged between 3.74 - 4.41 with a mean value of 3.92. The free acidity ranged between 10.17 and 22.67 with a mean value of 16.64.

Table (2):- Physical and chemical properties of Citrus honey collected from new desert regions in Egypt.

Regions	Location	Sample No.	Moisture%	T.S.S.*	EC* ppm	pH	Free acidity ml.eq/kg	Lacton ml.eq/kg	Total acidity ml.eq/kg	HMF mg/kg
North Sinai	Shiekh Zowied	1	21.5	78.5	70	3.76	22.67	10.33	33	1.048
	Rafah	2	21.5	78.5	60	3.81	17.67	3.25	20.92	3.039
Ismailia	Ismailia	3	19	81	110	4.41	10.17	16.67	26.83	2.894
	Fayed	4	19.5	80.5	70	3.74	17.33	6	23.33	0.349
	Qantara East	5	20.5	79.5	110	3.78	19.5	9	28.5	1.597
Al Menofia	Ashmoun	6	20	80	60	4	12.5	7.834	20.33	2.815
Variation limits			19 - 21.5	78.5 - 81	60 - 110	3.74- 4.41	10.17 - 22.67	3.25 - 16.67	20.33 - 33	0.349- 3.039
Average values			20.33	79.67	80	3.92	16.64	8.85	25.85	1.96

T.S.S.* total soluble solid
EC*ppm Electric conductivity

The lacton content ranged between 3.25 and 16.67 with a mean value of 8.85. While the total acidity value (free acidity + lacton content) ranged between 20.33 and 33 with a mean value of 25.85. The content of citrus honey sample hydroxy methyl furfural (HMF) showed low values as the values ranged between 0.349 and 3.039 with a mean value of 1.96.

Physical and chemical properties of clover honeys:-

Data in table (3) show that clover honey moisture ranged between 18.75 and 20.25 (%) with mean value of 19.25(%). Concerning the electrical conductivity (EC) the citrus honey ranged between 60 - 90(ppm), where as total soluble solids (T.S.S.) ranged between 79.75 and 81.25(%) with a mean value of 80.75(%). Data presented in table (3), indicated the chemical properties of Clover honey sample. The pH value ranged between 3.7and 3.98 with a mean value of 3.83. The free acidity ranged between 11.5 and 20.25 with a mean value of 14.44.

Table (3):- Physical and chemical properties of Clover honey collected from new desert regions in Egypt.

Regions	Location	Sample No.	Moisture%	T.S.S.%*	EC* ppm	pH	Free acidity ml.eq/kg	Lacton ml.eq/kg	Total acidity ml.eq/kg	HMF mg/kg
New Valley	Al Dakhlah	7	19	81	60	3.85	12.33	2	14.33	2.121
Ismailia	Ismailia	8	18.75	81.25	80	3.98	13.67	11	24.67	1.425
Al Menofia	Ashmoun	9	20.25	79.75	90	3.79	20.25	8.625	28.88	1.362
Al Dakahlia	Qlabasho	10	19	81	80	3.7	11.5	17.17	28.67	0.369
Variation limits			18.75 – 20.25	79.75 – 81.25	60 – 90	3.7– 3.98	11.5 – 20.25	2 - 17.17	14.33 – 28.88	0.369 – 2.12
Average values			19.25	80.75	77.5	3.83	14.44	9.7	24.14	1.32

T.S.S.%* total soluble solid
 EC*ppm Electric conductivity

The lacton content ranged between 2 and 17.17 with a mean value of 9.7. While the total acidity value (free acidity + lacton content) ranged between 14.33 and 28.88 with a mean value of 24.14. The content of clover honey sample with hydroxy methyl furfural (HMF) showed low values. The values ranged between 0.369 and 2.12 with a mean value of 1.32.

The data in table (4) showed clear differences in free acidity values among the tested honey samples.

The free acidity in citrus honey followed by clover honey samples the highest mean value among the tested honey sample (mean 16.64 and 14.44 milli/kg), respectively. These results were not accepted by Egyptian standard (2005). The high value of lacton content were recorded in clover honey while the low values of lacton content were recorded in citrus. The values of total acidity in all tested honey samples were higher in the Egyptian standard (2005).

Citrus samples showed the highest HMF content (mean 1.96 mg/ kg)

Than clover honey (mean 1.32mg/ kg). This may be due to the effect of climatic conditions in different locations especially the high temperature in Rafah region.

Table (4):- Means of physical and chemical properties for all honey sample (Citrus and Clover).

Honey type	physical and chemical properties							
	Moisture%	T.S.S.%*	EC* ppm	pH	Free acidity ml.eq/kg	Lacton ml.eq/kg	Total acidity ml.eq/kg	HMF mg/kg
Citrus honey	20.33	79.67	80	3.92	16.64	8.85	25.85	1.96
Clover honey	19.25	80.75	77.5	3.83	14.44	9.7	24.14	1.32
Range	19.25 - 20.33	79.67 - 80.75	80	3.83 - 3.92	14.44 - 16.64	8.85 - 9.7	24.14 - 25.85	1.32 - 1.96
Mean	19.79	80.21	78.75	3.875	15.54	9.275	24.995	1.64

T.S.S.%*

EC*ppm

total soluble solid

Electric conductivity

Generally, the physical properties of the most honey samples were not in the range values as mentioned by the specification of the Egyptian honey standard 2005. This may be due to the abnormal practices occurred by the beekeepers towards their honey bee colonies during the honey flow, or gathering the honey from these colonies before complete ripening. Many authors discussed the physical properties.

Water content in Egyptian honeys was found by (Ibraliim *et al.*, 1977) to be 18 – 19.5 %, by (vorwohl *et al.*, 1989) to be 16.3 – 22.3 % with mean 18 % and by (Gomaa, 2002) to be 15.4 to 22 %.

REFERENCES

- A. O. A. C. (1980) Official methods of analysis. Washington, DC: Ass off. Ana. chem..
- A. O. A. C. (1995) Official methods of analysis. (16thed).Washington, DC: USA: Ass. Off. Ana. Chem..
- A. O. A. C. (1998) The 16th edition of the Official Methods of Analysis, 4th Revision, current through March, Washington, DC: USA: Ass. Off. Ana. Chem..
- Codex Alimentations Commission (1998) Recommended European Regional standard for honey (CAC/RS 12-1969).
- Doner, L. W. (1977) The sugars of honey-A review, J. Sci. Food Agric., 28, 443–456.
- Gomaa, W. M. K. (2002) studies on honey quality with special reference to the Egyptian honey standard, M.Sc. thesis, Fac. Agric., Alex. Uni. Egypt, 156pp.

- Ibrahim, S. H.; N. Z. Soliman and H. Wissa.(1977) studies on the properties of the major Egyptian honey types and on honey ripening Agric. Res. Rev., 55:125 -129.
- Vorwhol, G. (1964a) Die Messung der elektrischen Leitfähigkeit des Honig und die Verwendung der Messwerte zur Sortendignose und zum Nachweis von Verfälschungen mit Zuckerrückführung Honig, Bienenforschung, 7,37-47.
- Vorwhol, G.; M. S. Salem and M. E. Nour (1989) Chemical and Physical properties of Egyptian honeys. Proc. 4 Int. Conf. Apic. Trop. Climates, Cairo,:240 – 244.
- Wang, X. H.; Andrae, L. and Engeseth, N. J. (2002) Antimutagenic effect of various honeys and sugars against, J. Agric. Fd. Chem., 50, 6923-6928.
- White, J. W.; Riethof, M. L.; Sobers, M. H. and Kushnir, I. (1962) Composition of American honeys, U.S., Dept. Agric., Tech. Bull., 1-124.

بعض الخصائص الطبيعية والكيميائية للأعسال التقليدية في المناطق الصحراوية الجديدة في مصر

- حسن محمد فتحى^١ ، السيد إبراهيم حجاج^٢ ، مصطفى إبراهيم سند^٣ و محمد رمضان محمد عبد الدايم^٢
- ١- قسم وقاية النبات – كلية الزراعة – جامعة المنصورة – مصر
 - ٢- "قسم بحوث النحل – معهد بحوث وقاية النباتات – مركز البحوث الزراعية – الجيزة – مصر
 - ٣- قسم الكيمياء الزراعية – كلية الزراعة – جامعة المنصورة – مصر

أجرى هذا البحث لدراسة بعض الخصائص الطبيعية والكيميائية للأعسال التقليدية (البرسيم، الموالح) والمفروزة من المناطق الصحراوية الجديدة في محافظات (شمال سيناء – الإسماعيلية – المنوفية – الدقهلية – الوادي الجديد). أجريت التحاليل المختلفة في معمل تحاليل منتجات نحل العسل بقسم بحوث النحل بالجيزة.

ولقد أظهرت التحاليل النتائج التالية:-

بالنسبة لعينات عسل الموالح ، البرسيم كان متوسط كل من الرطوبة ٢٠.٣٣% - ١٩.٢٥% ، درجة التوصيل الكهربي ٨٠ - ٧٧.٥ و الأملاح الذائبة الكلية ٧٩.٦٧ - ٨٠.٧٥ ودرجة الـ pH ٣.٩٢ - ٣.٨٣ والحموضة الحرة ١٦.٦٤ - ١٤.٤٤ واللاكتون ٨.٨٥ - ٩.٧ والحموضة الكلية ٢٥.٨٥ - ٢٤.١٤ والهيدروكسي مثيل فورفورال ١.٩٦ - ١.٣٢ وذلك على التوالي.

قام بتحكيم البحث

كلية الزراعة – جامعة المنصورة
كلية الزراعة – جامعة الاسكندرية

أ.د / سمير صالح عوض الله
أ.د / اسامه محمد الانصارى