

A NEW PASTEURIZED NECTARS PREPARED FROM HUSK TOMATO AND CARROT JUICES

Matter, Amal A.M.; Samaa M. E. Mahmoud and Rabab H. Salem
Food Science & Technology Dept., Fac. of Home Economics, Al- Azhar Univ., Tanta. Egypt

ABSTRACT

The present study proposed to produce a new pasteurized blended nectars from husk tomato (*Physalis peruviana* L.) and carrot (*Daucus carota*) juice in order to determine whether it could prefer to the consumer.

Results showed that husk tomato juice contained high acidity, 1.43% (pH, 3.79), high content of ascorbic acid (31.0 mg/100 ml) and total sugars (6.41%). Meanwhile, carrot juice contained high carotenoids content (160.0 µg/100 ml). So, attention was paid to make use of the valuable constituents and quality attributes of each variety by prepared blended nectars.

Treatments comprised three pasteurized blended nectars representing (1:1), (1:2), and (2:1) (w:w) from husk tomato and carrot juices respectively. It could be observed that, the addition of carrot juice to husk tomato juice improved taste and over all acceptability meanwhile the addition of husk tomato juice increased ascorbic acid, total acidity and total sugars compared to carrot juice alone.

The storage at ambient temperature (20-25°C) of pasteurized nectars for 3 months, decreases the total sugars, ascorbic acid as well as carotenoids. No microorganisms were detected in all samples during storage period. Highest organoleptic scores were recorded for nectars prepared from husk tomato and carrot juices in ratio (1:2) followed by nectars prepared in ratio (1:1) and (2:1) respectively.

Keywords: *Physalis peruviana*, carrot, natural juices, nectar, pasteurized.

INTRODUCTION

Husk tomato (*Physalis peruviana* L.) fruit belong to solanaceae family. It native to tropical and subtropical American region. Husk tomato fruit are nutritious, containing particularly high levels of niacin, carotenoids and minerals (Elsheikha, 2004). Husk tomato having yellow flowers with purple centers and an inflated calyx enclosed an edible yellow or orange berry used to make jam, sauces and desserts.

Fruit of the herbaceous perennial Chinese lantern *Physalis peruviana*, *P. pubescens*, and *P. edulis*, resembling small cherry, surrounded by dry, bladder like calyx. Also known as golden berry, peruviana cherry and ground tomato. Dwarf cape gooseberry (Morton, 1904).

Husk tomato shows its constituents, possibly polyphenols carotenoids, which demonstrate anti-inflammatory and antioxidant properties. In folk medicine, husk tomato has been used as a medicinal herb for cancer, malaria, asthma, hepatitis, dermatitis and rheumatism (Wu, Sj., et al; 2006; Franco et al., 2007 and Pardo, et al., 2008).

Husk tomato is one of the most promising exotic fruits, which can be used for developing interesting functional products. The pulp was yellowish or orange color with a yield 64%. The pulp and fresh juice had a light sweet and acidic taste (pH 3.7). The titratable acidity was 1.23%, polyphenols, 82 mg/100g and vitamin C, 32.7 mg per 100g fruit. Husk tomato fruit and its juice were rich in carotenoids (19.6 µg/g and 70 µg/ml, respectively). Juice had

high levels (in mg/ml) of minerals such as phosphorous (๐๗๗), potassium(๑๑๖), zink (๕,๕) and boron (๑). Essential amino acids as Leucin, Lysine, Isoleucine, valine and tryptophan were either higher than those recommended by FAO/WHO/UNU (๑๑๗๐).

Carrot (*Daucus carota* L.) is a popular vegetable and is increasing in importance, owing to the fact that its value in the diet is better understood than it was known. It is rich in β -carotene, a precursor of vitamin A, and appreciable quantities of thiamine and riboflavin (Thamposan and Kelly, ๑๑๗๗).

Carrots had the highest content of carotene compared to several raw fruits and vegetables (Simon, ๑๑๖๐). In general, carrot contains ๖,๑ to ๑๐,๗ mg carotenoids per ๑๐๐ g (Heinonen, ๑๑๖๐). It has been reported that β carotene constitutes a large portion (๖๐-๗๐%) of carotenoids in carrots, followed by α -carotene (๑๐-๕๐%), lutein (๑-๐%), and the other minor carotenoids (๐.๑-๑%) (Baloch, *et al.*, ๑๑๗๗; Bushway and Wilson, ๑๑๗๕; Heinonen, ๑๑๖๐; Munsch and Simard, ๑๑๗๕; Seifert and Buttery, ๑๑๗๗).

The carrot is an important root vegetable and is often used for juice production (Yoon, *et al.*, ๒๐๐๐). Carrot juice had a high nutritional value as it is an important source of carotene, and carrot juice is preferably used as a natural source of pro-vitamin A in the carotenoids drinks (Demir, *et al.*, ๒๐๐๕; Yoon *et al.*, ๒๐๐๐). In many countries, a steady increase of carrot juice consumption has been reported (Schieber, *et al.*, ๒๐๐๑). The consumption of carrot juice concentrate (CJC) is increasing very fast in China, which reached ๐,๕๗, ๐,๕๕ and ๐,๐๕ million tons with ๕๐Brix in the year of ๒๐๐๑, ๒๐๐๒ and ๒๐๐๕, respectively (Wang and Chen, ๒๐๐๕; Wang, *et al.*, ๒๐๐๕). It was known that many juices darken during thermal processing and storage (Wang *et al.*, ๒๐๐๖).

According to Simon (๑๑๖๐) vitamin A was the most common dietary deficiency in the world. Vitamin A is derived from carotene, which are abundantly available in vegetables and fruits. So food-based strategies have been regarded as important measures for combating vitamin A deficiency (VAD) in many countries, especially in an agriculture-based country (Udomkesmalee, ๑๑๖๒). Among the carotenoids, β -carotene theoretically possesses ๑๐๐% vitamin A activity while α -carotene possesses between ๐๕% (Bushway and Wilson, ๑๑๗๕) and ๐๐% (Heinonen, ๑๑๖๐). β -carotene provides ๗๐% of vitamin A value measured as retinol equivalents (Chen, *et al.*, ๑๑๖๐; Chou and Breene, ๑๑๗๒). Besides, being a remedy for VAD, the carotenoid pigments also serve as effective antioxidants that quench free radicals provide protection against oxidative damages to cells, multitude of degenerative diseases and also stimulate immune function (Ames, ๑๑๗๕; Anon, ๒๐๐๐a; Craig, ๑๑๖๖; Hudson, ๑๑๗๐).

Carrot juices are preferably used as a natural source of pro-vitamin A in the production of alpha-tocopherol beta-carotene drinks (ATBC-drinks) leading to superior physical and chemical stability (Carle, ๑๑๖๖; Marx, *et al.*, ๒๐๐๐). Carrot juice is one of the most popular vegetable juices (Marx, *et al.*, ๒๐๐๕), and it is preferably used as a natural source of pro-vitamin A in the carotenoid drinks (Yoon, *et al.*, ๒๐๐๐). In many countries, a steady increase of carrot juice consumption has been reported. From ๑๑๖๐ to ๑๑๖๖, German

carrot juice production increased by 69% (Schieber, *et al.*, 2001). The consumption of carrot juice is increasing very fast in China (Wang *et al.*, 2006). Since carrot juice is a low-acid food of approximately pH 6.0, it has a higher risk of bacterial contamination than other acidic foods (Park, *et al.*, 2002). Hence, it requires severe heat treatment (100 to 121°C) for protection from spoilage (Kim and Gerber, 1988; Chen, *et al.*, 1990).

The quality and shelf life of fruit juice depend upon a range of internal parameters related to the product and several external factors, packaging being one of them, (Mannheim *et al.*, 1987).

Additionally, no attention has been paid to the utilization of husk tomato fruit in food industries, and there is little available data in the literature regarding its nutritional properties and industrial utilization. So this work aims to produce new pasteurized nectars prepared from husk tomato and carrot juice with high quality and rich in carotenoids specially β -carotene and providing useful information for storage properties and industrial application of husk tomato. Also the effect of pasteurization and storage at ambient temperature (20-30°C) for 3 months on the chemical composition, microbial evaluation and organoleptic characteristics of blended juice were investigated.

MATERIALS AND METHODS

Materials:

Ripe husk tomato (*Physalis peruviana* L.), fruits were obtained from local market from Ismailia governorate, Egypt during April 2009. Samples were carefully selected according to the degree of ripeness which observed by berries colour (brilliant orange). The berries were peel back the husk and rins it, then sorting, washed, crushed and blended to obtain juice.

Carrot (*Daucus carota*) was obtained from local market from Ismailia. Carrot were washed, trimmed and chopped lightly.

The juice was obtained from the chopped carrot heated in water at 90°C. The carrot was mixed with water in the ratio 1:1 (w/w) to facilitate the extraction of the juice.

The juice of husk tomato fruits and carrot was used to prepare five different blended nectars as the following:

- 1-husk tomato juice alone
- 2-Carrot juice alone
- 3-husk tomato and carrot blend (1:1) (w/w)
- 4-husk tomato and carrot blend (1:2) (w/w)
- 5- husk tomato and carrot blend (2:1) (w/w)

Each of the blended treatment was adjusted to 14% total soluble solids using sucrose solution (20%). All prepared nectars were filled in white glass bottles, closed tightly, pasteurized at 80°C for 10 minutes. Then cooled suddenly in cold water bath. Processed nectars were stored at ambient temperature (20-30°C) for 3 months.

Methods:

Moisture content, total soluble solids, free amino nitrogen, total acidity and pH value were determined according to methods described in the A.O.A.C (2000).

Colour index was determined colorimetrically at 420 nm according to the methods described by Ranganna (1979). Ascorbic acid was determined by 2,6 dichlorophenolindophenol as described by A.O.A.C. (2000). Total carotenoids were determined spectrophotometrically (in diethyl ether) at wavelength 440 nm as mentioned in the methods of Beadle and Zscheile (1952). Total and reducing sugars content were determined colorimetrically according to the methods of Bernfeld, (1900) and Miller, (1909)..

Total plate count, moulds & yeast, coliform group and spore former bacteria were determined according to Oxoid (1992).

Sensory evaluations were carried out by 10 trained panelists to evaluate the quality of the final product according to the method of Larmond, (1970). Ten panelists were asked for their decision concerning the organoleptic attributes using a scale from 1 to 10. The score 6 was considered as the limit of acceptability.

Statistical analysis for data obtained were subjected to analysis of variance according to SPSS, (1997). Significant differences among individual means were analyzed by Duncan's multiple range test (Duncan, 1900).

RESULTS AND DISCUSSION

The chemical composition of fresh juices and prepared nectars from husk tomato and carrot juice were evaluated and presented in Table (1). Results revealed that, husk tomato juice contained 1.42% total acidity (as citric acid), 31.0 mg/ 100ml ascorbic acid and Total sugars 6.41%. These values were higher compared to that in carrot juice. Meanwhile, carotenoids content of carrot juice was 66.0 µg/ 100ml compared with 3843 µg/ 100ml for husk tomato juice. Furthermore, the color index and free amino nitrogen content in the prepared husk tomato nectar was 0.22 and 2.78 mg/ 100ml respectively.

Table (1): Chemical composition of fresh juices and nectar prepared from husk tomato and carrot juice

Treatment	Husk tomato		Carrot		**1:1		**1:2		**2:1	
	Juice	nectar	*Juice	Nectar	Juice	nectar	Juice	nectar	Juice	nectar
Chemical composition										
Total soluble solids (T.S.S.) %	12.2	18.0	4.3	18.0	8.1	18.0	6.9	18.0	9.2	18.0
Total acidity (as citric acid) %	1.42	1.22	0.32	0.30	0.53	0.50	0.330	0.31	0.79	0.74
pH value	3.79	3.81	0.96	6.01	4.06	4.12	4.08	4.60	3.96	4.99
Ascorbic acid (mg/100ml)	31.0	29.8	3.0	2.70	19.00	13.30	16.00	10.12	20.8	19.20
Free amino nitrogen mg/100ml)	2.8	2.78	1.21	1.19	2.02	2.49	2.24	2.20	2.80	2.77
Total sugars %	6.41	12.30	2.01	12.21	4.04	12.30	3.38	12.19	4.86	12.38
Reducing sugars %	3.61	10.32	1.1	10.03	2.00	10.22	1.82	10.18	2.82	10.28
Non reducing sugars %	2.80	2.03	.91	2.18	1.99	2.08	1.06	2.01	2.04	2.10
Carotenoids (µg/100ml)	3843	3340	66.0	612.0	0.72	42.00	0.840	0.800	400.0	410.0
Color index (O.D. at 420 nm)	0.216	0.220	0.270	0.270	0.200	0.202	0.230	0.230	0.200	0.208

* diluted carrot / juice with water at ratio (1/1)

** husk tomato / carrot blend ratio (w/w)

These results are in agreement with those found by (El- Sheikha *et al.*, 2008; Sanad, 1991) and Mattuk *et al.*, 1996).

Results in Table (2) showed that the effect of processing and storage at ambient temperature (20- 25°C) for 3 months on some chemical composition of the pasteurized nectars prepared from husk tomato and carrot juices. It could be included that, there was decreased of pH values during storage period. The pH values decreased conversely to total titratable acidity. Ascorbic acid content sharply decreased, it may be due to the effect of heat treatment of pasteurization, and oxidation these results are in agreement with those found by Gerschenson and Rojas (1997). The free amino nitrogen content also decreased slightly.

Table (2): Effect of storage at ambient temperature (20- 25°C) for 3 months on the total acidity, pH, ascorbic acid and free amino nitrogen of husk tomato and carrot pasteurized nectar

Treatment	Husk tomato	*Carrot	**1:1	**1:2	**2:1
Period/ month					
Total acidity					
0	1,20	0,03	0,00	0,30	0,74
1	1,28	0,10	0,64	0,44	0,92
2	1,42	0,28	0,82	0,71	1,00
3	1,68	0,46	1,10	0,90	1,64
pH value					
0	3,81	6,01	4,11	4,66	4,99
1	3,80	0,94	4,06	4,64	4,92
2	3,70	0,89	4,01	4,60	4,89
3	3,70	0,82	3,96	4,08	4,82
Ascorbic acid					
0	18,60	1,70	1,00	8,00	11,10
1	17,30	1,31	9,70	8,10	10,70
2	16,40	1,11	8,90	7,44	10,30
3	10,02	0,92	8,31	6,30	9,80
Free amino nitrogen					
0	0,60	2,80	3,90	3,20	4,10
1	0,08	2,02	3,08	2,80	3,80
2	0,20	2,24	2,80	2,40	3,20
3	4,80	2,00	2,40	2,20	2,80

* diluted carrot / juice with water at ratio (1/1)

** husk tomato / carrot blend ratio (w/w)

Data given in Table 2 show the effect of processing and storage at ambient temperature for 3 months on total sugars, reducing sugars, carotenoids and color index of husk tomato and carrot pasteurized nectars. Total and reducing sugars were decreased. These decreased in sugars and free amino nitrogen may be due to the Millard reaction, this also in agreement with those reported by Tressler and Joslyn (1961).

Carotenoids constitute is the major coloring substances in juices. The results were decreased in carotenoids after processing of husk tomato and carrot nectars, the percentage of carotenoids decreases were 18,81% and 7,40% respectively. This decrement reached to 20,00% and 10,00% of the initial

amount of carotenoids after 3 months of storage. It could be explained by oxidation reactions that developed during storage and also by Millard reactions this results in agreement with those reported by Nicoli *et al.*, (1999) Singleton *et al.*, (1961) reported that the total carotenoids content is unchanged and these changes which occur during processing could be attributed to the isomerization, they added also that the decrease could be occurred due to the leaching of pigments in the blanching solution

Table (3): Effect of storage at ambient temperature (20- 30°C) for 3 months on the total sugars, reducing sugars, carotenoids and color index of husk tomato and carrot pasteurized nectar

Treatment Period/ month	Husk tomato	*Carrot	**1:1	**1:2	**2:1
Total sugars					
0	12,30	12,21	12,30	12,19	12,38
1	12,07	12,12	12,08	12,00	12,10
2	11,40	11,87	11,77	11,02	11,74
3	11,00	11,78	11,34	11,34	11,22
Reducing sugars					
0	10,32	10,03	10,22	10,18	10,28
1	10,02	9,94	10,00	9,92	10,00
2	9,88	9,00	9,78	9,76	9,70
3	9,74	9,34	9,00	9,42	9,30
Carotenoids					
0	3120	7108	4180	0000	4122
1	3030	7099	4090	0049	4000
2	2970	7011	4020	4979	3970
3	2880	0941	3900	3899	3890
Color index					
0	0,220	0,271	0,200	0,237	0,208
1	0,290	0,340	0,281	0,277	0,280
2	0,330	0,470	0,310	0,290	0,330
3	0,040	0,711	0,480	0,341	0,477

* diluted carrot / juice with water at ratio

** husk tomato / carrot blend ratio (w/w)

The color index of all samples of the pasteurized nectars was slightly increased during storage period. This increasing may be attributed to browning reactions. These findings are in accordance with those found by (Speers *et al.*, 1987; Aguilera *et al.*, 1987; Avil and Silva, 1990). El-Sheikha (2004). Reported that the increasing of the juice optical density is usually an index for undesirable changes in its quality and safety. The darkening may be due to non- enzymatic browning reactions, as cited by Buglione and Lozano (2002).

Microbial counts (total bacterial count, moulds & yeasts and spore former bacterial count) and coliform group of pasteurized blended nectars were studied and the obtained results were given in Table (4). The total bacterial counts were 47×10^{-1} , 8×10^{-1} , 20×10^{-1} , 19×10^{-1} and 38×10^{-1} CFU/g in husk tomato, carrot, 1:1, 1:2 and 2:1 nectars respectively. While these values were 108×10^{-1} , 3×10^{-1} , 70×10^{-1} , 09×10^{-1} and 93×10^{-1} for moulds &

yeasts. All samples were free from spore former bacterial count and coliform group. These results revealed the effect of following sanitation during processing and may also due to the anti-microbial effect of husk tomato juice against some serious pathogenic microbes. The anti-microbial activity of methanolic extracts of juice was significant against *E. coli* O157:H7 and *Bacillus subtilis*, complete against *Fusarium solani* and had no effect against *Candida albicans* (El-Sheikha et al., 2008).

Table (4): Effect of storage at ambient temperature (25- 30°C) for 3 months on total bacterial count, moulds & yeasts and spore former bacterial count of pasteurized nectar

Treatment Period/ month	Husk tomato	*Carrot	**1:1	**1:2	***2:1
Total bacterial count					
***	47x10 ⁶	8x10 ⁶	20x10 ⁶	19x10 ⁶	38x10 ⁶
1	Not detected	Not detected	Not detected	Not detected	Not detected
2	Not detected	Not detected	Not detected	Not detected	Not detected
3	Not detected	Not detected	Not detected	Not detected	Not detected
Moulds & Yeasts					
***	108x10 ³	3x10 ³	60x10 ³	09x10 ³	93x10 ³
1	Not detected	Not detected	Not detected	Not detected	Not detected
2	Not detected	Not detected	Not detected	Not detected	Not detected
3	Not detected	Not detected	Not detected	Not detected	Not detected
Spore former bacterial count					
***	Not detected	Not detected	Not detected	Not detected	Not detected
1	Not detected	Not detected	Not detected	Not detected	Not detected
2	Not detected	Not detected	Not detected	Not detected	Not detected
3	Not detected	Not detected	Not detected	Not detected	Not detected
Coli form group					
***	Not detected	Not detected	Not detected	Not detected	Not detected
1	Not detected	Not detected	Not detected	Not detected	Not detected
2	Not detected	Not detected	Not detected	Not detected	Not detected
3	Not detected	Not detected	Not detected	Not detected	Not detected

* diluted carrot / juice with water at ratio (1/1)

** husk tomato / carrot blend ratio (w/w)

*** freshly prepared nectars before pasteurising

Data in Table (5) show the organoleptic evaluation of pasteurized nectar prepared from husk tomato and carrot juice. Results revealed that, there are no significant differences in taste between the samples (husk tomato, 1:1 and 2:1) respectively and the best treatment was, (1:2). Data also show that the treatment, (1:2) had a higher score for odor characteristic and there are no significant differences between the samples, (husk tomato, carrot, 1:1 and 2:1). The significantly, (≤ 0.05) lowest score of color was given for samples, (husk tomato and 2:1) respectively and the best color quality was samples, 1:2, which also showed the higher overall acceptability significantly.

Finally, it could be concluded that, it is possible to produce pasteurized nectar from husk tomato and carrot juice at the ratio (1:2) (w/w). These blended have high nutritional value and preferred by consumer

In a row, means followed by the same superscript letters are not significantly different at 5% level

Table (°): Organoleptic evaluation of pasteurized nectar prepared from husk tomato and carrot juice

Attribute	Treatment				
	Husk tomato	carrot	1:1	1:2	2:1
Taste	7.0 ^{bc}	5.0 ^a	7.0 ^{bc}	7.5 ^c	6.0 ^b
Odor	6.0 ^b	5.6 ^b	6.3 ^b	7.9 ^a	6.0 ^b
Color	7.0 ^a	8.0 ^b	7.7 ^b	9.0 ^c	7.0 ^a
Overall acceptability	7.0 ^b	6.2 ^a	7.1 ^b	8.9 ^c	5.9 ^a

REFERENCES

- Aguilar, J.M.; Oppermann, K. and Sanchez, F. (1987). Kinetics of browning of Sultana Grapes. *J. Food Sci.*, 52 (4): 990-994.
- Ames, B. N. (1983). Dietary carcinogens and anticarcinogens (oxygen radicals and degenerative diseases). *Science*, 221, 1206-1274.
- Anonymous. (2006). Nutritional and health properties of the carrot. <http://website.lineone.net/stolarczyk/nutrition.html>. As retrieved on 0/1/2006.
- A. O. A. C. (2000). "Official Method of Analysis" 17th Ed. Association of Official Analytical Chemists, Published by Association of Official Analysis Chemists, Inc., USA.
- Avila, I.M.L.B. and Silva, C.L.M. (1999). Modeling Kinetics of Thermal Degradation of Colour in Peach Puree *J. Food Eng.*, 29: 161-166
- Baloch, A. K., Buckle, K. A., & Edwards, R. A. (1977). Separation of carrot carotenoids on hyflo super-cel-magnesium oxide-calcium sulfate thin layers. *Journal of Chromatography*, 129, 149-150.
- Beadle, M. and Zscheile, A. (1942). *J. Biol. chem.*, 144:21. C. F. Hawk P.B.; Oser, B. L. And Summer son, W. H. 1904, practical physiological chemistry, 13th Ed. MCGRAW- Hill Book Co., Inc. New york. Toronto And London P. 1121
- Bernfeld, F. (1900). *Methods In Enzymology*. Acad. Press, Inst., New York. In. S. P. Colowick And N. O. Kaplan (Eds.) pp 149-154.
- Buglion, M. and J. Lozana, (2002). Nonenzymatic browning and hemical changes during grape juice storage, *J. Food Sci.*, 67: 538- 1043
- Bushway, R. J., & Wilson, A. M. (1982). Determination of α and β carotene in fruit and vegetables by HPLC. *Canadian Institute of Food Science and Technology Journal*, 10, 160-169.
- Carle, R. (1999). Physical and chemical stability of ATBC-drinks. *Fruit rocessing*, 9, 342-349.
- Chen, B. H., Peng, H. Y., & Chen, H. E. (1990). Changes of carotenoids, color, and vitamin A contents during processing of carrot juice. *Journal of Agriculture and Food Chemistry*, 38, 1912-1918.
- Chou, H. E., & Breene, W. M. (1972). Oxidative decolouration of β -carotene in low moisture model systems. *Journal of Food Science*, 37, 66-68.
- Craig, W. J. (1999). Health promoting properties of common herbs. *The American Journal of Clinical Nutrition*, 70, 491S-499S
- Demir, N., Acar, J., & Bahceci, K. S. (2004). Effect of storage on quality of carrot juices produced with lacto fermentation and acidification. *European Food Research and Technology*, 218(5), 460-468.

- Duncan, D. (1900). Multiple range and multiple F-test. Biometric, 11:1- 42. Edition, ASSOC , Office, Anal. Chem., Arlington
- Elsheikha, A. (2004). Technological, chemical and microbiological studies on some packed food M. Sc. Faculty of Agriculture. Minofiya University, Egypt.
- Elsheikha, A. F, Zaki M. S, Bakr A. A, El- Habashy M. M. And Montet (2008). Physico- chemical properties and biochemical composition of physalis (*Physalis pubescens* L.) fruits. Global Science Book L td., U K, Food. 2 (2), 124-130.
- FAO, (1980). Food And Agriculture Organization of United Nations , Amino acids content of food and biological data on proteins. FAO nutrition studies No. 28.
- Franco, L. A., Matiz G. E., Calle J., Pinzon R. and Ospina F.(2007). Anti-inflammatory activity of extracts and fractions obtained from *Physalis peruviana* L. calyces Biomedica 11:110-0.http://www. ncbi. nlm. nih. gov/ pubmed/ 170467228
- Gerscherson, L. N. and A. M. rojas (1997). Influence of System composition on ascorbic acid destruction at processing temperatures. J. Sci. Food Agric., 74:369- 378
- Heinonen, M. I. (1990). Carotenoid and provitamin A activity of carrot (*Daucus carota* L.) cultivars. Journal of Agricultural and Food Chemistry, 38, 609-612
- Hudson, B. J. F. (1990). Food antioxidants (pp. 99). England: Applied Science Publishers
- ICMCF (1978). International Commission on Microbiological Specification for Foods Microorganisms in Foods. Their Significance and Methods of Enumeration 2nd Ed., Univ. of Toronto Press, Toronto ,Canada.
- Kim, H. Y., & Gerber, L. E. (1988). Influence of processing on quality of carrot juice. Korean Journal of Food Science and Technology, 20, 683-690.
- Larmond, S. (1970). Method for sensory evaluation of food publication No. 1284, Cana. Dept. of Agric., Ottawa.
- Mainheim, C. H., Miltz J. and Letzter(1987). Interaction between polyethylene laminated cartons and aseptically packed citrus juices J. Food Sci., 92(3) :737-740.
- Marx, M., Schieber, A., & Carle, R. (2000). Quantitative determination of carotene stereoisomers in carrot juices and vitamine supplemented (ATBC) drinks. Food Chemistry, 70, 403-408.
- Marx, M., Stuparic, M., Schieber, A., & Carle, R. (2003). Effects of thermal processing ontrans-cis-isomerization of β -carotene in carrot juices and carotene-containing preparations. Food Chemistry, 83, 609-617.
- Matuk, H. I., A. M. Ali and K. Sh. Ibrahim (1996). Chemical and technological studies on production of carrot nectar. Egypt, J. Appl. Sci., 11 (9)
- Miller, G. L. (1909). Use of Dinitrosalysilic Acid Reagent for Determination of Reducing Sugars. Anal. Chem., 31,426-428.
- Morton, J. F. and Russell, O. S.(1904). The cape gooseberry and the Mexican husk tomato. Florida State Horticultural Society 17:261- 266.
- Munsch, M. H., & Simard, R. E. (1983). Relationships in colour and carotene content of carrot juices. Canadian Institute of Food Science and Technology Journal, 16, 173-178.

- Nicoli, M. C., Anese M. And M. Parpinel (1999). Influence of processing on the antioxidant properties of fruit and vegetables. Trends Food Sci. Technol., 10:94-100.
- Oxoid, M. (1992). The oxoid manual of culture media and other laboratory services. 5th Edition Basingstock : oxoid Hd
- Pardo, J. M., Fontanilla M. R., Ospina L. F. and Espinosa L.(2008). Determining the pharmacological activity of *Physalis peruviana* fruit juice on rabbit eyes and fibroblast primary cultures Invest Ophthalmol Vis Sci., 49:3074-3079
- Park, S. J., Lee, J. I., & Park, J. (2002). Effects of a combined process of high-pressure carbon dioxide and high hydrostatic pressure on the quality of carrot juice. Journal of Food Science, 67:1827-1834.
- Ranganna, S. (1999). Manual of analysis of fruits and vegetables products. Tata Mc Graw- Hill. Publishing Company Ltd., New Delhi, pp. 84-87
- Robards K., Prenzler P.D., Glover W., Tucker G. and P. Swatsitang (1999). Phenolic compounds and their role in oxidative processes in fruit. food chem., 66: 401- 436.
- Sannad, A. F.(1991). Technological studies on some sugar product Standrized apricot and carrot juices M. Sc. Thesis, Fac. Agric.; Zagazig Univ.
- Schieber, A., Stintzing, F. C., & Carle, R. (2001). By-products of plant food processing as a source of functional compounds-recent development. Trends in Food Science and Technology, 12, 401-413.
- Seifert, R. M., & Buttery, R. G. (1978). Characterization of some previously unidentified sesquiterpenes in carrot roots. Journal of Agricultural and Food Chemistry, 26, 161-163
- Simon, P. W. (1990). Carrots and other horticultural crops as a source of provitamin A carotenes. HortScience, 20, 1490-1499.
- Speers, R. A., M. A. Tung and R. L. Jackman (1987). Production of Color Determination in Strawberry Juice. Candian Institute of Food Sci. And Techn. J., 20 (1), 10- 18
- SPSS, (1997). Spss users Gide statistics version 8 copy right Spss Inc., USA, Washengton, D. C. USA.
- Tressler, D. K. and M. A. Joslyn (1961). Fruit And Vegetable Juice Processing Technology. The Avi. Pub, Co. Inc., Westport, Connecticut
- Thompson, H. C. and W. C. Kelly (1988). Vegetable Crops. Tata Mc Graw- Hill Pub, Co., Ltd, New Delhi
- Udomkesmalee, E. (1992). Overview of vitamin A: Global situation and the experience. In P. Winichagoon, Y. Kachondham, G. Attig, & K. Tontisirin (Eds.), Integrating food and nutrition into development: Thailand's experiences and future visions (pp. 147-152). Nakorn Pathom, Thailand: Inst. of Nutrition, Mahidol University
- Wang, H. Y., Hu, X. S., Chen, F., Wu, J. H., Zhang, Z. H., and Liao, X. J. (2006). Kinetic analysis of non-enzymatic browning in carrot juice concentrate during storage. European Food Research and Technology, 223, 282-289.
- Wang, X. F., and Chen, Z. (2002). Yearbook of China agricultural products processing industries (p. 474). China Agriculture Press.
- Wang, X. F., and Chen, Z. (2004). Yearbook of China agricultural products processing industries (p. 516). China Agriculture Press.

- Wang, H. Y., Hu, X. S., Chen, F., Wu, J. H., Zhang, Z. H., and Liao, X. J. (۲۰۰۶). Kinetic analysis of non-enzymatic browning in carrot juice concentrate during storage. *European Food Research and Technology*, in press, doi:10.1007/S00217-005-0202-2
- Wu, S. J., Tsal J. Y., Chang S. P., Lin D. L., Wang S. S. and Huang S. N. (۲۰۰۶). Super critical carbon dioxide extract exhibits enhanced antioxidant and anti-inflammatory activities of *Physalis peruviana*. *J. Ethnopharmacol* ۱۰۸ (۳): ۴۰۷- ۱۳.
- Yoon, K. Y., Cha, M., Shin, S. R., and Kim, K. S. (۲۰۰۵). Enzymatic production of a soluble fiber hydrolyzate from carrot pomace and its sugar composition. *Food Chemistry*, ۹۲, ۱۵۱-۱۵۷.

المشروب المبستر المحضر من عصائر الحرنكش والجزر

امل عبد الله محمد مطر، سماء محمود السيد محمود و رباب حسن سالم
قسم علوم وتكنولوجيا الاغذية - كلية الاقتصاد المنزلى - جامعة الأزهر - طنطا

يهدف هذا البحث الى انتاج مشروبات جديدة مبسترة محضرة من مخاليط عصائر الحرنكش والجزر لتحديد افضلها لدى المستهلك وقد اوضحت نتائج هذا البحث ارتفاع محتوى عصير الحرنكش من كل من الحموضة الكلية والسكريات الكلية مقارنة بلب الجزر بينما يحتوى عصير الجزر على نسبة عالية من الكاروتينات مما دفعنا للاستفادة من مكونات وعناصر الجودة لكل منهما وانتاج مخاليط منهما بالنسب التالية: (۱:۱) و (۲ : ۱) و (۲ : ۱) (و/و) من عصير الحرنكش والجزر على التوالى وكذلك مشروبات مبسترة من كل منهما على حدة وقد اظهرت نتائج هذا البحث ان اضافة عصير الجزر لعصير الحرنكش ادى لتحسين كل من الطعم والقابلية العامة وكذلك وجد ان اضافة عصير الحرنكش ادى لزيادة محتوى فيتامين ج والحموضة الكلية وكذلك السكريات الكلية مقارنة بعصير الجزر بمفرده. وقد ادى التخزين على درجة حرارة الغرفة (۲۵ - ۳۰ م °) لمدة ۳ شهور لانخفاض المحتوى من السكريات الكلية وفيتامين ج وكذلك الكاروتينات لكل العينات. كما اعطيت اعلى درجات للتقييم الحسى للمشروب المحضر من خلط عصير الحرنكش والجزر بنسبة ۲:۱ واتبعه كل من ۱:۱ ثم ۱:۲ (و/و) على التوالى .

قام بتحكيم البحث

كلية الزراعة - جامعة المنصورة
كلية الاقتصاد المنزلى - جامعة الأزهر

أ.د / منى محمود خليل
أ.د / سعاد محمد الدسوقي بدوى