

EFFECTS OF HIGH FAT DIET ON SOME PHYSIOLOGICAL PARAMETERS IN BROILER CHICKEN UNDER HIGH ENVIRONMENTAL TEMPERATURE

Wehailsh, F. E.*; Abou-Ismael, U.A.** and Ageez, A. M.

*: Department of Physiology, Faculty of Veterinary Medicine, Mansoura University, Egypt.

** : Department of Animal Husbandry, Faculty of Veterinary Medicine, Mansoura University, Egypt.

ABSTRACT

The objectives of this study were to investigate the effects of increasing the level of soybean oil in the diet of broiler chicken on some physiological parameters including growth performance, some blood parameters, some blood metabolites and hormonal changes during the exposure to stress of high environmental temperature. A total of 60 one-day-old chicks were divided into three groups with each group containing 20 chicks of both sexes. One group served as a control group whereas the other two groups served as experimental groups. At day 21 of age, the control group; Normal Temperature Low Fat (NTLF) was transferred to a separate room and remained under ideal temperature (25°C) and the same fat level (3%) in the diet. In the other two experimental groups in which temperature was not adjusted and depended on the outdoor climatic condition of the summer season, one group remained under the same fat level (3%) (High Temperature Low Fat group (HTLF)) and fat level was increased in the diet of the other group (6%) (High Temperature High Fat group (HTHF)). No significant differences between the groups regarding body weight, body weight gain, feed conversion ratio and feed intake were found. A significant decrease in the total red blood cells count (RBCs) was observed in the HTLF group at 42 day of age and in the total leucocytic count (WBCs) at both 35 and 42 days of age. Heterophile/Lymphocyte ratio (H/L) was significantly higher in HTLF and HTHF group at 28 days of age. Serum total cholesterol (TCOL), triacylglycerol (TG), total lipids and high density lipoprotein cholesterol (HDL-c) were significantly higher in HTHF and HTLF group while there was a significant decrease in low density lipoprotein cholesterol (LDL-c) in these groups. Serum total protein was significantly decreased in HTHF and HTLF group. Serum triiodothyronin (T3) was significantly decreased in HTHF and HTLF group at 28 days of age. These results suggest that the addition of soybean oil to broiler diets at a level of 3 and 6% during the period of heat stress can improve growth performance parameters without affecting serum TCOL, TG, total lipids and T3. Increase serum HDL-c and reduce LDL-c.

INTRODUCTION

Broiler industry is increasing dramatically

throughout the developing countries. It has been developed very fast in the last two dec-

ades to become one of the most important sectors in animal production industry. Fat is frequently included in poultry diets to increase the energy density (Pinchasov and Nir, 1992; Sanz et al., 1999). As no special problems are associated with feeding fats to poultry, the use of different vegetable oils allows the enrichment of poultry meat with poly-unsaturated fatty acids (Lopez-Ferrer et al., 1999). Also and Balnave, (1985) investigated the nutritional significance of different dietary fat level (sunflower oil, maize oil, poultry fat and animal tallow) for the growing broilers and concluded that increasing the dietary supplementation of fats and oils from 3 to 9% improved the growth rate of broilers. Also, optimum growth responses were obtained with vegetable oils. It was found that diets contained saturated fat had poorer feed conversion ratio than those contained polyunsaturated fats (Pinchasov and Nir, 1992; Zollitsch et al., 1997). Nitsan et al., (1997) observed that the addition of 3% soybean oil in the diet of broilers improved the live weight gain significantly than the diet containing 0% soybean oil. They also observed higher (but not significant) live weight gain of 3.4% in group of broiler fed on 6% soybean oil than 3% soybean oil added diet during heat exposure.

Tony (1997) investigated the effect of dietary fat sources (rich in essential fatty acids) on some blood constituents of broiler chickens. The effects of feeding high energy diets containing supplemented fat on some blood constituents in broilers were investigated by Wafaa and El-Sadawy, (1997). No significant changes in serum total protein, albumin and globulin among different treatments were re-

ported while serum total lipids and cholesterol significantly decreased in the broilers fed low fat supplemented diets. High content of HDL-c was found in broilers fed 6% corn oil and soybean oil respectively. Cholesterol level was also lower in groups fed 6% sunflower oil and soybean oil respectively (Özdoğan and Aksit, 2003). Dietary supplementation with 3 or 6% soybean oil did not significantly affect serum total lipid in the experimental broiler chicks (Mansour, 2006). In addition, El-Katcha and Walaa, (1996) reported that blood metabolites in broilers did not significantly affected by dietary fat level. Bartha (1993) showed that adding fat to isocaloric diet decreased T3 and increased T4 levels in the serum of broiler chicken. Kahl et al., (1998) and Rosenbrough et al., (1999) reported no differences in plasma thyroid hormone in broiler chickens fed diets with different dietary fat levels. These findings indicate that there is a close relationship between thyroid hormone metabolism and fat metabolism but the effect of dietary fat type needs more investigation.

In Egypt high ambient temperature during the summer generates a status of stress and evokes a combination of behavioural, biochemical, immunological and physiological changes (Abd-Elsamee, 2005 and Abdel-Fattah, 2006). The expression of heat stress in poultry production can be described as "acute" or "chronic". Acute heat stress refers to short and sudden periods of extremely high temperature whereas chronic heat stress refers to extended periods of elevated temperature (Emery, 2004). High environmental temperature have deleterious effects such as reducing rate of growth, feed intake, live

weight gain, feeding efficiency and digestibility of nutrients (Donkoh, 1989 and Mills et al, 1999). Chronic heat stress has detrimental effects on the performance of broiler birds reared in open sided poultry houses, principally through reducing feed intake and growth rate, negatively affect feed efficiency as well as health (Carmen et al., 1991 and Har et al., 2000).

Heat stress may shorten the RBC's life span, resulting in a decrease total numbers of RBCs (Sturkie, 1986; Sahota et al., 1994 and Rautela et al., 1994). Mashaly et al., (2004) found that birds exposed to chronic heat stress had a lower WBCs count compared to those exposed to chronic cycle or controlled temperature. They also reported that the total WBCs counts were significantly inhibited in hens (31 weeks old) exposed to heat stress (constant heat stress and humidity for 5 weeks). The heterophile/ lymphocyte ratio has been shown to be a highly heritable (Al-Murrani et al, 1997 and Zulkifli et al., 2003) and a reliable index for determining stress in poultry (Gross and siegel, 1983). Borgea et al., (2004) showed that heat stress altered the proportion of heterophile (increased) to lymphocyte (decreased) in blood. Puvadolpirod and Tharton, (2000) reported that stress increased serum cholesterol levels in chickens. Sosnowka-Czajka et al., (2005) exposed chicks to elevated ambient temperature from the 21st day of age to the 28th day of age and found that the highest level of cholesterol was found in these chickens on last day of rearing (day 42). They also reported a high level of triacylglycerol in these birds.

Gursu et al., (2004) investigated the ef-

fects of high environmental temperature, vitamin C, and folic acid supplementation on serum metabolites in Japanese quails and found an increase in the level of cholesterol, high density lipoprotein cholesters (HDL-c) and triglyceride in the blood of the heat stressed birds. Sahin et al., (2001) recorded a reduction in the concentration of total protein, albumin and globulin in broilers as a result of the exposure to chronic heat stress from 21 to 42 days of age. It is generally accepted that neuron- endocrine factors act as an interface between input factors such as genotype, sex, environmental conditions and nutrition (Buyse et al., 2001). Heat stress caused a series of physiological and metabolic changes in broiler chicken (Deyhim and Teeter, 1991) such as a reduced metabolic status which was reported to occur due to decreased levels of plasma triiodothyronine (T3). Similarly, Saurate et al., (1987) indicated that the plasma T3 concentration decreased, but the T4 concentration increased during exposure to high temperature. On contrary to that, Geeraert et al., (1996) reported a significant reduction in plasma T3 concentration whereas T4 concentration did not decrease as much or even remained unchanged under chronic heat exposure. Blood corticosterone concentration has been widely used as a measure of environmental stress in broilers (Zulkifli et al., 2003 and McFarlane and Curtis, 1989). Lin et al., (2006) reported that corticosterone concentration in broilers that had been exposed to acute heat stress (32°C and 40% relative humidity) for 6 hours varied from 3.17 to 4.28 ng/ml on day 42. This experiment was therefore carried out to investigate the effects of (increasing the level of soybean oil in the diet of broiler chicken on some physiological

parameters including growth performance, some blood parameters, some blood metabolites and hormonal changes during the exposure to stress of high environmental temperature.

MATERIALS AND METHODS

A total number of 60 one-day-old chicks (Cobb breed) were divided into three groups with each group containing 20 chicks of both sexes. During the first 3 days of age, the brooding temperature was maintained at 35 to 32°C with a constant lighting (24 hours / day). The room temperature was decreased by 1°C every 2 days reaching 22-25°C by the beginning of the third week of age. At day 21 of age, the control group was transferred to a separate room and remained under ideal temperature (25°C) and the same fat level in the diet till the end of the experiment. In the other two experimental groups the temperature was not adjusted and depended on the outdoor climatic condition of the season, one group remained under the same fat level (Low fat group) whereas the fat level was increased in the diet of the other group (High fat group) (table 1). Corn-Soybean based broiler diet (starter-grower and finisher diet) was used with two levels of soybean oil (3% and 6%) in isonitrogenous and isocaloric diet. The diet was formulated according to NRC (1994) recommendation (Table 2).

Body weight, body weight gain, feed efficiency and feed intake were calculated for each group at 21, 28, 35 and 42 days of age. Blood samples were collected from the birds at these ages. Five birds from each group were bled from the wing vein. Each collection was divided into two samples. The first one

was taken with sodium citrate 3.8% freshly prepared as anticoagulant (Shum and Grimmer, 1972) and was used immediately for the determination of the total erythrocytic count, total leucocytic count (Feldman et al., 2000) and differential leucocytic count (Gross and Siegel, 1983). The second sample was collected without anticoagulant and was left to clot at room temperature. It was then centrifuged at 3000 rpm for 15 minutes to separate clear serum sample which was stored frozen at -20°C for later determination of serum total cholesterol (Allain et al., 1974), triacylglycerol (Passati and Prencipe, 1982), HDL-c (Lopez-Virella et al., 1977), LDL-c (Fradwald et al., 1972). Total lipids (Zollner and Kirsch, 1982), serum total protein (Gornal et al., 1949), albumin (Dumas et al., 1971) using commercial available kits and T3 (Klee, 1996), T4 (Albertini and Ekris, 1982) and corticosterone hormone (Donia, 1967) using ELISA kit. Statistical analysis was done using SPSS (1994). Mean and standard errors were calculated for the obtained data and the level of significance for all means was determined using general linear model (GLM) of Univariate analysis (full factorial design).

RESULTS AND DISCUSSION

Growth performance parameters: There were no significant differences between the two experimental groups (High temperature-low fat (HTLF) and high temperature-high fat (HTHF) and between the two experimental groups and the control one (normal temperature-low fat (NTLF)) during the last three weeks at which the experimental conditions were applied in respect to body weight (table 3), body weight gain (table 4), feed conversion

ratio (table 5) and amount of feed consumed (table 6). These results agree with those of **Whitehead and Keller, (2003)** and **Ghazalah et al., (2007)** who reported that nutrient modification affects productivity, health and physiological processes occurring in the body of broilers reared in elevated ambient temperature. On the other hand, the performance of broiler chicks was improved with using fat in broiler chick diet. The improvement seen in the birds of the two experimental groups that were exposed to the chronic heat stress for the last three weeks of age can be explained by **Ferket (1995)** who found that fat could improve energetic efficiency of a diet fed during hot weather in 3 ways. First, dietary fat gives 2.25 times more energy per unit of weight than carbohydrate or protein so, it can be used to increase the flexibility of feed formulation by allowing more inclusion possibilities for another crucial nutrients. Second, digestion and metabolism of dietary fat generates less body heat per gram when absorbed and used for growth than dietary carbohydrate and protein. Third, the rate of food passage is reduced by dietary fat which may increase the digestibility of other ingredients (**Mateos et al., 1982**). Higher, but not significant, live weight gain was seen in heat exposed chicken fed diet contain 6% soy bean oil than heat exposed chickens fed diet contain 3% soybean oil and this finding agrees with the result obtained by **Nitsan et al., (1997)**.

Blood parameters: The results shown in table (7) revealed a significant decrease in RBCs count in the HTLF and HTHF group at the 42nd day of age. Reduced numbers of red blood cells in heat stressed birds was ob-

served by **Sahota et al., (1994)** and **Rautela et al., (1994)**. A higher RBCs count was seen in HTHF than HTLF group due to increasing the fat level from 3 to 6% which is supported by the finding of **Mateos et al., (1982)** who found that the digestibility of other nutrient including protein might increase which it needed in the RBCs building. The reduction in the RBCs count might be due to demodulation due to increased water consumption during heat stress period. Furthermore, **Sturkie, (1986)** stated that the numbers of RBCs are dependent on both the numbers of newly produced RBCs and the average life span of the old cells, hence heat stress may shorten the RBCs life span, resulting in a reduction in the total numbers of RBCs. The significant reduction in the WBCs count that is seen in the current experiment in table (8) is in hand with the finding of **Mashaly et al., (2004)** who reported that birds exposed to chronic heat stress had a lower WBCs count compared with birds exposed to chronic or controlled temperature. The results in table (8) revealed a significant increase in heterophile (H %), H/L ratio and a decrease in lymphocyte (L %) in both the HTLF and the HTHF groups during the period of the last 3 weeks of age. Broiler exposed to heat stress in summer showed an increase in heterophiles and a decrease in lymphocytes, which led to an increase in the H:L ratio. These results are in an agreement with those of **Borges et al., (2004)** and **Turkyilmaz,, (2008)** who reported that increased circulating (H/L) ratio is one of the most accepted indicators of the stress condition in birds as heat stress altered (increased) the proportion of heterophile to lymphocyte (decreased) in the blood and there was also a significant

increase in H/L ratio in birds after the application of heat stress.

Serum metabolites: Results presented in table (8) showed significant increases in the level of serum total cholesterol and triacylglycerol in the HTLF and HTHF than NTLF group. These results are in harmony with those of **Siegel, (1968)** and **Özbey et al., (2004)**. High temperature increased levels of serum total cholesterol and triglyceride. It could be said that as a consequence of increased serum total cholesterol and triglyceride levels, levels of total lipids were subsequently increased in the two heat stressed groups. From the results recorded in table (8) it can be observed that increasing fat level in the diet from 3 to 6% in the HTHF group did not increase cholesterol and triglyceride levels in this group compared to the HTLF group. The significant increases in the levels of HDL-c in heat stressed groups observed in this study are in an agreement with the findings of **Gursu et al., (2004)**. Increasing fat level from 3 to 6% in the HTHF group increased the level of HDL-C in the HTHF group compared to the HTLF group at the 28th and the 35th days of age. It also decreased levels of LDL-e in the HTHF in comparison to the LTLF group at the 28th, 35th and the 42nd days of age. This is in an agreement with the results of **Kinsella et al., (1990)** who stated that dietary PUFA of vegetable oils containing mostly linoleic acid are effective in counter-acting the effects of dietary saturated FAs. Thus n-3 PUFAs may reduce plasma lipids and alter the cell and tissue PUFAs. In respect of serum total protein the results shown in table (9) revealed that serum total protein levels were reduced in the two heat stressed

groups (HTLF and HTHF) compared to the control group (NTLF). This finding is in an agreement with that of **Makind and Fatunmbi (1985)**; **Faitas et al., (1987)**; **Kutlu and Forber, (1993)** and **Berrong and Washburn, (1998)** who reported that concentrations of blood protein were decreased significantly when birds were exposed to heat stress. These results may be due to the reduction in the amount of protein consumed and the reduced protein digestibility with the exposure of broiler chickens to high temperatures (**Bonnet et al., 1997**; **Ozbey and Ozcelik, 2004**; **Abu-Dlayeh, 2006a,b**). The non significant decreases in the levels of serum albumin in HTLF and HTHF group and the non significant changes in the levels of globulin in comparison with the control group as shown in table (9) disagree with the results of **Faisal et al., (2008)** and **Sahin et al., (2001)**. This might be due to differences in the condition of heat stress application.

Serum hormonal parameters: Data illustrated in table (10) revealed decreased levels of T3 in both HTLF and HTHF group at the 28th days of age and levels are non significantly changed than the control group at the 35th and the 42nd days of age. These results are supported by those reported by **Geraert et al., (1996)** who found a significant reduction in plasma T3 concentration under chronic heat stress. It is well known that thyroid hormones play a crucial role in thermoregulation in avian species and that plasma T3 levels are positively correlated with heat production (**Decuyper and Kuhn, 1984, 1988**). T4 concentrations remained significantly unchanged throughout the 3 weeks of applying the heat stress condition. These findings are

In harmony with those of **Geraert et al., (1996)**. The finding that T3 levels returned to a level similar to that of the control group at the 35th and the 42nd days of age might be due to various factors impact on T3 and T4 levels including species (**Scanes et al., 1983**, **Gonzales et al., 1999**), age (**Newcomer, 1978**; **Decuyper and Buyse, 1988**; **Rendon et al., 1994**) and energy intake and dietary composition (**He et al., 2000**).

Blood CORT concentration has been widely used as a measure of environmental stress in broilers (**Mcfarlane and Curtis, 1989** and **Zulkifli et al., 2003**). **Lin et al., (2006)** reported that CORT concentration in broilers that had been exposed to acute heat stress (32°C and 40% relative humidity) for 6 hours varied from 3.17 to 4.28 ng/ml on day 42. It was also reported that heat stress caused the

blood CORT level to increase in cockerels. The results shown in table (10) disagree with those of **Lin et al., (2006)**. This difference might be due that in the current work broiler chickens were exposed to chronic heat stress from the 21st day to the 42nd days of age. So, if serum CORT levels were increased as a result of heat exposure this would have been in the first days after heat exposure but after 1, 2 and 3 weeks (28, 35, 42 days of age) serum CORT levels in the heat stresses groups would have been the same as the control group. This finding is in harmony with that of **Edens, (1978)** who stated that in heat stressed broilers, a sharp increase was followed by a rapid decline in plasma CORT levels as involvement of corticosteroids has been investigated under thermal conditions, but not in chronically heat-exposed chickens (**Geraert et al., 1996**).

Table 1: The experimental design

Group	Name	Number of chicks	Fat level	Temperature
NTLF (control)	Normal Temperature-Low Fat	20	3%	Ideal (22-25 °C)
HTLF	High Temperature-Low Fat	20	3%	Natural temperature of Summer season from 21 day of age till 42 day of age. (25°C -37°C)
HTHF	High Temperature-High Fat	20	6%	Natural temperature of Summer season from 21 day of age till 42 day of age. (25°C -37°C)

Table 2: The Physical and chemical composition of the experimental diet

Ingredient	Starter-grower diet		finisher diet	
	0-4 weeks		4-6 weeks	
	Soy bean oil level %			
	3	6	3	6
Yellow corn	59.54	53.52	69.50	62.54
Soybean meal	22.38	29.98	17.90	25.62
Corn gluten	4.6	--	3.43	--
Soy bean oil	3.00	6.00	3.00	6.00
Fish meal	7.25	7.5	2.90	2.00
Lime stone	1.38	1.01	1.08	1.30
Di Calcium phosphate	1.20	1.59	1.51	1.93
Sodium chloride	0.30	0.30	0.30	0.30
Sodium vit. premix ^a	0.25	0.25	0.25	0.25
Methionine	0.1	0.1	0.03	0.06
Crude protein %	23	22.96	18	18.02
ME (Kcal / kg)	3130	3160	3182	3200
Ether extract %	6.07	9.16	6.02	8.76
Calcium %	1.03	1.00	0.91	1.06
Total phosphorus %	0.71	0.73	0.68	0.77

Table (3): Effect of fat level and high environmental temperature on body weight (gm) of broiler chickens. (Means ± SE)

Age/day	NTLF	HTLF	HTHF
21 st	788.5 ^a ±16.53	787.5 ^a ±13.72	745.5 ^a ±25.64
28 th	1329 ^a ±28.83	1294 ^a ±26.85	1295 ^a ±40.31
35 th	1787 ^a ±40.31	1713 ^a ±39.32	1762 ^a ±57.86
42 th	2103 ^a ±48.41	2051.4 ^a ±47.41	2145.7 ^a ±29.26

Table (4): Effect of fat level and high environmental temperature on body weight gain (gm) of broiler chickens. (Means \pm SE).

Age/day	NTLF	HTLF	HTHF
21 st	412 ^a \pm 16.5	412.5 ^a \pm 7.15	384.5 ^a \pm 15.66
28 th	540.5 ^a \pm 128.46	507 ^a \pm 14.74	550 ^a \pm 16.81
35 th	458 ^a \pm 16.29	418.5 ^a \pm 15	466.6 ^a \pm 21.79
42 th	309.86 ^a \pm 126.2	290.71 ^a \pm 12.97	372 ^a \pm 34.72

Table (5): Effect of fat level and high environmental temperature on feed conversion ratio/gm in broiler chickens. (Means \pm SE)

Age/day	NTLF	HTLF	HTHF
21 st	1.4429 ^a \pm 0.08	1.4432 ^a \pm 0.02	1.4884 ^a \pm 0.06
28 th	1.6361 ^a \pm 0.07	1.8192 ^a \pm 0.05	1.8243 ^a \pm 0.09
35 th	2.1953 ^a \pm 0.07	2.2117 ^a \pm 0.07	2.0411 ^a \pm 0.09
42 th	3.5943 ^a \pm 0.2	3.4112 ^a \pm 0.14	2.8473 ^a \pm 0.33

NTLF (control) (normal temperature-low fat) HTLF (high temperature low fat) HTHF (high temperature high fat)

Table (6): Effect of fat level and environmental temperature on the amount of feed consumed gm/bird/week.

Age/day	NTLF	HTLF	HTHF
21 st	583.25	593.75	563
28 th	865	855	815
35 th	995	915.5	934.5
42 th	1100	980.56	990

NTLF (control) (normal temperature-low fat) HTLF (high temperature low fat) HTHF (high temperature high fat)

Table (7): Effect of fat level and environmental temperature on blood cells count and differential leucocytic count (Means \pm SE).

Parameter	Age/day	NTLF	HTLF	HTHF
RBCs $\times 10^4$	28	2.452 ^{def} ± 0.19	2.486 ^{def} ± 0.27	2.88 ^{cde} ± 0.23
	35	2.762 ^{def} ± 0.22	2.46 ^{def} ± 0.03	2.756 ^{de} ± 0.13
	42	2.998 ^{adc} ± 0.21	1.784 ^f ± 0.16	2.324 ^{ef} ± 0.27
WBCs $\times 10^3$	28	18.470 ^c ± 0.35	15.640 ^{dc} ± 1.6	16.300 ^d ± 0.89
	35	18.520 ^c ± 0.17	13.590 ^e ± 1	18.560 ^e ± 1.4
	42	18.720 ^c ± 0.18	14.480 ^{ef} ± 1.9	16.480 ^d ± 1.4
H%	28	39 ^e ± 0.89	48.8 ^a ± 2.22	49.4 ^a ± 2.37
	35	36.4 ^h ± 1.07	48.8 ^a ± 1.74	46.8 ^e ± 0.91
	42	32.4 ⁱ ± 1.07	41.6 ^k ± 1.2	43.6 ^d ± 0.67
L %	28	58.8 ^c ± 0.96	41.6 ^k ± 2.24	44.4 ^d ± 2.73
	35	62.8 ^d ± 0.66	48.4 ^b ± 1.63	48 ^{gh} ± 2.48
	42	65.4 ⁱ ± 1.66	55.2 ^c ± 1.31	51.2 ^f ± 2.08
H/L ratio	28	0.664 ^{bcd} ± 0.02	1.197 ^a ± 0.12	1.179 ^a ± 0.12
	35	0.58 ^{cd} ± 0.02	1.017 ^{abc} ± 0.06	1.029 ^{abc} ± 0.08
	42	0.498 ^d ± 0.02	0.757 ^{abcd} ± 0.03	0.853 ^{abcd} ± 0.03

Table (8): Effect of fat level and high environmental temperature on serum lipid metabolites (Means \pm SE).

Parameter	Age/day	NTLF	HTLF	HTHF
Tcol (mg/dl)	28	131.718 ^{cdet} ± 2.9	135.917 ^{ab} ± 3.29	136.917 ^a ± 3.61
	35	131.472 ^{def} ± 2.26	133.864 ^{bc} ± 3.43	133.864 ^{bc} ± 3.34
	42	131.865 ^{def} ± 2.26	132.613 ^{bcd} ± 3	132.372 ^{bcd} ± 3.89
TG (mg/dl)	28	86.516 ^{ab} ± 4.07	88.087 ^a ± 4.27	88.965 ^a ± 4.2
	35	85.835 ^{ab} ± 4.72	87.12 ^{ab} ± 4.51	87.589 ^a ± 4
	42	86.604 ^{ah} ± 5.05	86.211 ^{ab} ± 4.87	86.264 ^{ab} ± 4.44
Tot. Lipids (mg/dl)	28	866.568 ^{bc} ± 93.67	890.573 ^a ± 82.28	890.129 ^a ± 81.34
	35	832.155 ^{dc} ± 93.36	862.763 ^{bc} ± 108.3	873.498 ^{ab} ± 76.54
	42	865.342 ^{bc} ± 92.93	875.006 ^{ab} ± 105.4	893.401 ^a ± 88.62
HDL (mg/dl)	28	83.134 ^e ± 5.97	84.11 ^{de} ± 5.77	89.368 ^{abc} ± 5.51
	35	83.56 ^{de} ± 8.18	85.748 ^{bcd} ± 7.19	90.456 ^{ab} ± 3.92
	42	84.39 ^{cd} ± 8.17	93.337 ^a ± 3.62	93.251 ^a ± 3.98
LDL (mg/dl)	28	31.282 ^{abcd} ± 5.73	34.314 ^a ± 4.86	29.081 ^{abcd} ± 3.35
	35	31.655 ^{abcd} ± 6.23	30.411 ^{abcd} ± 4.39	25.890 ^{cde} ± 1.55
	42	26.633 ^{bcd} ± 6.23	22.034 ^e ± 1.69	21.868 ^e ± 1.76

- Tcol= Total cholesterol, TG= Triacylglycerol, Tot.Lipid= Total lipids

- Means carrying different small letters in the same row are significantly different at the level of $p < 0.05$.

- NTLF (control) (normal temperature-low fat) HTLF (high temperature low fat) HTHF (high temperature high fat)

- Means carrying different small letters in the same row are significantly different at the level of $p < 0.05$

- NTLF (control) (normal temperature-low fat) HTLF (high temperature low fat) HTHF (high temperature high fat).

Table (9): Effect of fat level and high environmental temperature on serum total protein, albumin and globulin (Means \pm SE).

Parameter	Age/day	NTLF	HTLF	HTHF
Protein (g/L)	28	2.804 ^{abc} \pm 0.3	2.52 ^{bc} \pm 0.28	2.531 ^{bc} \pm 0.23
	35	3.148 ^{ab} \pm 0.31	2.516 ^{bc} \pm 0.19	2.704 ^{bc} \pm 0.22
	42	3.006 ^{abc} \pm 0.11	2.688 ^{bc} \pm 0.17	2.705 ^{bc} \pm 0.11
Albumin (g/L)	28	1.52 ^a \pm 0.17	1.27 ^a \pm 0.06	1.215 ^a \pm 0.06
	35	1.698 ^a \pm 0.15	1.329 ^a \pm 0.14	1.293 ^a \pm 0.11
	42	1.645 ^a \pm 0.17	1.375 ^a \pm 0.14	1.386 ^a \pm 0.12
Globulin (g/L)	28	1.285 ^a \pm 0.2	1.226 ^a \pm 0.3	1.316 ^a \pm 0.25
	35	1.408 ^a \pm 0.22	1.187 ^a \pm 0.19	1.411 ^a \pm 0.18
	42	1.361 ^a \pm 0.08	1.313 ^a \pm 0.14	1.316 ^a \pm 0.12

- Means carrying different small letters in the same row are significantly different at the level of $p < 0.05$

- NTLF (control) (normal temperature-low fat) HTLF (high temperature low fat) HTHF (high temperature high fat)

Table (10): Effect of fat level and environmental temperature on serum hormones (Means \pm SE).

Parameter	Age/day	NTLF	HTLF	HTHF
T3 ng/ml	28	1.068 ^{abc} \pm 0.07	0.838 ^c \pm 0.02	0.832 ^c \pm 0.02
	35	1.128 ^{abc} \pm 0.09	1.006 ^{abc} \pm 0.06	0.892 ^{bc} \pm 0.03
	42	1.134 ^{abc} \pm 0.08	1.11 ^{abc} \pm 0.08	1.092 ^{abc} \pm 0.11
T4 ng/ml	28	5.934 ^a \pm 0.87	5.99 ^a \pm 0.98	6.06 ^a \pm 0.89
	35	5.906 ^a \pm 1.25	6 ^a \pm 0.82	5.716 ^a \pm 1.08
	42	5.940 ^a \pm 1.05	5.88 ^a \pm 0.84	5.886 ^a \pm 0.84
CORT ng/ml	28	1.508 ^a \pm 0.15	2.066 ^a \pm 0.35	2.132 ^a \pm 0.23
	35	1.534 ^a \pm 0.16	1.892 ^a \pm 0.38	1.918 ^a \pm 0.2
	42	1.63 ^a \pm 0.08	1.752 ^a \pm 0.2	1.756 ^a \pm 0.3

Means carrying different small letters in the same row are significantly different at the level of $p < 0.05$

NTLF(control) (normal temperature-low fat) HTLF (high temperature low fat) HTHF (high temperature high fat)

REFERENCES

- Abdel-Fattah, S. A. (2006)** : Physiological and immunological adjustment of dietary ascorbic acid and acetyl salicylic acid in heat stressed Japanese quail. *Egypt. Poult. Sci*; 26: 1395 - 1418.
- Abd-Elsamee, M. O. (2005)** : Influence of different levels of lysine and vitamin (E) on broiler performance under normal or heat stress condition. *Egyptian. J. Nutr. feeds*; 8: 827 - 838 .
- Abu-Dieyeh, Z. H. M. (2006a)** : Effect of high temperature per se on growth performance of broilers. *Int. J. poult. Sci*; 5:19 - 21.
- Abu-Dieyeh, Z. H. M. (2006b)** : Effect of chronic heat stress an long-term feed restriction on broiler performance. *Int. J. poult. Sci*; 5: 185-190.
- Allain, C.; Poon, L. and chan, C. (1974)** : Enzymatic determination of serum total cholesterol . *clin. chem*; 20:470.
- Alao, S. J. and Balnave, D. (1985)** : Nutritional significance of different fat sources for growing broilers. *Poult. Sci*; 64: 1602-1604.
- Albertini, A. and Eknis, R. P. (1982)** : Determination of serum total thyroxin (T4) using (ELISA) kit. Free hormone in blood - Amsterdam. Elsevier Biochemical Press.
- Al-Murrani, W. K. A.; Al-Sam, H. Z. and Al-Athari, A. M. (1997)** : Heterophile/lymphocyte ratio as a selection criterion for heat resistance in domestic fowl. *Br. Poult.Sci*; 38: 159-163.
- Bartha, T. (1993)** : Thyroid hormone metabolism in broiler chickens as influenced by exogenous and endogenous factors. *Dissert. Agr. 233*, Katholieke Universiteit Leuven, Belgium.
- Berrong, S. L. and Washburn, K. W. (1998)** : Effects of genetic variation on total plasma protein, body weight gains and body temperature responses to heat stress. *poult. sci*; 77: 379-385.
- Bonnet, S.; Geraert, P. A.; Lessire, M.; Carre, B. and guillaumin, S. (1997)** : Effect of high ambient temperature on feed digestibility in broilers. *Poult. Sci*; 76: 857 - 863.
- Borges, S. A.; Fischer da Silva, A.V.; Majorjorka, A.; Hooge, D. M. and Cummings, K. R. (2004)** : Physiological responses of broiler chickens to heat stress and electrolyte balance (sodium plus potassium minus chloride, milliequivalents per kilogram) 1,2. *Poult. Sci*; 83 1551 1558.
- Buyse, J.; Darass, V. M.; Vleurick, L.; Kühn, E. R. and Decuyper, E. (2001)** : Nutritional regulation of the somatotropic axis and intermediary metabolism in the chicken. *Avian. Endocrinol*; 303-313.
- Carmen, A.; France, M.; Macleod, G. and Julie, E. (1991)** : Alleviation of acute heat stress by food withdrawal or darkness. *Br. Poult. Sci*; 32: 219-225.
- Decuyper, E. and Buyse, J. (1988)** : Thyroid hormones, corticosterone, growth hormone and somatomedines in avian species: General effects and possible implications

in fattening. Pag 298-302 in *Leanness in Domestic Birds: Genetic, Metabolic and Hormonal Aspects*. B. Leclercq, and C.C. Whitehead, ed. Butterworths, London, UK.

Decuyper, E., Kühn, E. R. (1984) : Effect of fasting and feeding time on circadian rhythms of serum thyroid hormone concentrations, glucose, liver monodeiodinase activity and rectal temperature in growing chickens. *Domest. Anim. Endocrinol*; 1: 251 - 262.

Deyhim, F. and Teeter, R. G. (1991) : Sodium and potassium chloride drinking water supplementation effects on acid-base balance and plasma corticosterone in broilers reared in broilers in thermoneutral and heat-distressed environment. *poult. Sci*; 70: 2551 - 2553.

Donia, V. (1987) : Determination of serum cortisol using (ELISA) kit. *Clinical Chemistry*; 33:33-42.

Donkoh, A. (1989) : Ambient temperature: a factor affecting performance and physiology response of broiler chickens. *Int.J. Biomet*; 33: 259-265.

Doumas, B., Weston, W., and Biggs, H. (1971): Albumin standards and the measurements of serum albumin with bromocresol green. *clin. Chim. Acta*; 31 (1): 87 - 96.

Edens, F. W. (1978) : Adrenal cortical insufficiency in young chickens exposed to a high ambient temperature. *Poult. Sci*; 75: 1746 - 1750.

El-Katcha, M. I. and Walaa Abd El-Razik

M. (1996) : Effect of feeding over and unheated oil with or without vitamin E supplementation on growth and reproductive performance of Japanese quail. *Alex. J Vet Sci*;12(2): 145-158.

Emery, J. (2004) : Heat stress on poultry solving the problem. Defra publications (ADAS). <http://www.alpha.com/ahd>.

Faisal, B. A.; Abdel - Fattah, S. A. K.; El-Hommosany, M.; Nermin M-Abdel - Gawad, and Male, F. M. Ali (2008) : Immunocompetence, hepatic heat shock protein 70 and physiological responses to feed restriction and heat stress in two body weight lines of Japanese quail. *Int. J. poult. sci*; 7, 2: 179-183.

Faitas, A. A.; Abd-Moty, A. K. I. and Mohamed, Kh. A. (1987) : The effect of exposure at 40 °C on the physiological response of the domestic fowl rearing at 16 °C. *Minia. J. Agric. Res and Dev*; 9(1):383-389.

Fassati, P. and prencipe L. (1982) : Enzymatic determination of serum triglycerides. *clin. chem.* ; 28, 2077.

Feldman, B. F.; Zinkl, J. G. and Jain, I. C. (2000) : *Schalm's veterinary haematology* 5th ed Lippincott Williams and Wilkins, Canada.

Ferket, P. R. (1995) : Nutrition of turkeys during hot weather. In *Proceedings of Turkey 18th Technical Turkey Conference* Renfrew, U.K. Turkey, Leyburn, UK.

Friedewald, W.; Levy, R. and Frederickson, D. (1972) : Estimation of the concentra-

tion of low density lipoprotein cholesterol in plasma without use of the ultracentrifuge. Clin. chem.; 18:499-502.

Geraert, P. A.; Padilha, J. C. F. and Guillaume, S. (1998) : Metabolic and endocrine changes induced by chronic heat exposure in broiler chickens: Biological and endocrinological variables. Br.J. Nutr; 75: 205-216.

Ghazalah, A. A.; Soliman, A. Z. M.; Bouious, N. Z.; Samla and Mobarez, M. (2007): Effect of using dry fat on performance, nutrients digestibility, carcass traits and blood constituents of broiler chicks. Egypt. Poult. Sci; 27: 363-382.

Gonzales, E.; Buyse, J.; Sartori, J. R.; Lodd, M. M. and Decuyper, E. (1999) : Metabolic disturbances in male broilers of different strains.2. Relationship between the thyroid and somatotropic axes with growth rate and mortality. Poult.Sci; 78:516-521.

Gornal, A. C.; Bardawill, C. J. and David M. M. (1949) : Determination of serum total protein content in serum. Biol. J. chem; 177-751.

Gross, W. B. and Siegel, H. S. (1983): Evaluation of the heterophile/Lymphocyte ratio as a measure of stress in chickens. Av. Diseases; 27:972-979.

Gursu, M. F.; Onderel, M.; Gulcu, F. and Sabir, K. (2004) : Effects of vitamin C and folic acid supplementation on serum paraoxonase activity and metabolites induced by heat stress in vivo . Nutr, Res; 24: 157 - 164. coronary heart disease. American. J. Clin. Nutr;

45: 1168 - 1175.

Har, L.; Rong, D. and Zhang, Z. A. (2000) ; effect of thermal environment on the digestion of broilers. Amin. Physiol; 83: 61-75.

He, J.; Ohtsuka, A. and Hayashi, K. (2000) ; Selenium influences growth via thyroid hormone status in broiler chickens. Br. J. Nutr; 84: 727-732.

Kahl, S.; Rosebrough, R. W. and Elsas-ser, T. H. (1998) : Hepatic iodothyronine 5-Monodetodnase activity in the broiler chicken: Effect of dietary fat and triiodothyronine supplementation. Nutr. Res; 18: 1043 - 1047.

Kinsella, J. E.; Lokesh, B. and Stone, R. (1990) : Dietary n-3 polyunsaturated fatty acids and amelioration of cardiovascular disease: possible mechanisms. American. J. Clin. Nutr; 52: 1 - 28.

Klee, G. G. (1996) : Clinical usage recommendation and analytic performance goals for total and free triiodothyronine measurements. Clinical Chemistry;42: 155-159.

Kutlu, H. R. and Forbes, J. M. (1993) : Changes in growth and blood parameters in heat-stressed broiler chicks in response to dietary ascorbic acid. Livestock produce Sci; 36:335-350.

Lin, H.; Decuyper, E. and Buyse, J. (2008) : Acute heat stress induces oxidative stress in broiler chickens. Comp. Biochem. Physiol. A Mol.Integr.Physiol; 144: 11-17.

Lopez-Ferrer, S., Baucells, M.D., Barro-

ta, A. C. and Grashorn, M. A. (1999) : Influence of vegetable oil sources on quality parameters of broiler meat. *Archiv fur Geflugelkunde*; 63: 29-35.

Lopez-Virella, M.; stome, P. and ElHo, S. (1977) : Determination of HDL in serum. *clin. chem*; 23: 822.

Makind, M. O. and fatunmbi, O. O. (1985) : Some hematological and biological values of turkeys in Ibadan. *Bulletin of animal health and production in Africa*; 33: 245 - 248.

Mansour, M. E. (2006) : Effect of fats in poultry diet. M. V. Sc. Thesis. Fac. vet. Mansoura University Egypt.

Mashaly, M. M.; Hendricks, G. L.; Kalam, M. A.; Gehad, A. E.; Abbas, A. O. and patterson, P. H. (2004) : effect of heat stress on production parameters and immune responses of commercial laying hens. 1. *Poult. Sci*; 83: 889-894 .

Mateos, G. C.; Sell, J. L. and Eastwood, J. A. (1982): Rate of food passage (transit time) as influenced by level of supplemental fat. *Poult. Sci*; 63: 1170 - 1174.

McFarlane, J. M. and Curtis, S. E. (1989) : Multiple concurrent stressors in chicks.3. Effects on plasma Corticosterone and the heterophile/lymphocyte ratio. *Poult. Sci*; 68: 522-527.

Mills, L. J.; Mitchell, M. A. and Mahon, M. (1998) : comparison of thermoregulatory ability in fast and slow growing strain of tur-

key during acute heat stress. *Br. Poult. Sci*; 40: 51-2.

Newcomer, W. S. (1978) : Diurnal rhythms of thyroid function in chicks. *Gen. comp. endocrinol*; 24: 65-73.

Nitsan, Z.; Dvorin, A.; Zoref, Z. and Mokady, S. (1997) : Effect of added soybean oil and dietary energy on metabolizable and net energy of broiler diets. *Br. poult. Sci*; 38: 101 - 106.

NRC (1994) : Nutrient requirements of poultry. National Academy Press. Washington. DC. USA.

Ozbey, O. and Ozcelik, M. (2004) : The effect of high environmental temperature on growth performance of Japanese quails with different body weight. *Int. J. poult. Sci*; 3: 468 - 470.

Ozdogan, M. and Aksit, A. (2003) : Effects of feeds containing different fats on carcass and blood parameters of broilers *J. Appl. pouth. Res*; 12: 251 - 256.

Pinchasov, Y. and Nir, I. (1992) : Effect of dietary polyunsaturated fatty acid concentration on performance, fat deposition and carcass fatty acid composition in broiler chickens. *Poult. Sci*; 71: 1504-1512.

Puvadolpirod, S. and Thaxton, J. P. (2000) : Model of physical stress in chickens 2. Dosimetry of adrenocorticotropin. *Poult. Sci*; 79: 370 - 376.

Rautela, M. A.; Khatl, D. S. and Joshp,

M. C. (1994) : Effect of high temperature on some haematological values of gallus domesticus at 1650 m altitude. *Indian J. Anim. Sci*: 64: 752 - 753.

Rendon, J.A., Lien, R.J., Oates, S.S. and Bilgili, S.F. (1994): Plasma concentration of corticosterone and thyroid hormones in broilers provided various lighting schedules. *Poult. Sci*: 73:186-193.

Rosebrough, R. W.; Mcmurty, J. P. and Vasilatos-Younken, R. (1999) : Dietary fat and protein Interactions in the broilers. *poult. Sci*: 78: 992 - 998.

Sahin, N. V.; Sahin, K. and Kucuk O. (2001) : Effects of vitamin E and Vitamin A supplementation on performance, thyroid status and serum concentrations of some metabolites and minerals in broilers reared under heat stress (32°C). *Vet. Med- Czech*, 46, (11-12):286-292.

Sahota, A. W.; Gillani, A. H. and Ullah, M. F. (1994) : Hematological studies on heat stress chickens supplemented with ascorbic acid. *Pakistan. Vet. J*:14: 30-33

Sanz, M.; Florves, A. and Lopez, C. L. (1999) : Effect of fatty acid saturation in broiler diets on abdominal fat and breast muscle fatty acid composition and susceptibility to lipid oxidation. *Poult. Sci*: 64 : 1602 - 1604.

Scanes, C. G.; Marsh, J.; Decuypere, E. and Rudas, P. (1983) : Abnormalities in the plasma concentrations of thyroxine, triiodothyronine and growth hormone in sex-linked

dwarf and autosomal dwarf white leghorn domestic fowl. *J. Endocrinol*: 97 : 127-135.

Shum, Y. S. and Griminger, P. (1972) : Prothrombin time stability of avian and mammalian plasma: effect of anticoagulants. *Lab. Anim. Sci*: 22: 384.

Siegel, H. S. (1968) : Blood cells and chemistry of young chickens during daily ACTH and cortisol administration. *Poult. Sci*: 47 : 1811 - 1817.

Sinurate, A. P.; Balnave, D. and McDowell, G. H. (1987) : Growth performance and concentrations of thyroid hormones and growth hormone in plasma of broilers at high temperatures. *Aust. J. Biol. Sci*: 40:443-450.

Sosnowka-Czajka, E.; Skomorucha, I. and Herbut, E. (2005) : Effect of dietary vitamin supplementation on productivity and physiological parameters of broiler chickens exposed to elevated ambient temperature Warsaw, Poland; 2 : 111 - 114.

SPSSPC+ (1994) : SPSS for window release 10.0 SPSS Inc., USA.

Sturkie, P. D. (1986) : Pages 235 - 239 in 'Avian physiology'. 4th ed. Springer -Verlag, Inc., NewWork , NY.

Türkyilmaz, M. K. (2008) : The effect of stocking density on stress reaction in broiler chicks during summer. *Turk. J. Vet. Anim. Sci*: 32 (1): 31 - 36.

Wafaa, Eleraky, A. and El-Sadawy, H. A. (1997) : Effect of feeding high energy diets containing supplemental fat on growth performance, some blood parameters yield and body composition of broiler chicks. Egypt. Poult. Sci; 17(1): 151-169.

Whitehead, C. C. and Keller, T. (2003) : An update on ascorbic acid in poultry. J. World's poult. Sci; 59: 161 - 184.

Zollitsch, W.; Knaus, W.; Aichinger, F. and Lettner, F. (1997) : Effect of different dietary fat sources on performance and carcass characteristics of

broilers. Anim. Feed Sci. Technol; 66 : 63-73.

Zollner, N. and Kirsch, K. (1982) : Determination of the total lipid content in serum. zentralblattfür Gesamte experimental medicine · 135 : 545 - 549.

Zulkifli, I.; Liew, P. K.; Israf, D. A.; Omar, A. R. and Hair-Bejo, M. (2003) : effect of early age feed restriction and heat conditioning on heterophile/lymphocyte ratios, heat shock protein 70 expression and body temperature of heat stressed broiler chickens. J. Therm. Biol; 28:217-22.

الملخص العربي

تأثيرات العليقة عالية الدهون على بعض السمات الفسيولوجية في بدارى التسمين تحت درجات الحرارة العالية

د/ فهيم البهى وحيش*
د/ أسامه أحمد أبوإسماعيل**
د/ أماني محمود عجيز*

قسم الفسيولوجيا - كلية الطب البيطرى - جامعة المنصورة - مصر*

قسم الرعاية وتنمية الثروة الحيوانية - كلية الطب البيطرى - جامعة المنصورة - مصر**

أجريت الدراسة الحالية لاستبيان تأثير زيادة مستوى زيت الصويا فى علائق بدارى التسمين على بعض السمات الفسيولوجية لهذه الطيور تحت درجات الحرارة العالية فى فصل الصيف وقد تم استخدام ستين كتكوت عمر يوم من سلالة (كوب) وتقسيمهم عشوائياً إلى ثلاث مجموعات متساوية كل مجموعة تحتوى على عشرين كتكوتاً، المجموعة الضابطة (درجة حرارة عادية ومستوى دهن منخفض) والتي تم تربيتها (درجة حرارة عالية ومستوى دهن منخفض) والتي تم تربيتها تحت درجة حرارة عالية فى فصل الصيف وتم تغذيتها على عليقة تحتوى على ٣٪ زيت صويا وذلك من عمر ٢١ حتى ٤٢ يوم والمجموعة الثالثة (درجة حرارة عالية ومستوى دهن عالى) والتي تم تربيتها تحت درجة حرارة عالية فى فصل الصيف وتم زيادة نسبة زيت الصويا فى العليقة من ٣٪ إلى ٦٪ وذلك من عمر ٢١ حتى ٤٢ يوم، تم حساب متوسط الأوزان وكمية العليقة المستهلكة إسبوعياً وأيضاً تسجيل معدل الزيادة وكفاءة التحريل لكل مجموعة وتم تجميع عينات الدم من أوردة الجناح عن عمر ٢٨ و ٣٥ و ٤٢ يوم وتم تقسيم العينات لعينات دم وعليها موانع تجلط لعد خلايا الدم الحمراء، وخلايا الدم البيضاء، وعمل مسحات لتسجيل كلاً من إعداد الهيتروفيل والليمفوسيت ونسبة الهيتروفيل إلى الليمفوسيت (H/L Ratio) كما تم فصل مصل الدم لقياس مستويات الكوليمترول الكلى والجليسريدات الثلاثية والدهون الكلية والبروتينات الدهنية (الليپوروتينات) عالية وقليلة الكثافة والبروتينات الكلية والألبومينات والجلوبولينات والترأى أيدوثيرونين (T3) والثيروكسين (T4) والكورتيكوستيرون (CORT) وكانت النتائج كالتالى :

- معدلات النمو : لوحظ عدم وجود فرق معنوية بين المجموعات فى وزن الجسم، معدل الزيادة فى وزن الجسم، كفاءة التحريل الغذائى وكمية العليقة المستهلكة.

- مكونات الدم : كان هناك إنخفاض معنوى فى متوسط عدد كرات الدم الحمراء، فى عينة الدم فى المجموعة عالية الحرارة قليلة الدهون عند عمر ٤٢ يوم مقارنة بالمجموعة عالية الحرارة عالية الدهون والمجموعة الضابطة عند نفس العمر، بالنسبة لمتوسط عدد خلايا الدم البيضاء، فى الدم كان هناك انخفاض معنوى فى المجموعة عالية الحرارة قليلة الدهون عن عمر ٣٥، ٤٢ يوم والمجموعة عالية الحرارة عالية الدهون عند

عمر ٤٢ يوم مقارنة بالمجموعة الضابطة عن عمر ٢٨، ٣٥، ٤٢ يوم، بالنسبة لتوسط نسبة الهيمتروفيل إلى الليمفوسيت كان هناك زيادة معنوية في المجموعة عالية الحرارة منخفضة الدهون والمجموعة عالية الدهون عند عمر ٢٨ يوم وكان هناك انخفاض معنوي في المجموعة الضابطة عند عمر ٤٢ يوم.

- نواتج أبيض الدهون في الدم : الصيف كان متوسط الكوليسترول في مصل الدم لبدارى التسمين المجموعة عالية الحرارة عالية الدهون عند عمر ٢٨ يوم أعلى معنوياً منه في المجموعة الضابطة والمجموعة عالية الحرارة منخفضة الدهون عند عمر ٢٨ يوم وفى المجموعة عالية الحرارة عالية الدهون عند عمر ٢٨ و ٣٥ يوم مقارنة بالمجموعة الضابطة.

بالنسبة لمستوى الدهون الكلية في مصل الدم كان هناك زيادة معنوية في المجموعة عالية الحرارة منخفضة الدهون عند عمر ٢٨ يوم وفى المجموعة عالية الحرارة عالية الدهون عن عمر ٢٨ و ٣٥ يوم مقارنة بالمجموعة الضابطة عند عمر ٢٨ و ٣٥ و ٤٢ يوم.

كانت متوسطات الليبوبروتينات عالية الكثافة في مصل الدم لبدارى التسمين أعلى معنوياً في المجموعة عالية الحرارة قليلة الدهون والمجموعة عالية الحرارة عالية الدهون عند عمر ٤٢ يوم وكانت متوسطات المجموعة الضابطة أقل معنوياً من المجاميع الأخرى عند عمر ٢٨، ٣٥، ٤٢ يوم، بالنسبة لليبوبروتينات منخفضة الكثافة كانت المتوسطات منخفضة معنوياً في المجموعة عالية الحرارة قليلة الدهون والمجموعة عالية الحرارة عالية الدهون عند عمر ٤٢ يوم مقارنة بالمجموعة الضابطة.

البروتينات الكلية والالبومين والجلوبولين : كان هناك انخفاض معنوي في قيمة البروتينات الكلية في المجموعة عالية الحرارة قليلة الدهون والمجموعة عالية الحرارة عالية الدهون عند عمر ٢٨، ٣٥، ٤٢ يوم مقارنة بالمجموعة الضابطة عند نفس الأعمار، لم تكن هناك فروق معنوية في متوسطات الألبومين والجلوبولين في مصل الدم بين المجموعات.

- الهرمونات :

(تراى أيودوثيرونين (T3)، الثيروكسين (T4) والكورتيكوستيرون (CORT) كان متوسط هرمون (T3) في مصل الدم أقل معنوياً في المجموعة عالية الحرارة قليلة الدهون والمجموعة عالية الحرارة عالية الدهون عند عمر ٢٨ يوم مقارنة بالمجموعة الضابطة عند نفس العمر وعند عمر ٣٥ و ٤٢ يوم لوحظ عدم وجود فروق معنوية بين المجاميع، بالنسبة لتوسط هرمون (T4) في السيروم لم تكن هناك فروق معنوية عند عمر ٢٨، ٣٥ و ٤٢ يوم بين المجاميع، بالنسبة لتوسط هرمون (CORT) في مصل الدم كانت هناك زيادة غير معنوية عند عمر ٢٨ يوم في كلاً من المجموعة عالية الحرارة قليلة الدهون والمجموعة عالية الحرارة عالية الدهون مقارنة بالمجموعة الضابطة وعند عمر ٣٥ و ٤٢ يوم لوحظ عدم وجود فروق معنوية بين المجموعات.